

# Some new results on the Chowgroups of quadrics

Markus Rost

Regensburg, January 1990

## § 1 Introduction

The computation of some  $K$ -cohomology groups of certain normvarieties plays an essential role in the investigation of the bijectivity of the Galois symbol

$$K_n^M F/p \rightarrow H^n(F; \mu_p^{\otimes n}).$$

In the case  $p = 2$  these normvarieties are quadrics associated with Pfisterforms.

In this note we describe some new results and conjectures about such quadrics. In short we have the following results:

It turns out that for a  $n$ -fold Pfisterform  $\varphi$  the Chow-motive of its associated quadric  $X_\varphi$  can be described as

$$X_\varphi \simeq M_\varphi \times \mathbb{P}^{d_\varphi}, \quad d_\varphi = 2^{n-1} - 1,$$

where  $M_\varphi$  is a certain Chow-motive associated with  $\varphi$ . We give a complete description of the Chowgroups of  $M_\varphi$  and of the generators of  $H^p(M_\varphi; K_{p+1})$ ; moreover we have a precise conjecture about (the Milnor- $K$ -theory version of)  $H^p(M_\varphi; K_{p+r})$  for  $r \leq 2$ .

This note contains no proofs. Detailed proofs will be prepared as soon as possible.

## § 2 Motivic decomposition of certain quadrics

We work in the category of Chow-motives over a field  $F$  ( $\text{Char } F \neq 2$ ) (see e.g. [Fulton; Intersection Theory, § 16]). For a motive  $M = (X, p)$  let  $CH_k(M) = p_*(CH_k(X))$ , where  $CH_k(X)$  is the group of  $k$ -dimensional cycles modulo rational equivalence. We denote by  $L = (\mathbb{P}^1, p)$  the Tate motive and by  $L^i$  its  $i$ -th power. For a quadratic form  $\varphi$  over  $F$  we denote by  $X_\varphi$  the corresponding projective quadric ( $\dim X_\varphi = \dim \varphi - 2$ ) and by  $\overline{X}_\varphi$  its associated Chow-motive.

One first observation is



The motive  $M_\varphi$  can be described more precisely as follows: Let  $\varphi = \mu \otimes \langle\langle a \rangle\rangle$  where  $\mu$  is a Pfisterform of degree  $n_\mu = n_\varphi - 1$  and put  $\rho = \mu \perp \langle -a \rangle$ . Then  $M_\varphi = (X_\rho, p)$  for a certain projector  $p \in \text{End}(\overline{X}_\rho)$  such that  $p_*$  is the identity on  $CH_{d_\varphi}(X_\rho)$  and  $CH_0(X_\rho)$  (note that  $d_\varphi = \dim X_\rho$ ). Hence  $CH_{d_\varphi}(M_\varphi) = \mathbb{Z}$  with the cycle  $X_\rho$  as canonical generator; we denote this generator by  $[M_\varphi]$ . Moreover  $CH_0(M_\varphi) = CH_0(X_\rho)$ ; the homomorphisms  $N$  in iii) are given by the degree of a zero-cycle.

The motive of  $X_\rho$  decomposes as

$$\overline{X}_\rho = M_\varphi \oplus \overline{X}_{\mu'} \otimes L,$$

where  $\mu'$  is the pure subform of  $\mu$ , i.e.  $\mu = \mu' \perp \langle 1 \rangle$ . More generally, one has

#### Proposition 4

Let  $\varphi$  be a Pfisterform and let  $\rho$  be a subform of  $\varphi$  with  $\dim \rho = \frac{1}{2} \dim \varphi + k$ ,  $k > 0$ . Suppose that

$$\rho_{F(X_\varphi)} \simeq \eta_{F(X_\varphi)} \perp h,$$

where  $\eta$  is a form over  $F$  and  $h$  is hyperbolic of dimension  $2k$ . Then

$$\overline{X}_\rho = \bigoplus_{i=0}^{k-1} M_\varphi \otimes L^i \oplus \overline{X}_\eta \otimes L^k.$$

Hence, if  $\rho$  is an excellent form, i.e. the class of  $\varphi$  in the Witt ring of  $F$  is the alternating sum of a sequence  $\varphi_0, \dots, \varphi_n$  of Pfisterforms  $\varphi_i$  with  $\varphi_i$  a subform of  $\varphi_{i+1}$ , one has a decomposition of the motive  $X_\rho$  in terms of the motives  $M_{\varphi_i}$ .

### § 3 On the $K$ -cohomology of $M_\varphi$

For a variety  $X$  over  $F$  we denote by  $A_p(X, K_n^M)$  the homology of the complex

$$\bigoplus_{v \in X_{(p+1)}} K_{n+p+1}^M \kappa(v) \xrightarrow{d} \bigoplus_{v \in X_{(p)}} K_{n+p}^M \kappa(v) \xrightarrow{d} \bigoplus_{v \in X_{(p-1)}} K_{n+p-1}^M \kappa(v),$$

where  $K_n^M$  denotes the  $n$ -th Milnor  $K$ -group,  $X_{(p)}$  denotes the set of all points of  $X$  of dimension  $p$  and  $d$  is given by the tame symbol.

Using the deformation to the normal cone (see Fulton) one can define intersection theory for the groups  $A_p(X; K_n^M)$ . Hence the functors  $A_p(\ , K_n^M)$  are defined on the category of Chow-motives.

Note that  $A_p(X; K_{-p}^M) = CH_p(X)$  and that for a smooth variety  $X$  of dimension  $d$  there is a natural homomorphism

$$A_p(X; K_n^M) \rightarrow H^{d-p}(X; K_{n+d})$$

induced by the canonical map from Milnor- $K$ -theory to Quillen's  $K$ -groups (which is an isomorphism for  $n + p \leq 2$ ).

For a nonsingular quadratic form  $\varphi$  let  $D_0(\varphi) \subset K_0F = \mathbb{Z}$  be the subgroup

$$D_0(\varphi) = \begin{cases} K_0F & \text{if } \varphi \text{ is isotropic} \\ 2K_0F & \text{if } \varphi \text{ is nonisotropic} \end{cases}$$

and for  $n \geq 1$  let  $D_n(\varphi) \subset K_n^M F$  be the subgroup generated by symbols in which one entry is represented by  $\varphi$ .

One can show that  $D_n(\varphi)$  is exactly the image of the normmap

$$N : A_0(X_\varphi K_n^M) \rightarrow K_n^M F.$$

### Theorem 5

For a Pfisterform  $\varphi$  of degree  $n_\varphi \geq 2$  one has

$$A_p(M_\varphi; K_{-p}) = CH_p(M_\varphi) = \begin{cases} K_0F & \text{for } p = d_\varphi = 2^{n_\varphi - 1} - 1 \\ K_0F/D_0(\varphi) & \text{for } p = 2^k - 1; k = 1, \dots, n_\varphi - 2 \\ D_0(\varphi) & \text{for } p = 0 \\ 0 & \text{else} \end{cases}$$

The generators of  $CH_p(M_\varphi)$  can be described as follows.

For a Pfistersubform  $\psi$  of  $\varphi$  there is a natural morphism

$$i_{\psi, \varphi} : M_\psi \rightarrow M_\varphi,$$

compatible with the norm maps  $CH_0 \rightarrow \mathbb{Z}_0$ . It is induced by inclusion  $X_{\tilde{\rho}} \rightarrow X_\rho$  for appropriate choices of representations  $M_\varphi = (X_\rho, p)$ ,  $M_\psi = (X_{\tilde{\rho}}, \tilde{p})$  as described above.

The generator of  $CH_p(M_\varphi)$ ,  $p = 2^k - 1$  for some  $k \in \{1, \dots, n_\varphi - 1\}$ , is then given by the image  $(i_{\psi, \varphi})_*([M_\psi])$  of the fundamental cycle  $[M_\psi] \in CH_{d_\psi}(M_\psi)$ , where  $\psi$  is any Pfistersubform of  $\varphi$  of degree  $n_\psi = k + 1$ .

### Theorem 6

Let  $\varphi$  be a Pfisterform of degree  $n_\varphi \geq 2$ .

- i) If  $p = 2^k + 2^\ell - 1$ ,  $0 < \ell < k < n_\varphi - 1$ , then  $A_p(M_\varphi; K_{1-p})$  is cyclic of order at most 2.
- ii) If  $p = 0$ , then the normmap  $A_0(M_\varphi; K_1) \rightarrow K_1F$  is injective with image  $D_1(\varphi)$ .
- iii) For all other values of  $p$  the multiplication map

$$CH_p(M_\varphi) \otimes K_1F \rightarrow A_p(M_\varphi; K_{1-p})$$

is surjective.

To give some information about the generators for the groups in i) we describe now a general conjecture about certain elements in the groups  $A_p(M_\varphi; K_n)$ .

Let  $\ell : \mathbb{N} \rightarrow \mathbb{N}$  be the function with the property

$$p + 1 = \sum_{i=0}^{\ell(p)} 2^{k_i} \quad \text{for some } 0 \leq k_0 < k_1 < \dots < k_{\ell(p)}$$

### Conjecture 7

For Pfisterforms  $\varphi$  of degree  $n_\varphi \geq 2$  there exist unique classes

$$\gamma_p(\varphi) \in A_p(M_\varphi; K_{\ell(p)-p}^M) \quad \text{for } 0 < p \leq d_\varphi, p \text{ odd,}$$

such that

I)  $\gamma_{d_\varphi}(\varphi) \in CH_{d_\varphi}(M_\varphi)$  is the generator  $[M_\varphi]$ .

II) For a Pfister-subform  $\psi < \varphi$  one has

$$(i_{\psi, \varphi})_*(\gamma_p(\psi)) = \gamma_p(\varphi) \quad \text{for } 0 < p \leq d_\psi, p \text{ odd.}$$

III) For a Pfister-subform  $\psi < \varphi$  with  $\varphi = \psi \otimes \langle\langle a \rangle\rangle$  one has

$$(i_{\psi, \varphi})^*(\gamma_p(\varphi)) = \{a\} \cdot \gamma_{p-d_\varphi+d_\psi}(\psi) \quad \text{for } d_\psi < p < d_\varphi, p \text{ odd.}$$

For III) note that  $\ell(p - d_\varphi + d_\psi) = \ell(p) - 1$ .

For  $\ell(p) = 0$  the classes  $\gamma_p(\varphi)$  are exactly the generators of  $CH_p(M_\varphi)$  as described above (this follows from I) and II)). I have constructed the classes  $\gamma_p(\varphi)$  for  $\ell(p) \leq 1$  (which are the generators for  $A_p(M_\varphi; K_{1-p})$ ) and it is probable that I can do this for  $\ell(p) \leq 2$ . Moreover our methods should lead to a proof of

### Conjecture 8

For a Pfisterform  $\varphi$  of degree  $n_\varphi \geq 2$  one has for  $n \leq 2$ :

$$A_p(M_\varphi; K_{n-p}^M) = \begin{cases} K_n^M F & \text{for } p = d_\varphi \\ [K_{n-\ell(p)}^M F / D_{n-\ell(p)}(\varphi)] & \text{for } 0 < p < d_\varphi, p \text{ odd,} \\ 0 & \text{for } 0 < p < d_\varphi, p \text{ even} \\ D_n(\varphi) & p = 0 \end{cases}$$

Here we understand  $K_r^M F = 0$  for  $r < 0$ . For  $0 < p \leq d_\varphi, p$  odd, the isomorphism is given by multiplication with  $\gamma_p(\varphi)$  and for  $p = 0$  by the normmap to  $K_n^M F$ .

Conjecture 8 is definitely false for  $n \geq 3$  (e.g. for  $n_\varphi = 2$  the motive  $M_\varphi$  is the conic corresponding to  $\varphi$  and  $K_3^M F \xrightarrow{[M_\varphi]} A_1(M_\varphi; K_2^M)$  is neither surjective nor injective in general). Nevertheless the classes  $\gamma_p(\varphi)$  should form a (part of a) fundamental set of generators of the groups  $A_p(M_\varphi; K_n^M)$ .