A NOTE ON THE QUILLEN-LICHTENBAUM CONJECTURE AND THE ARITHMETIC OF SQUARE RINGS

G.BANASZAK, W.GAJDA, P.KRASOŃ AND P.ZELEWSKI

ABSTRACT. In this paper we consider the Quillen-Lichtenbaum conjecture for number fields using description of the higher K -theory in terms of SK_1 groups.

1. INTRODUCTION

Let F be a number field and let n be a non-negative integer. Fix an odd prime l . We define the wild kernel $WK_n(F)$ as the kernel of the localization map in K -theory of F :

$$\lambda_n : K_n(F) \rightarrow \prod_v K_n(F_v).$$

The symbol $\prod_v$ denotes the usual direct product. When n is even, the image of λ_n is contained in $\bigoplus_v K_n(F_v)$. However for n odd, the image is not contained in the direct sum, cf. diagram (3.4) below and [AB, Theorem 1]. The group $WK_n(F)$ is interesting because we have:

Proposition A. *[Prop. 5.1.] For each $n \geq 0$*

$$WK_n(F) \subseteq K_n(\mathcal{O})_{tor},$$

where $\mathcal{O}$ denotes the ring of integers of F . In particular, the group $WK_n(F)$ is finite.

By [B, theorem 2] there is the following long exact sequence.

$$(1.1) \quad 0 \rightarrow K_{2n}^w(\mathcal{O})_l \rightarrow K_{2n}(F)_l \rightarrow \bigoplus_v W^n(F_v) \rightarrow W^n(F) \rightarrow 0,$$

where v runs over finite places of F . In [B] the group $K_{2n}^w(\mathcal{O})_l$ was called the l -part of the wild kernel for $K_{2n}(F)$. We expect that the groups $WK_{2n}(F)_l$ and $K_{2n}^w(\mathcal{O})_l$ coincide. It is unknown in the moment, since we do not know the K -theory with coefficients in $\mathbb{Z}/l^k$ of a local field in the case when l is divisible by the residue characteristic, except for the case of $\mathbb{Q}_l$ which follows by results of Bökstedt and Madsen [BM].

1991 *Mathematics Subject Classification.* Primary 19F27; Secondary 19Fxx.
This work has been sponsored in part by a KBN grant 2P03A 005 11.

We propose the following conjecture.

Conjecture B. *For all number fields F and for all $n \geq 0$ we have:*

$$WK_n(F)_l = [\text{div}K_n(F)]_l$$

where $\text{div}K_n(F)$ is the group of divisible elements in $K_n(F)$.

In section 5 of the paper we give the following condition for equivalence of the above conjecture with the Quillen-Lichtenbaum conjecture for the number field F [DF, p. 273].

Theorem C. [Th. 5.2.]

(a) *Assume that the following maps:*

$$K_{2m}(F_v)_l \rightarrow K_{2m}^{et}(F_v)_l$$

are isomorphisms for all v which divide l and for all $m > 0$. Then the Quillen-Lichtenbaum conjecture for F is equivalent to the conjecture B for all indices $n > 1$.

(b) *If $F = \mathbb{Q}$, then the Quillen-Lichtenbaum conjecture and conjecture B are equivalent.*

Theorem D. [Th. 5.5.] *The conjecture B holds true for all number fields F and for $0 \leq n \leq 3$. In addition, it holds true for $n = 4$ and $F = \mathbb{Q}$.*

For another reformulation of the Quillen-Lichtenbaum conjecture see [BZ1, p. 390].

By [BZ2, theorem, p.161] and a result of Dayton and Weibel [DW, Corollary 5.3] we get the following exact sequence involving the wild kernel (see Section 6).

$$0 \rightarrow WK_n(F) \rightarrow SK_1(F_n) \rightarrow \prod_v SK_1(F_{vn}^h).$$

In this way we reformulate the conjecture B in terms of elementary linear algebra, number fields and algebraic properties of the square rings F_n . In the last section of the paper we investigate $WK_n(F)$ using the description of the higher K -theory in terms of SK_1 .

NOTATION

- (1) We fix an odd rational prime l .
- (2) Let F be a number field and let $\mathcal{O}$ be its ring of integers.
- (3) Let v be a finite place of $\mathcal{O}$ and let k_v be the residue field for v .
- (4) Put F_v to be the completion of F at v and $\mathcal{O}_v$ to be the ring of integers of F_v .
- (5) Put $\mathcal{O}_v^h$ to be the henselization of $\mathcal{O}_F$ at v and F_v^h to be the quotient field of $\mathcal{O}_v^h$.

- (6) For a commutative ring with identity R and for $n > 0$ we denote by R_n the square ring:

$$R[t_1, t_2, \dots, t_n]/((t_1^2 - t_1) \dots (t_n^2 - t_n)).$$

- (7) Let $K_*(-)$ ($K_*(\cdot, \mathbb{Z}/l^k)$ resp.) denote the Quillen K -theory (K -theory with coefficients defined by Browder and Karoubi, resp.).
- (8) Let $K_*^{et}(\cdot)$ ($K_*^{et}(\cdot, \mathbb{Z}/l^k)$, resp.) denote the etale K -theory (etale K -theory with finite coefficients, resp.) defined by Dwyer and Friedlander.
- (9) For an abelian group G we denote by G_l its l -torsion part and by $G[l^k]$ the elements of G with orders dividing l^k .
- (10) $\text{div } A = \bigcap_{r \geq 1} A^r$ denotes the subgroup of infinitely divisible elements inside an abelian group A .

ACKNOWLEDGMENTS

The first three authors thank the Max-Planck-Institut für Mathematik in Bonn and Fakultät für Mathematik Bielefeld Universität in Germany where parts of this work were done. The second author was supported by a research fellowship of Alexander von Humboldt Foundation at Bielefeld Universität when this paper was completed. He thanks both institutions for support and hospitality. The authors thank the referee for careful reading of the manuscript and useful remarks.

2. INTRODUCTORY COMPUTATIONS

The computations below follow easily from well-known results but we include some detailed explanations for the convenience of the reader. By [Su1] we have the following isomorphism for all l different from the residue characteristic of k_v , for all $j > 0$ and all nonnegative n .

$$K_n(\mathcal{O}_v; \mathbb{Z}/l^j) \rightarrow K_n(k_v; \mathbb{Z}/l^j)$$

Using the Bockstein exact sequences for K -theory with finite coefficients we get the following diagram.

$$(2.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & K_{2n}(\mathcal{O}_v)/l^j & \longrightarrow & K_{2n}(\mathcal{O}_v; \mathbb{Z}/l^j) & \xrightarrow{\sim} & K_{2n-1}(\mathcal{O}_v)[l^j] \longrightarrow 0 \\ & & \downarrow \sim & & \sim \downarrow & & \downarrow \sim \\ 0 & \longrightarrow & 0 & \longrightarrow & K_{2n}(k_v; \mathbb{Z}/l^j) & \xrightarrow{\sim} & K_{2n-1}(k_v)[l^j] \longrightarrow 0 \end{array}$$

The diagram (2.1) shows that the group $K_{2n}(\mathcal{O}_v)$ is l -divisible for all $n > 0$. Consider also the following commutative diagram:

$$(2.2) \quad \begin{array}{ccccccc} 0 & \longrightarrow & K_{2n+1}(\mathcal{O}_v)/l^j & \xrightarrow{\sim} & K_{2n+1}(\mathcal{O}_v; \mathbb{Z}/l^j) & \longrightarrow & K_{2n}(\mathcal{O}_v)[l^j] \longrightarrow 0 \\ & & \sim \downarrow & & \sim \downarrow & & \sim \downarrow \\ 0 & \longrightarrow & K_{2n+1}(k_v)/l^j & \xrightarrow{\sim} & K_{2n+1}(k_v; \mathbb{Z}/l^j) & \longrightarrow & 0 \longrightarrow 0 \end{array}$$

Consider $n > 0$. From the diagram (2.1) we see that if j is big enough, then we have:

$$(2.3) \quad K_{2n-1}(\mathcal{O}_v)_l = K_{2n-1}(k_v)_l = \text{finite}$$

Hence, the l -torsion part of the group $K_{2n-1}(\mathcal{O}_v)_l$ can not be divisible. In addition, we see that:

$$K_{2n-1}(k_v)_l = K_{2n-1}(k_v)/l^j$$

so the left vertical arrow in the diagram (2.2) is surjective for j big enough. This implies that the right vertical arrow in (2.2) is an isomorphism. Hence, $K_{2n}(\mathcal{O}_v)_l = 0$. This shows that the left square in the diagram (2.2) consists of isomorphisms. So the group $K_{2n}(\mathcal{O}_v)$ is uniquely l -divisible. Let us consider the following localization exact sequence.

$$(2.4) \quad 0 \rightarrow K_{2n}(\mathcal{O}_v) \rightarrow K_{2n}(F_v) \rightarrow K_{2n-1}(k_v) \rightarrow 0$$

We know by [BZ1, p. 386] that:

$$K_{2n}(F_v) = K_{2n}(F_v)_l \bigoplus l^{j_0} K_{2n}(F_v)$$

and the group $l^{j_0} K_{2n}(F_v)$ is uniquely l -divisible. We can also write:

$$K_{2n-1}(k_v) = K_{2n-1}(k_v)_l \bigoplus (\text{non } l\text{-part}).$$

Hence, we observe that the boundary map in the localization sequence (2.4) sends the subgroup $K_{2n}(F_v)_l \bigoplus 0$ of $K_{2n}(F_v)$ into:

$$K_{2n-1}(k_v)_l \bigoplus 0 \subseteq K_{2n-1}(k_v)$$

and it sends $0 \bigoplus l^{j_0} K_{2n}(F_v)$ into:

$$0 \bigoplus (\text{non } l \text{ part}) \subseteq K_{2n-1}(k_v).$$

It follows that the map:

$$(2.5) \quad K_{2n}(F_v)_l \xrightarrow{\sim} K_{2n-1}(k_v)_l.$$

is surjective. We know that this map is injective because $K_{2n}(\mathcal{O}_v)_l = 0$.

3. THE CASE OF ODD INDICES

Observe that by [DF, Theorem 8.7] and [So2, Theorem 1, p. 375] we have the following isomorphism:

$$(3.1) \quad K_{2n+1}(\mathcal{O}[1/l]) \otimes \mathbb{Z}_l/torsion \rightarrow K_{2n+1}^{et}(\mathcal{O}[1/l])/torsion.$$

This implies that we have the following surjection:

$$K_{2n+1}(\mathcal{O}[1/l])_l \rightarrow K_{2n+1}^{et}(\mathcal{O}[1/l])_l.$$

Assume that v does not divide l . Consider the following commutative diagram:

$$(3.2) \quad \begin{array}{ccc} K_{2n+1}(\mathcal{O})_l & \longrightarrow & K_{2n+1}(k_v)_l \\ \downarrow & & \downarrow \sim \\ K_{2n+1}^{et}(\mathcal{O}[1/l])_l & \longrightarrow & K_{2n+1}^{et}(k_v) \\ \downarrow \sim & & \downarrow \sim \\ H^1(\mathcal{O}[1/l]; \mathbb{Z}_l(n+1))_l & \longrightarrow & H^1(k_v; \mathbb{Z}_l(n+1)) \\ \downarrow \sim & & \downarrow \sim \\ H^0(\mathcal{O}[1/l]; \mathbb{Q}_l/\mathbb{Z}_l(n+1)) & \longrightarrow & H^0(k_v; \mathbb{Q}_l/\mathbb{Z}_l(n+1)) \end{array}$$

It is easy to see that the bottom horizontal arrow in the diagram (3.2) is an imbedding for all v which do not divide l . This shows that the kernel of the upper horizontal arrow is equal to the kernel of the upper, left vertical arrow. Hence, we can conclude the following result.

Lemma 3.3. *The Quillen-Lichtenbaum conjecture holds for $K_{2n+1}(\mathcal{O})$ if and only if the reduction map:*

$$\pi : K_{2n+1}(\mathcal{O})_l \rightarrow K_{2n+1}(k_v)$$

is injective for some v which does not divide l . The Quillen-Lichtenbaum conjecture for $K_{2n+1}(\mathcal{O})$ is also equivalent to the injectivity of the map:

$$\pi : K_{2n+1}(\mathcal{O})_l \rightarrow \prod_v K_{2n+1}(k_v).$$

Let us consider the following diagram:

$$(3.4) \quad \begin{array}{ccc} K_{2n+1}(F) & \longrightarrow & \prod_v K_{2n+1}(F_v) \\ \parallel & & \parallel \\ K_{2n+1}(\mathcal{O}) & \longrightarrow & \prod_v K_{2n+1}(\mathcal{O}_v) \\ \parallel & & \downarrow \prod_v \pi_v \\ K_{2n+1}(\mathcal{O}) & \xrightarrow{\pi} & \prod_v K_{2n+1}(k_v) \end{array}$$

By the lemma 3.3, the diagram (3.4) and the isomorphism (2.3) we obtain the following proposition.

Proposition 3.5. *The Quillen-Lichtenbaum conjecture holds for K_{2n+1} , $n > 0$, if and only if the following localization map:*

$$K_{2n+1}(F)_l \rightarrow \prod_v K_{2n+1}(F_v)_l$$

is injective.

4. THE CASE OF EVEN INDICES

Consider the following commutative diagram.

$$(4.1) \quad \begin{array}{ccc} K_{2n}(F) & \longrightarrow & \bigoplus_v K_{2n}(F_v) \\ \parallel & & \bigoplus_v \delta_v \downarrow \\ K_{2n}(F) & \xrightarrow{\delta} & \bigoplus_v K_{2n-1}(k_v) \end{array}$$

The maps δ and δ_v come from Quillen localization sequences in K-theory [Q1]. We can write $\bigoplus_v$ instead of $\prod_v$ for the upper horizontal arrow in diagram (4.1), because of the isomorphism (2.5). The diagram shows that:

$$WK_{2n}(F) \subseteq K_{2n}(\mathcal{O}).$$

The wild kernel $K_{2n}^w(\mathcal{O})_l$ was defined by (1.1) cf. [B, p. 288]. At the moment we can not compare the groups $K_{2n}^w(\mathcal{O}_F)_l$ and $WK_{2n}(F)_l$, since we do not know the groups $K_n(\mathcal{O}_v; \mathbb{Z}/l^k)$ for v dividing l except for the case when $F = \mathbb{Q}$ which is known by results of [BM]. Observe that the definition of the group $WK_n(F)$ makes sense for every l , including $l = 2$ and that for $n = 2$ it agrees with the definition of the classical wild kernel for the functor K_2 . By [DF, theorem 8.5, p.275] and by Bockstein sequences we obtain the following natural surjection:

$$K_{2n}(F_v)_l \rightarrow K_{2n}^{et}(F_v)_l$$

for all v . This surjection is an isomorphism for every v which does not divide l , as we can easily check using results of [Su1] which have been already mentioned. In addition, due to [DF, Prop. 5.1, p.260] and [T, prop. 2.3] we have natural isomorphisms:

$$K_{2n}^{et}(F_v)_l \cong H^2(F_v; \mathbb{Z}_l(n+1))_l \cong H^1(F_v; \mathbb{Q}_l/\mathbb{Z}_l(n+1))/Div.$$

Let $D_{n+1}(F)$ denote the group of divisible elements in $H^2(F; \mathbb{Z}_l(n+1))_l$ cf. [Sch, p. 192]. Consider the following commutative diagram.

$$(4.2) \quad \begin{array}{ccccccc} 0 & \longrightarrow & WK_{2n}(F)_l & \longrightarrow & K_{2n}(F)_l & \longrightarrow & \bigoplus_v K_{2n}(F_v)_l \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & D_{n+1}(F) & \longrightarrow & H^2(F; \mathbb{Z}_l(n+1))_l & \longrightarrow & \bigoplus_v H^2(F_v; \mathbb{Z}_l(n+1))_l \end{array}$$

Lemma 4.3. *Assume that the natural maps:*

$$K_{2n}(F_v)_l \rightarrow K_{2n}^{et}(F_v)_l$$

are isomorphisms for all v which divide l . Then the Quillen-Lichtenbaum conjecture for K_{2n} , $n > 0$, is equivalent to the statement that the left vertical arrow in the diagram (4.2) is an isomorphism.

Proof. Observe that by our assumption the right vertical arrow in (4.2) is an isomorphism. Simple diagram chasing in the diagram (4.2) shows that the left vertical arrow is an isomorphism if and only if the middle vertical arrow is an isomorphism. Hence, our lemma follows by [B, remark 7, p. 295]. $\square$

5. COMPARISON OF CONJECTURES

The diagrams (3.4), (4.1) and [AB, Theorem 1] give us the following statement.

Proposition 5.1. *For each $n \geq 0$:*

$$WK_n(F) \subseteq K_n(\mathcal{O})_{tor},$$

hence the group $WK_n(F)$ is always finite.

Our computations of the previous sections suggest that we should expect that the following conjecture holds true.

Conjecture B. *Up to 2-torsion:*

$$WK_n(F) = \text{div} K_n(F)$$

for any n and for any number field F .

Theorem 5.2.

(a) *Assume that the following maps:*

$$K_{2m}(F_v)_l \rightarrow K_{2m}^{et}(F_v)_l$$

are isomorphisms for all v which divide l and for all $m > 0$. Then the Quillen-Lichtenbaum conjecture for F is equivalent to the conjecture B for all indices $n > 1$.

(b) *If $F = \mathbb{Q}$, then the Quillen-Lichtenbaum conjecture and conjecture B are equivalent for $n > 1$.*

Proof. a)

We observe that $K_{2n+1}(F) = K_{2n+1}(\mathcal{O})$ for $n > 0$ due to [Q1, Theorem 5] and [So1, Theoreme 3, p. 274]. In addition, $K_{2n+1}(\mathcal{O})$ is a finitely generated abelian group [Q2, Theorem 1, p. 179]. Hence $\text{div} K_{2n+1}(F) = 0$ for $n \geq 0$. Note that $K_1(F) = F^\times$ [Mi, p. 28] so $\text{div} K_1(F) = 0$. So the theorem follows by proposition 3.5 and lemma 4.3.

b) By the results of Bökstedt and Madsen [BM, p. 29], see also [HM, p. 30] the assumption of point (a) holds true for $F = \mathbb{Q}$. $\square$

Proposition 5.3. *For every odd prime l we have:*

$$\begin{aligned} \operatorname{div} K_n(\mathbb{Q})_l &\subset WK_n(\mathbb{Q})_l \quad \text{for } n \geq 0 \\ K_{2n}^w(\mathbb{Z})_l &= WK_{2n}(\mathbb{Q})_l \quad \text{for } n > 0. \end{aligned}$$

Proof. Let us first prove the first statement. For $F = \mathbb{Q}$ and n even, the right vertical arrow in diagram 4.2 is an isomorphism. In addition the group $H^2(F_v; \mathbb{Z}_l(n+1))_l$ is finite [Sch, Satz 4(i) p. 188] for every l odd and every finite prime v of F . Hence, the group $\operatorname{div} K_{2n}(\mathbb{Q})_l$ is contained in the kernel of the upper, right horizontal arrow of diagram 4.2. For n odd the first statement is trivial since $\operatorname{div} K_{2n+1}(F) = 0$ for all $n \geq 0$. The second statement follows from diagram 4.2 and the definition of $K_{2n}^w(\mathbb{Z})_l$ cf. [B]. $\square$

Remark 5.4. It follows by [B, theorem 3, p. 289] that:

$$\operatorname{div} K_{22}(\mathbb{Z})_{\text{odd}} = \mathbb{Z}/691.$$

In addition we have:

$$\operatorname{div} K_{22}(\mathbb{Z})_l \subset WK_{22}(\mathbb{Q})_l \subset K_{22}(\mathbb{Z})_l$$

Let us use the isomorphism $K_{22}(\mathbb{Z}) = SK_1(\mathbb{Z}_{22})$. We can find a matrix $A \in SL(\mathbb{Z}_{22})$ corresponding to an element of order 691 in $\operatorname{div} K_{22}(\mathbb{Z})$, such that its inverse can be found by the standard formula involving adjoint matrix, but we cannot find the inverse of A using Gauss-Jordan algorithm. In addition let us consider A as a matrix in $SL((\mathbb{Z}_l)_{22})$. Then by proposition 5.3 we are able to find the inverse of A using Gauss-Jordan algorithm in $SL((\mathbb{Z}_l)_{22})$ for every l .

Theorem 5.5. *Conjecture B holds true for all number fields F and for $0 \leq n \leq 3$. In addition, it holds true for $n = 4$ and $F = \mathbb{Q}$.*

Proof. It is obvious for $n = 0$ and $n = 1$. For $n = 2$ it was proven by Tate cf. [Ba, p.250]. For $n = 3$ this is a consequence of the well-known result of Merkurjev and Suslin [MS, p.342] which verifies the Quillen-Lichtenbaum conjecture for K_3 , so we can see that our conjecture B holds in this case as follows by the diagram (3.2). By [RW] we know that $K_4(\mathbb{Z}) = 0$. Hence, $\operatorname{div} K_4(\mathbb{Q}) = 0$ and $WK_4(\mathbb{Q}) = 0$ as subgroups of $K_4(\mathbb{Z})$. $\square$

6. CONJECTURE B AND MATRICES

In this section, for the convenience of the reader, we enclose some results from the paper [BZ2]. Let R denote any commutative ring with identity. For $n > 0$ let us denote by R_n the ring:

$$R[t_1, t_2, \dots, t_n] / ((t_1^2 - t_1) \dots (t_n^2 - t_n)).$$

For $n = 0$ we put $R_0 = R$. The ring R_n is called the square ring. To simplify notation we put $T = (t_1, t_2, \dots, t_n)$ and

$$\alpha(T) = (t_1^2 - t_1) \dots (t_n^2 - t_n).$$

Observe that:

$$\alpha(T) = t_1^2 t_2^2 \dots t_n^2 + (\text{terms of total degree smaller than } 2n).$$

Lemma 6.1. [BZ2, Lemma 1] *If $R \subseteq S$ are two commutative rings with identity, then the induced map $R_n \rightarrow S_n$ is an imbedding.*

Lemma 6.2. [DW, Corollary 5.3] *Let R be a domain. Then $R_n^* = R^*$ for $n > 1$ and $R_1^* = R^* \oplus R^*$, where by the upper star we denote the group of units.*

Remark 6.3. Let R be a commutative, regular ring with identity. Then by [DW, Corollary 5.4] or by [Su2, p. 351], we have for $n > 0$:

$$K_n(R) = K_1(R_n)/K_1(R).$$

Remark 6.4. Let R be a field, the ring of integers in a number field or a local ring. Then $K_1(R) = R^*$ [Mi, p. 28]. Hence, if R is a field or the ring of integers in a number field or R is a regular, local ring which is a domain, then by the remark 6.3 and lemma 6.2 we get:

$$K_n(R) = K_1(R_n)/K_1(R) = (SK_1(R_n) \oplus R^*)/R^* = SK_1(R_n)$$

cf. [DW, Corollary 5.3]. In particular, the definition of the group $WK_n(F)$ can be rewritten as the following exact sequence.

$$0 \rightarrow WK_n(F) \rightarrow SK_1(F_n) \rightarrow \prod_v SK_1(F_{vn})$$

Remark 6.5. Let $A \in SL(F_n)$. Let $[A]$ denote the class of the matrix A in the group $SK_1(F_n)_{tor}$. Then $[A] \in WK_n(F)$ if and only if A is row- or column-equivalent or both row- and column-equivalent to the identity matrix in $SL(F_{vn})$ for each v . It is easy to show that in the case of even indices we have the following exact sequence.

$$0 \rightarrow WK_{2n}(F) \rightarrow K_{2n}(\mathcal{O}[1/l]) \rightarrow \prod_{v|l} K_{2n}(F_v)$$

Remark 6.6. Comparison of the local and global localization sequences shows that we have the following exact sequence for all $n > 0$.

$$0 \rightarrow WK_n(F) \rightarrow SK_1(\mathcal{O}_n) \rightarrow \prod_v SK_1(\mathcal{O}_{vn})$$

It was proven in [BZ2] that the natural maps:

$$K_n(\mathcal{O}_v^h) \rightarrow K_n(\mathcal{O}_v)$$

$$K_n(F_v^h) \rightarrow K_n(F_v)$$

are injective. Hence, we obtain the following exact sequences.

$$(6.7) \quad 0 \rightarrow WK_n(F) \rightarrow SK_1(F_n) \rightarrow \prod_v SK_1(F_{vn}^h)$$

$$(6.8) \quad 0 \rightarrow WK_n(F) \rightarrow SK_1(\mathcal{O}_n) \rightarrow \prod_v SK_1(\mathcal{O}_{v_n}^h)$$

Observe that for any field extension $K \subset L$ we have the following exact sequence.

$$0 \rightarrow SL(K_n) \cap E(L_n)/E(K_n) \rightarrow SK_1(K_n) \rightarrow SK_1(L_n)$$

From (6.7) and (6.8) we obtain the following equality.

$$WK_n(F) = \left(SL(F_n) \cap \bigcap_v E(F_{v_n}^h) \right) / E(F_n) = \left(SL(\mathcal{O}_n) \cap \bigcap_v E(\mathcal{O}_{v_n}^h) \right) / E(\mathcal{O}_n)$$

Remark 6.9. Observe that the conjecture B says that for each odd prime l we should have:

$$WK_n(F)_l = \bigcap_{r \geq 1} (SK_1(F_n)_l)^r.$$

We can also look at the question of descent using the SK_1 description of K -theory. Let L/K be a Galois extension with the Galois group $G = G(L/K)$.

Proposition 6.10. *The natural map:*

$$K_n \xrightarrow{\sim} L_n^G$$

is an isomorphism.

Proof. Consider the following commutative diagram.

$$(6.11) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \alpha(T)K[T] & \longrightarrow & K[T] & \longrightarrow & K_n \longrightarrow 0 \\ & & \downarrow \sim & & \downarrow \sim & & \downarrow \\ 0 & \longrightarrow & \alpha(T)L[T]^G & \longrightarrow & L[T]^G & \longrightarrow & L_n^G \longrightarrow H^1(G; \alpha(T)L[T]) \end{array}$$

We observe that:

$$\alpha(T)L[T] \xrightarrow{\sim} \bigoplus L$$

as G -modules, where the direct sum is over all monomials $t_1^{r_1} t_2^{r_2} \dots t_n^{r_n}$ of $L[T]$ for $r_i \geq 0$. By [CF, chapter V.2.6, Proposition 2.1] we know that $H^1(G; L) = 0$. Hence:

$$H^1(G; \alpha(T)L[T]) = H^1(G; \bigoplus L) = \bigoplus H^1(G; L) = 0$$

and the proposition follows from the five lemma applied to the diagram (6.11). $\square$

Corollary 6.12. *The isomorphism from proposition 6.10 induces the following isomorphism.*

$$SL(K_n) \xrightarrow{\sim} SL(L_n)^G$$

Remark 6.13. Observe that we only have $E(K_n) \subseteq E(L_n)^G$. Consider the following commutative diagram.

$$(6.14) \quad \begin{array}{ccccccc} 0 & \longrightarrow & E(K_n) & \longrightarrow & SL(K_n) & \longrightarrow & SK_1(K_n) \longrightarrow 0 \\ & & \downarrow & & \downarrow \sim & & \downarrow \\ 0 & \longrightarrow & E(L_n)^G & \longrightarrow & SL(L_n)^G & \longrightarrow & SK_1(L_n)^G \end{array}$$

Remark 6.15. The diagram (6.14) implies that we have the following exact sequence.

$$0 \rightarrow SL(K_n) \cap E(L_n) / E(K_n) \rightarrow K_n(K) \rightarrow K_n(L)^G \rightarrow SK_1(L_n)^G / \text{Im} SL(K_n) \rightarrow 0$$

7. CONDITIONS FOR TRIVIALITY OF ELEMENTS IN $WK_n(F)$

Let R be a commutative ring with identity. Let $A \in E(R)$ and let us write $A = \prod_{i,j,\lambda} e_{i,j}^\lambda$ in terms of elementary matrices.

Lemma 7.1. *Let A be a product of elementary matrices $e_{i,j}^\lambda$ that are pairwise commutative. Then all entries of the matrix A which are on the main diagonal are equal to 1 and all other entries are linear homogeneous polynomials in λ 's. The polynomials have integral coefficients.*

Proof. The proof is by induction on the number of elementary matrices. It is obvious for one such a matrix. Let $B = (b_{ij})_{1 \leq i,j \leq m} \in GL(R)$ which commutes with elementary matrices $e_{i,j}^\lambda \in E(R)$. For our computations we assume that $i < j$. The case $i > j$ is similar. Consider products of matrices:

$$Be_{i,j}^\lambda = \begin{pmatrix} b_{11} & b_{12} & \dots & b_{1j-1} & b_{1j} + \lambda b_{1i} & \dots & b_{1m} \\ b_{21} & b_{22} & \dots & b_{2j-1} & b_{2j} + \lambda b_{2i} & \dots & b_{2m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ b_{m1} & b_{m2} & \dots & b_{mj-1} & b_{mj} + \lambda b_{mi} & \dots & b_{mm} \end{pmatrix}$$

On the other hand:

$$e_{i,j}^\lambda B = \begin{pmatrix} b_{11} & b_{12} & \dots & b_{1m} \\ b_{21} & b_{22} & \dots & b_{2m} \\ \vdots & \vdots & \vdots & \vdots \\ b_{(i-1)1} & b_{(i-1)2} & \dots & b_{(i-1)m} \\ \vdots & \vdots & \vdots & \vdots \\ b_{i1} + \lambda b_{j1} & b_{i2} + \lambda b_{j2} & \dots & b_{im} + \lambda b_{jm} \\ \vdots & \vdots & \vdots & \vdots \\ b_{m1} & b_{m2} & \dots & b_{mm} \end{pmatrix}$$

Comparison of entries gives: $b_{kj} + \lambda b_{ki} = b_{kj}$ for $k \neq i$. So, $\lambda b_{ki} = 0$. In the same way $b_{il} + \lambda b_{jl} = b_{il}$ for $l \neq j$ and $\lambda b_{jl} = 0$. At the end we compare the ij entries: $b_{ij} + \lambda b_{ii} = b_{ij} + \lambda b_{jj}$. This implies that $\lambda(b_{ii} - b_{jj}) = 0$. So in particular if $b_{ii} = b_{jj}$, the ij entries of the matrices in question are equal. To continue observe that if $A = Be_{i,j}^\lambda$, where B is a product of pairwise commuting elementary matrices, then by inductive hypothesis B is a matrix with 1's on the main diagonal and the other entries are linear homogeneous polynomials of λ 's. Let us denote a_{ij} the entries of A . Hence the above comparison of entries shows that all the entries of A on the main diagonal are equal to 1. It is so because $a_{rr} = b_{rr} = 1$ for $r \neq j$. In addition $a_{jj} = b_{jj} + \lambda b_{ji} = b_{jj} = 1$. Also all entries of A beyond the j -th column are just the same as the corresponding entries of B . Using again the comparison of entries due to the equation $Be_{i,j}^\lambda = e_{i,j}^\lambda B$ we see that in the j -th column $a_{kj} = b_{kj} + \lambda b_{ki} = b_{kj}$ for $k \neq i$ so this case follows by inductive hypothesis. On the other hand for $k = i$ we have $a_{ij} = b_{ij} + \lambda b_{ii} = b_{ij} + \lambda$ because $b_{ii} = 1$ by the induction. $\square$

Proposition 7.2. *Let $A \in SK_1(F_n)$. Assume that for some finite prime v , A can be written as a product of elementary, pairwise commuting matrices in $SL_1(F_{vn}^h)$. Then A determines a divisible element in $SK_1(F_n)$.*

Proof. Let $A = \prod_{i,j,\lambda} e_{i,j}^\lambda$ in $SL_1(F_{vn}^h)$ be a product of elementary, pairwise commutative matrices. For any natural number m consider the matrix $C = \prod_{i,j,\lambda} e_{i,j}^{\lambda/m}$. The matrices of the form $e_{i,j}^{\lambda/m}$ also commute pairwise. Hence by lemma 8.1 we see that C comes from an element of $SL(F_n)$. Let us denote this element C_0 . But $C_0^m = C^m = A$ in $SL(F_{vn}^h)$. Hence by lemma 6.1 we get $C_0^m = A$ in $SL(F_n)$. $\square$

Corollary 7.3. *Let $A \in SK_1(F_n)$ be an element of $WK_n(F)$. Assume that for some finite prime v , A can be written as a product of elementary, pairwise commuting matrices in $SL(F_{vn}^h)$. Then A determines a divisible element in $SK_1(F_n)$.*

Remark 7.4. Let $A \in SL(F_n)$ be a matrix whose class mod $E(F_n)$ is in $WK_n(F)_l$. Proposition 7.2 and corollary 7.3 show, that if a matrix A is a product of pairwise commutative matrices in $SL(F_{vn}^h)$, for some v , then this matrix A gives a trivial class in $SK_1(F_n)$. Indeed, computations in proposition 7.2 and corollary 7.3 show that for each power l^k of an odd prime number l we can find a matrix $B_k \in SL(F_n)$ such that $A = B_k^{l^k}$. Observe that actually $A \in SL(\mathcal{O}_{S,n})$ for some finite set S of prime numbers containing l and $B \in SL(\mathcal{O}_{S,n})$ by computations. Hence the class of A is divisible by all powers of l in a finite group $K_n(\mathcal{O}_S)_{tor}$. Hence, the class of A must be trivial. This also follows from the corollary 7.6 below.

Let us note first that every elementary matrix with coefficients in any commutative ring can be written as a product of at most four unipotent matrices:

$$U_1 L_1 U_2 L_2 \quad \text{or} \quad L_1 U_1 L_2 U_2,$$

where U_i (L_i , respectively) denote upper (lower, resp.) triangular matrix with 1 on the diagonal cf. [vdK, LNM 966, p.358]. Conversely every upper or lower triangular matrix with 1 on the diagonal is clearly a product of elementary matrices, since we can bring such a matrix to the identity matrix by a series of elementary operations.

Proposition 7.5. *Let R be a ring contained as a subring in S . Let A be a matrix in $SL(R)$ such that it is a product of at most two unipotent triangular matrices with coefficients in S . Then the unipotent matrices have coefficients in R .*

Proof. The case when A is a unipotent matrix is obvious. Let A be a product of two unipotent matrices: $A = UL$, where U is upper triangular and L is lower triangular. The case when $A = LU$ is handled similarly. Let us put:

$$U = \begin{pmatrix} 1 & u_{12} & u_{13} & \dots & \dots & \dots & \dots & u_{1m} \\ 0 & 1 & u_{23} & \dots & \dots & \dots & \dots & u_{2m} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & \dots & u_{i(i+1)} & \dots & u_{im} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & & & \dots & & & 1 \end{pmatrix}$$

$$L = \begin{pmatrix} 1 & 0 & 0 & & \dots & & & 0 \\ l_{21} & 1 & 0 & & \dots & & & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ l_{j1} & \dots & l_{j(j-1)} & 1 & \dots & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ l_{m1} & l_{m2} & \dots & \dots & \dots & \dots & l_{m(m-1)} & 1 \end{pmatrix}.$$

The product UL is the matrix A with coefficients in R . Observe that multiplying rows of U by last column of L we can figure out that last column of U has coefficients in R . Also multiplying last row of U by columns of L we see that last row of L has coefficients in R . Now the argument goes by induction. Assume that last k columns of U and last k rows of L are already in R . Multiplying the $m - k$ -th row of U by j -th column of L for $j < m - k$ we get:

$$l_{(m-k)j} + \sum_{i=m-k+1}^m u_{(m-k)i} l_{ij} \in R.$$

Hence, by the induction hypotheses $l_{(m-k)j} \in R$ for all $j < m - k$. In the same way, multiplying i -th row of U for $i < m - k$ by $m - k$ -th column of L we get:

$$u_{i(m-k)} + \sum_{j=m-k+1}^m u_{ij} l_{j(m-k)} \in R.$$

Hence, by the induction hypotheses $u_{i(m-k)} \in R$ for all $i < m - k$. We proved that last $k + 1$ columns of U and last $k + 1$ rows of L are in R . The claim follows by induction. $\square$

Corollary 7.6. *Let $A \in SL(F_n)$ be a matrix whose class modulo $E(F_n)$ is in $WK_n(F)$. Let for some v the matrix A be a product in $SL(F_{vn})$ (or $SL(F_{vn}^h)$) of at most two unipotent triangular matrices or A be a product of four unipotent triangular matrices such that the first and fourth factors of this product have coefficients in F_n . Then the class of A is trivial in $WK_n(F)$.*

Proof. Follows immediately from proposition 7.5. $\square$

Remark 7.7. A unipotent triangular matrix from $SL(F_{vn})$ can be written in the form $I + N$ where N is an upper or lower triangular with zero's on the main diagonal. Coefficients of N are elements of F_{vn} so for each natural number k every such coefficient can be written as a sum of an element of F_n and an element from $(\pi_v)^k \mathcal{O}_{vn}$, where π_v is a uniformizer of the maximal ideal of $\mathcal{O}_v$. Hence, we can write:

$$I + N = (I + N_0)(I + (\pi_v)^k N_1) \quad \text{or} \quad I + N = (I + (\pi_v)^k N'_1)(I + N_0),$$

where N_0 and N_1 are upper (lower) triangular and nilpotent. N_0 has coefficients in F_n and N_1, N'_1 have coefficients in $\mathcal{O}_{vn}$.

Proposition 7.8. *Let $[A] \in WK_n(F)$. For each prime v there is a matrix $B \in SL(\mathcal{O}_n)$ such that $[A] = [B]$ in $WK_n(F)$ and:*

$$B \equiv I \pmod{v}.$$

Proof. Let $A \in SL(\mathcal{O}_n)$ be a matrix whose class modulo $E(\mathcal{O}_n)$ is in $WK_n(F)$. Any element in $WK_n(F)$ can be represented in such a way by the exact sequence (6.8). Pick a prime ideal v and write:

$$A = U_1 L_1 U_2 L_2,$$

where the matrices in the product have coefficients in $\mathcal{O}_{vn}$. The case when the product on the right of the last matrix equation starts with a lower triangular matrix is analogous. By remark 7.7 we can write:

$$U_1 = U'_1 U''_1 \quad \text{and} \quad L_2 = L''_2 L'_2,$$

where U'_1 is a unipotent matrix with coefficients above main diagonal in the ring $\mathcal{O}_n$ and U''_1 is a unipotent matrix with coefficients above the main diagonal in $(\pi_v)^k \mathcal{O}_{vn}$. Similarly, L'_2 is a unipotent matrix with coefficients below the main diagonal in $\mathcal{O}_n$ and L''_2 is a unipotent matrix with coefficients below the main diagonal in $(\pi_v)^k \mathcal{O}_{vn}$. Hence, we get:

$$U'^{-1}_1 A L'^{-1}_2 = U''_1 L_1 U_2 L''_2.$$

Matrices A and $B = U'^{-1}_1 A L'^{-1}_2$ are in the same class modulo $E(\mathcal{O}_n)$. Again by remark 7.7 we can also write:

$$L_1 = L''_1 L'_1 \quad \text{and} \quad U_2 = U'_2 U''_2,$$

where U'_2 is a unipotent matrix with coefficients above main diagonal in the ring $\mathcal{O}_n$ and U''_2 is a unipotent matrix with coefficients above the main diagonal in $(\pi_v)^k \mathcal{O}_{vn}$. Similarly, L'_1 is a unipotent matrix with coefficients below the main diagonal in $\mathcal{O}_n$ and L''_1 is a unipotent matrix with coefficients below the main diagonal in $(\pi_v)^k \mathcal{O}_{vn}$. We have the following equality:

$$(7.9) \quad L'^{-1}_1 U'^{-1}_1 A L'^{-1}_2 U'^{-1}_2 = L'^{-1}_1 U''_1 L_1 U_2 L''_2 U'^{-1}_2.$$

Matrices A , $A_1 = U'^{-1}_1 A L'^{-1}_2$ and $A_2 = L'^{-1}_1 U'^{-1}_1 A L'^{-1}_2 U'^{-1}_2$ are in the same class modulo $E(\mathcal{O}_n)$. It is clear from (7.9) that:

$$A_2 \equiv I \pmod{v}.$$

□

REFERENCES

- [AB] D.Arlettaz and G.Banaszak, *On the non-torsion elements in the Algebraic K-theory of rings of integers*, J. Reine Angew. Math. **461** (1995), 63-79.
- [B] G.Banaszak, *Generalization of the Moore exact sequence and the wild kernel for higher K-groups*, Comp. math. **86**. (1993), 281-305.
- [BZ1] G.Banaszak and P.Zelewski, *Continuous K-theory*, K-theory **9** (1995), 379-392.
- [BZ2] G.Banaszak and P.Zelewski, *On the map between K-theory of henselization and completion of some local rings*, Journal of Pure and Applied Algebra **120** (1997), 161-165.
- [Ba] H.Bass, *K_2 des corps globaux*, Séminaire BOURBAKI, 23^e année n^o **394** (1970/71).
- [CF] J.W.S.Cassels and A.Frohlich, *Algebraic Number Theory*, Academic Press, 1967.
- [BM] M.Bökstedt and I.Madsen, *Algebraic K-theory of local number fields; the unramified case. Prospects in Topology.* (ed. F.Quinn), Ann. of Math. Stud. **138** (1995), Princeton Univ. Press, 28-57.
- [DW] B.H.Dayton and C.A.Weibel, *K-theory of hyperplanes*, Transactions of the AMS **257**, n.1 (1980), 119-141.
- [DF] W.Dwyer and E.Friedlander, *Algebraic and étale K-theory*, Transactions of the AMS **292**, n.1 (1985), 247-280.
- [HM] L.Hesselholt and I.Madsen, *On the K-theory of finite algebras over Witt vectors of perfect fields*, Topology **36** (1997), 29-101.
- [vdK] W.van der Kallen, *$SL_3(\mathbb{C}[x])$ does not have bounded word length*, Springer Lecture Notes in Mathematics **966**, 357-361.
- [MS] A.Merkurjev and A.Suslin, *On the K_3 of a field*, Math. USSR Izvestya **36**, n.3 (1991), 531-565.
- [Mi] J.Milnor, *Introduction to Algebraic K-theory*, Annals of Mathematics Studies, Princeton University Press, Princeton, N.J., 1971.
- [Q1] D. Quillen, *Higher Algebraic K-theory I*, Lecture Notes in Mathematics **341** (1973), 85-147.
- [Q2] D. Quillen, *Finite generation of the group K_i of ring of algebraic integers*, Lecture Notes in Mathematics **341** (1973), 179-198.
- [RW] J.Rognes and C.Weibel, *Two-primary algebraic K-theory of rings of integers in number fields*, preprint (1997).
- [Sch] P. Schneider, *Über gewisse Galoiskohomologiegruppen*, Math. Zeit. **168** (1979), 181-205.
- [So1] C. Soulé, *K-théorie de anneaux d'entiers de corps de nombres et cohomologie étale*, Inventiones math. **55** (1979), 251-295.
- [So2] C. Soulé, *On higher p-adic regulators*, Lecture Notes in Mathematics, **854** (1981), 372-401.
- [Su1] A.Suslin, *On the K-theory of local fields*, J. of Pure and Applied Algebra **34** (1984), 301-318.
- [Su2] A.Suslin, *Mennicke symbols and their applications in the K-theory of fields*, Lecture Notes in Mathematics **966** (1982), 334-356.
- [T] J. Tate, *Relations between K_2 and galois cohomology*, Inventiones math. **36** (1976), 257-274.

DEPARTMENT OF MATHEMATICS, ADAM MICKIEWICZ UNIVERSITY, POZNAŃ, POLAND
E-mail address: BANASZAK@math.amu.edu.pl

DEPARTMENT OF MATHEMATICS, ADAM MICKIEWICZ UNIVERSITY, POZNAŃ, POLAND
E-mail address: GAJDA@math.amu.edu.pl

DEPARTMENT OF MATHEMATICS, SZCZECIN UNIVERSITY, 70-451 SZCZECIN, POLAND
E-mail address: KRASON@uoo.univ.szczecin.pl

36-11 PIRIE DRIVE, DUNDAS, ONTARIO L9H 6X5, CANADA
E-mail address: PIOTR@icarus.math.mcmaster.ca