SOME REMARKS CONCERNING MOD- n K -THEORY

ERIC M. FRIEDLANDER AND MARK E. WALKER *

The spectral sequence predicted by A. Beilinson relating motivic cohomology to algebraic K -theory has been established for smooth quasi-projective varieties over a field (cf. [FS], [L1]). Among other properties verified, this spectral sequence has the expected multiplicative behavior (involving cup product in motivic cohomology and product in algebraic K -theory) and a good multiplicative “mod- n reduction” relating mod- n motivic cohomology to mod- n algebraic K -theory.

The purpose of this short paper is to draw two conclusions concerning mod- n algebraic K -theory of smooth, quasi-projective varieties over an algebraically closed field by comparing this spectral sequence to its “localization” which we show converges to mod- n étale K -theory. In Theorem 2.1, we verify that the map from mod- n algebraic K -theory to mod- n étale K -theory of a smooth quasi-projective variety X over an algebraically closed field in which n is invertible is surjective in degrees greater than or equal to twice the dimension of X . This refines the surjectivity result of [DFST] and extends an argument by A. Suslin in [S1].

Then, in Example 3.4, we show that certain projective smooth 3-folds X constructed by S. Bloch and H. Esnault have non-zero elements in $K_0(X, \mathbb{Z}/n)$ which are killed by multiplication by a sufficiently high power of the Bott element $\beta \in K_2(X, \mathbb{Z}/n)$. Although the existence of such examples is perhaps not surprising, there have been attempts to prove that multiplication by β is always injective.

The authors gratefully acknowledge useful conversations with H. Esnault and A. Suslin.

§1. COMPARING SPECTRAL SEQUENCES

One method for computing mod- n algebraic K -theory is to compare it to its topological analogue, the étale K -theory of [DF]. Since we shall be working over an algebraically closed field, this étale K -theory is that described in its simpler, earlier incarnation presented in [F1]. Indeed, in the case in which our base field is the complex field $\mathbb{C}$, the mod- n étale K -theory of the variety X is isomorphic to the mod- n topological K -theory of the underlying analytic space X^{an} .

We recall the following fundamental result of R. Thomason, specialized to quasi-projective varieties over an algebraically closed field.

Theorem 1.1. (cf. [T]) *Let X be a quasi-projective variety over an algebraically closed field k and let n be a positive integer invertible in k . Let $\beta \in K_2(k, \mathbb{Z}/n)$ be a choice of Bott element, an element which maps to a primitive n -th root of unity*

*Both authors were partially supported by the N.S.F. and the N.S.A.

in $_n(K_1(k)) = \mu_n(k)$. Then for any quasi-projective variety X over k there is a natural isomorphism (for $* \geq 0$)

$$K_*(X, \mathbb{Z}/n)[1/\beta] \xrightarrow{\simeq} K_*^{et}(X, \mathbb{Z}/n).$$

In the following proposition, we use a result of A. Suslin [S2] to identify the result of analogous localization in motivic cohomology. Let $H_{\mathcal{M}}^p(X, \mathbb{Z}(q))$ denote the motivic cohomology of an algebraic variety X , defined either as in [V1] or as the higher Chow group $CH^q(X, 2q - p)$. The equivalence of these definitions is proved in [V2]. Let $H_{\mathcal{M}}^*(X, \mathbb{Z}/n(q))$ denote the indicated mod- n motivic cohomology group defined, for example, as the mod- n homology of the Bloch complex of codimension q cycles.

Proposition 1.2. *Let X be a smooth quasi-projective variety over an algebraically closed field k and n a positive integer invertible in k . Let*

$$b \in H_{\mathcal{M}}^0(X, \mathbb{Z}/n(1)) = H_{et}^0(X, \mu_n)$$

be the primitive n -th root of unity used to define β . Then for any $i \geq 0$, multiplication by b

$$b \cdot (-) : H_{\mathcal{M}}^i(X, \mathbb{Z}/n(s)) \rightarrow H_{\mathcal{M}}^i(X, \mathbb{Z}/n(s+1))$$

is an isomorphism for $s \geq d = \dim(X)$ whereas multiplication by b

$$b \cdot (-) : H_{et}^i(X, \mu_n^{\otimes s}) \rightarrow H_{et}^i(X, \mu_n^{\otimes s+1})$$

is an isomorphism for all s . In other words, for any s the natural map

$$H_{\mathcal{M}}^i(X, \mathbb{Z}/n(*)) \rightarrow H_{et}^i(X, \mu_n^{\otimes *})$$

is equivalent to localization of $H^i(X, \mathbb{Z}/n())$ with respect to the action of multiplication by b .*

Proof. Suslin's comparison result [S2; 2.1],

$$H_{\mathcal{M}}^*(X, \mathbb{Z}/n(d)) \xrightarrow{\simeq} H_{et}^*(X, \mu_n^{\otimes d}),$$

is stated for a variety X over an algebraically closed field of characteristic 0. The characteristic 0 hypothesis was invoked in order to apply the techniques of [SV]. As is well known, these techniques also apply to varieties over algebraically closed field of characteristic $p > 0$ with $(n, p) = 1$, thanks to J. de Jong's result concerning the existence of smooth alterations for any singular variety [J].

On the other hand, if $s > d$, then homotopy invariance of motivic cohomology and étale cohomology implies the isomorphisms

$$H_{\mathcal{M}}^*(X, \mathbb{Z}/n(s)) \simeq H_{\mathcal{M}}^*(X \times \mathbb{A}^{s-d}, \mathbb{Z}/n(s)) \xrightarrow{\simeq} H_{et}^*(X \times \mathbb{A}^{s-d}, \mu_n^{\otimes s}) \simeq H_{et}^*(X, \mu_n^{\otimes s}).$$

Now observe that

$$b \cdot (-) : H_{et}^i(X, \mu_n^{\otimes s}) \rightarrow H_{et}^i(X, \mu_n^{\otimes s+1}),$$

is an isomorphism for all s in view of the non-canonical isomorphisms

$$\mu_n^{\otimes s} \simeq \mathbb{Z}/n \simeq \mu_n^{\otimes s+1}.$$

In particular, multiplication by b ,

$$b \cdot (-) : H_{\mathcal{M}}^i(X, \mathbb{Z}/n(s)) \rightarrow H_{\mathcal{M}}^i(X, \mathbb{Z}/n(s+1)),$$

is an isomorphism whenever $s \geq d$. $\square$

The following theorem shows how to relate multiplication by β in mod- n K -theory and multiplication by b in mod- n cohomology. The reader can find an alternate construction of such a map of spectral sequences given (implicitly) in [L2;12.13].

Theorem 1.3. *Let X be a smooth quasi-projective variety over an algebraically closed field k and n a positive integer invertible in k . Then there is a natural map of spectral sequences from*

$$E_2^{p,q} = H_{\mathcal{M}}^{p-q}(X, \mathbb{Z}/n(-q)) \Rightarrow K_{-p-q}(X, \mathbb{Z}/n)$$

to

$$'E_2^{p,q} = H_{et}^{p-q}(X, \mu_n^{\otimes -q}) \Rightarrow K_{-p-q}^{et}(X, \mathbb{Z}/n)$$

whose map on abutments is localization by the Bott element $\beta \in K_2(X, \mathbb{Z}/n)$ and whose map on E_2 -terms is localization by $b \in H_{\mathcal{M}}^0(X, \mathbb{Z}/n(1))$.

Proof. The first spectral sequence arises from a homotopy exact couple associated to a tower of fibrations

$$\dots \mathcal{K}^{(q)}(X \times \Delta^\bullet) \rightarrow \mathcal{K}^{(q-1)}(X \times \Delta^\bullet) \rightarrow \dots$$

The multiplicative structure on this exact couple as detailed in [FS] enables us to define a self-map of the exact couple associated to multiplication by β . The second spectral sequence is that of the exact couple obtained by localization of the first exact couple with respect to this map (i.e., by considering the colimit of an infinite sequence of this map), whose asserted form follows from Theorem 1.1 and Proposition 1.2. $\square$

§2. SURJECTIVITY IN MOD- n K -THEORY

In [S1], A. Suslin essentially assumed Theorem 1.3 and verified the following surjectivity result (among other things) in the special case in which X has dimension at most 2. As we see below, this dimension condition is not necessary to prove such a surjectivity result.

Theorem 2.1. *Let X be a smooth quasi-projective variety of dimension d over an algebraically closed field k and n a positive integer invertible in k . Then the natural map*

$$K_m(X, \mathbb{Z}/n) \rightarrow K_m^{et}(X, \mathbb{Z}/n),$$

is split surjective for $m \geq 2d$.

Proof. Note that the r th differentials, $d_r^{p,q}$ and $'d_r^{p,q}$, of the spectral sequences of 1.3 have bi-degree $(r, -r+1)$. As discussed in the proof of 1.2, a theorem of Suslin

[S2; 2.1] shows that the map $E_2^{p,q} \rightarrow {}'E_2^{p,q}$ is an isomorphism for $q \leq -d$. We claim that for $r \geq 2$, $p+q \leq -2d$, and $q \leq -d$, the image of the map

$$d_r^{p-r,q+r-1} : E_r^{p-r,q+r-1} \rightarrow E_r^{p,q}$$

is mapped isomorphically to the image of

$$'d_r^{p-r,q+r-1} : {}'E_r^{p-r,q+r-1} \rightarrow {}'E_r^{p,q}$$

under the map $E_r^{p,q} \rightarrow {}'E_r^{p,q}$, which is itself an isomorphism. To see this, consider the commutative square

$$\begin{array}{ccc} E_r^{p-r,q+r-1} \equiv H_{\mathcal{M}}^{p-2r-q+1}(X, \mathbb{Z}/n(-q-r+1)) & \xrightarrow{d_r} & E_r^{p,q} \equiv H_{\mathcal{M}}^{p-q}(X, \mathbb{Z}/n(-q)) \\ \downarrow & & \downarrow \\ {}'E_r^{p-r,q+r-1} \equiv H_{et}^{p-2r-q+1}(X, \mu_n^{\otimes(-q-r+1)}) & \xrightarrow{{}'d_r} & {}'E_r^{p,q} \equiv H_{et}^{p-q}(X, \mu_n^{\otimes(-q)}). \end{array}$$

If $q+r-1 \leq -d$, then both vertical maps of this diagram are seen to be isomorphisms by induction on $r \geq 2$, with the case $r = 2$ given by Suslin's theorem [S2; 2.1]. Thus the result is evident in this case. If $q+r-1 > -d$, then the lower left group is 0 for all $r \geq 2$, since $p-2r-q+1 < 0$, and so $'d_r = 0$. As before, by induction on $r \geq 2$ and using Suslin's theorem, one easily concludes that d_r is the zero map and the right-hand vertical map of the diagram is an isomorphism.

Letting $r \mapsto \infty$, it follows that for $p+q \leq -2d$, the map

$$E_{\infty}^{p,q} \rightarrow {}'E_{\infty}^{p,q}$$

is an isomorphism for $q \leq -d$ and has target 0 for $q > -d$. Thus, for any $m = -p-q \geq 2d$, the map

$$K_m(X, \mathbb{Z}/n) \rightarrow K_m^{et}(X, \mathbb{Z}/n)$$

admits a decreasing bounded filtration

$$F^s \equiv F^s K_m(X, \mathbb{Z}/n) \rightarrow {}'F^s \equiv {}'F^s K_m^{et}(X, \mathbb{Z}/n), \quad s \in \mathbb{Z},$$

such that $F^s/F^{s+1} \cong {}'F^s/{}'F^{s+1}$ for $s \geq d-m$ and $'F^s/{}'F^{s+1} = 0$ for $s < d-m$. Thus, the composite map $F^{d-m} \rightarrow {}'F^{d-m} \rightarrow K_m^{et}(X, \mathbb{Z}/n)$ is an isomorphism, whose inverse composed with $F^{d-m} \hookrightarrow K_m(X, \mathbb{Z}/n)$ is the desired splitting. $\square$

Using Theorem 2.1, we obtain further information concerning the map of spectral sequences of Theorem 1.3.

Proposition 2.2. *Adopt the hypotheses and notation of Theorem 1.3 with $d = \dim(X)$. If $(n, (d-1)!) = 1$, then there exists a commutative diagram*

$$\begin{array}{ccc} K_0(X, \mathbb{Z}/n) & \xrightarrow{ch} & \bigoplus_{s \geq 0} H_{\mathcal{M}}^{2s}(X, \mathbb{Z}/n(s)) \\ \downarrow & & \downarrow \\ K_0^{et}(X, \mathbb{Z}/n) & \xrightarrow{{}'ch} & \bigoplus_{s \geq 0} H_{et}^{2s}(X, \mu_n^{\otimes s}) \end{array}$$

whose horizontal arrows are isomorphisms, given as the reduction modulo n of the Chern character.

Proof. Explicit formulas for the Chern character in terms of (integrally defined) Chern classes enable us to conclude that ch and $'ch$ can be defined in terms of mod- n Chern classes provided $(n, (d-1)!) = 1$.

The only possible $r \geq 2$ such that d_r or $'d_r$ maps non-trivially to $E_r^{s,-s}$ or $'E_r^{s,-s}$ satisfy $r < s$. For if $r > s$, then the source of each of these maps is trivial, and if $r = s$, the source of d_r and $'d_r$ are each isomorphic to $H_{Zar}^1(X, \mu_n) \cong H_{et}^1(X, \mu_n)$, which admits a surjective map from $K_1(X, \mathbb{Z}/n)$ and thus must persist to infinity. Now observe that $E_2^{s,-s} \cong 'E_2^{s,-s} = H_{et}^{2s}(X, \mu_n^{\otimes s}) = 0$ whenever $2s > 2d$, by Suslin's theorem and since X has étale cohomological dimension of at most $2d$. Thus $E_r^{s,-s} = 'E_r^{s,-s} = 0$ for all $r \geq 2$ whenever $2s > 2d$. Since we may assume $r < s$ from above, the only non-trivial differentials $d_r, 'd_r$ with target $E_r^{s,-s}, 'E_r^{s,-s}$ satisfy $r < d$.

As shown in [L2;9.7], the spectral sequences of Theorem 1.3 admit Adams operations ψ^k commuting with differentials whose action on each of $E_r^{p,q}$ and $'E_r^{p,q}$ is multiplication by k^{-q} . As argued in [L2;11.7], this implies that the differentials d_r and $'d_r$ vanish provided that $n > r$.

In particular, since $n > d-1$, this shows that all differentials with target $E_r^{s,-s}$ or $'E_r^{s,-s}$ vanish. On the other hand, the only non-vanishing differentials with source $E_r^{s,-s}$ or $'E_r^{s,-s}$ must satisfy $r \geq n \geq d$. But we claim that even these differentials vanish. For if $s > 0$, then their targets, $E_r^{s+r,-s-r+1}$ and $'E_r^{s+r,-s-r+1}$, are zero, since $'E_r^{p,q} = 0$ for $p-q > 2d$ by considering étale cohomological dimension, and $E_r^{p,q} = 0$ for $p-q > -q+d$, since the motivic complex $\mathbb{Z}(-q)$ is concentrated in cohomological degrees $[-\infty, -q]$ (cf. [V1]) and X has Zariski cohomological dimension d . If $s = 0$, then $E_r^{0,0}$ and $'E_r^{0,0}$ clearly persist to infinity, and so d_r and $'d_r$ vanish in this case too.

Thus, the mod- n spectral sequences collapse at the E_2 -level in total degree 0, with splitting given by the Chern character maps ch and $'ch$. $\square$

§3. NON-INJECTIVITY OF MULTIPLICATION BY β

In this section, we show how examples of algebraic cycles which are neither n -torsion nor n -divisible and which are homologically trivial determine examples of classes in mod- n K -theory which are killed by multiplication by a sufficiently high power of the Bott element. This follows easily from Proposition 2.2.

The connection between homologically trivial cycles and elements in K -theory killed by multiplication by a power of the Bott element was first observed when contemplating the relationship between “algebraic vector bundles modulo algebraic equivalence” and “algebraic cycles modulo algebraic equivalence” as in [FW]. As we observe in Remark 3.3 below, one should be able to use a somewhat sophisticated version of this relationship to provide much sharper examples.

Our examples of non-surjectivity in Example 3.2 use the following proposition relating cycles homologically trivial to K -theory classes killed by a power of the Bott element. We remind the reader that for any smooth quasi-projective variety X , we have

$$H_{\mathcal{M}}^{2s}(X, \mathbb{Z}/n(s)) \simeq CH^s(X) \otimes \mathbb{Z}/n,$$

where $CH^s(X)$ denotes the Chow group of codimension s cycles on X modulo rational equivalence.

Proposition 3.1. *Let X be a smooth quasi-projective variety over an algebraically closed field k , n a positive integer invertible in k , and $d = \dim(X)$. Assume that there exists an algebraic cycle Z of codimension s on X with the property that*

$$[Z] \neq 0 \in CH^s(X) \otimes \mathbb{Z}/n, \quad [Z] = 0 \in H_{et}^{2s}(X, \mu_n^{\otimes s}). \quad (3.1.1)$$

Then there exists

$$0 \neq \alpha \in K_0(X, \mathbb{Z}/n), \quad \beta^e \cdot \alpha = 0$$

for some $e > 0$ provided that $(n, (d-1)!) = 1$.

Proof. By Proposition 2.2, the hypotheses of the proposition imply that there exists some non-zero class $\alpha \in K_0(X, \mathbb{Z}/n)$ mapping to $0 \in K_0^{et}(X, \mathbb{Z}/n)$. Thus, the proposition follows from Theorem 1.1. $\square$

Examples of varieties X and cycles Z satisfying the hypotheses of Proposition 3.1 should be “generic” in nature. In the following, we recall an explicit such example constructed by S. Bloch and H. Esnault.

Example 3.2. *We let $W \subset \mathbb{P}_{\mathbb{C}}^5$ be a smooth complex quadric hypersurface defined over $\mathbb{Q}$. For a prime $\ell \gg 0$ and integer $k \gg 0$, Bloch and Esnault show [BE; 4.1] that there exists a smooth hypersurface section X of W defined over an algebraic number field and an algebraic cycle Z of codimension 2 on X which satisfy (3.1.1) with $s = 2$, $n = \ell^k$:*

$$[Z] \neq 0 \in CH^2(X, \mathbb{Z}/\ell^k), \quad [Z] = 0 \in H_{et}^4(X, \mu_{\ell^k}^{\otimes 2}).$$

Thus, for such choices of X , ℓ , and k , Proposition 3.1 tells us that there exists

$$0 \neq \alpha \in K_0(X, \mathbb{Z}/\ell^k), \quad \beta^e \cdot \alpha = 0,$$

for some $e > 0$.

Remark 3.3. *As mentioned above, the relevance of the Bloch-Esnault examples to the non-injectivity of multiplication by the Bott element was motivated by considering the relationship between “semi-topological K -theory” (cf. [FW]) and “morphic cohomology” (cf. [FL]). In [CL], it is stated that these are rationally isomorphic for projective, smooth complex varieties. Moreover, it is shown in [CL] that multiplication by the Bott element corresponds under the Chern character to the s -operation in morphic cohomology. In [F2], examples are given of classes in morphic cohomology killed by a specific power of the s -operation. Thus, provided that one could show that the cycles occurring in these examples are not infinitely ℓ -divisible in the appropriate mod- ℓ Chow group, one could provide much sharper examples than those of Example 3.2. Namely, for $m > 1$ and suitable prime ℓ , one should be able to construct a projective smooth complex algebraic variety X and an element $\alpha \in K_0(X, \mathbb{Z}/\ell)$ such that*

$$\beta^{m-1} \cdot \alpha \neq 0, \quad \beta^m \cdot \alpha = 0.$$

REFERENCES

- [BE] S. Bloch and H. Esnault, *The coniveau filtration and non-divisibility for algebraic cycles*, Math. Ann **304** (1996), 303 - 314.
- [CL] R. Cohen and P. Lima-Filho, *Holomorphic K -theory, algebraic co-cycles, and loop groups*, Preprint, available at <http://www.math.uiuc.edu/K-theory>.
- [DF] W. Dwyer and E. Friedlander, *Algebraic and étale K -theory*, Trans. A.M.S. **292** (1985), 247-280.
- [DFST] W. Dwyer, E. Friedlander, V. Snaith, and R. Thomason, *Algebraic K -theory eventually surjects onto topological K -theory*, Inventiones Math **66** (1982), 481-497.
- [F1] E. Friedlander, *Étale K -theory II: Connections with algebraic K -theory*, Ann. Scient. Ecole Norml. Sup. **15** (1982), 231-256.
- [F2] E. Friedlander, *Relative Chow correspondences and the Griffiths group*, To appear in Annales de l'Institut Fourier.
- [FL] E. Friedlander and H.B. Lawson, Jr., *A theory of algebraic cocycles*, Annals of Math **136** (1992), 361-428.
- [FS] E. Friedlander and A. Suslin, *The spectral sequence relating algebraic K -theory to motivic cohomology*, Preprint, available at <http://www.math.uiuc.edu/K-theory>.
- [FW] E. Friedlander and M. Walker, *Semi-topological K -theory using function complexes*, Preprint, available at <http://www.math.uiuc.edu/K-theory>.
- [J] A.J. de Jong, *Smoothness, semi-stability, and alterations*, Pub. I.H.E.S. (1996), 51-93.
- [L1] M. Levine, *Techniques of localization in the theory of algebraic cycles*, Preprint, available at <http://www.math.uiuc.edu/K-theory>.
- [L2] M. Levine, *K -theory and motivic cohomology of schemes*, Preprint, available at <http://www.math.uiuc.edu/K-theory>.
- [S1] A. Suslin, *Algebraic K -theory and motivic cohomology*, Proc. 1994 I. C. M. Zurich, 342-351.
- [S2] A. Suslin, *Higher Chow groups and étale cohomology*, Cycles, Transfers and Motivic Homology Theories (V. Voevodsky, A. Suslin and E. Friedlander, eds.), Annals of Math. Studies, 2000, pp. 237—252..
- [SV] A. Suslin and V. Voevodsky, *Singular homology of abstract algebraic varieties*, Inventiones Math **123** (1996), 61-94.
- [T] R. Thomason, *Algebraic K -theory and étale cohomology*, Ann. Sci. Ecole Norm. Sup. **18** (1985), 437-552.
- [V1] V. Voevodsky, *The Milnor Conjecture*, Preprint, available at <http://www.math.uiuc.edu/K-theory>.
- [V2] V. Voevodsky, *Motivic cohomology are isomorphic to higher Chow groups*, Preprint, available at <http://www.math.uiuc.edu/K-theory>.

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, IL, 60208, U.S.A.
E-mail address: eric@math.nwu.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEBRASKA – LINCOLN,
 LINCOLN, NE, 68588-0323, U.S.A.
E-mail address: mwalker@math.unl.edu