

A counterexample to generalizations of the Milnor-Bloch-Kato conjecture*

By Michael Spieß and Takao Yamazaki

Abstract

We construct an example of a torus T over a field K for which the Galois symbol $K(K; T, T)/nK(K; T, T) \rightarrow H^2(K, T[n] \otimes T[n])$ is not injective for some n . Here $K(K; T, T)$ is the Milnor K -group attached to T introduced by Somekawa. We show also that the motive $M(T \times T)$ gives a counterexample to another generalization of the Milnor-Bloch-Kato conjecture (proposed by Beilinson).

1 Introduction

Let K be a field, m a positive integer and n an integer prime to the characteristic of K . The Milnor-Bloch-Kato conjecture asserts that the Galois symbol

$$(1) \quad K_m^M(K)/nK_m^M(K) \longrightarrow H^m(K, \mathbb{Z}/n\mathbb{Z}(m))$$

from Milnor K -groups to Galois cohomology is bijective. Recently, Rost and Voevodsky have announced a proof (special cases have been obtained earlier by Merkurjev-Suslin, Rost and Voevodsky).

In [So], Somekawa has introduced certain *generalized Milnor K -groups* $K(K; A_1, \dots, A_m)$ attached to semi-abelian varieties A_1, \dots, A_m . If $A_1 = \dots = A_m = \mathbb{G}_m$ is the one-dimensional split torus they agree with the usual $K_m^M(K)$. If $m = 2$, $A_1 = \text{Jac}_X$ and $A_2 = \text{Jac}_Y$ are the Jacobians of smooth, projective and connected curves X and Y over K having a K -rational point, then $K(K; A_1, A_2)$ is the kernel of the Albanese map $\text{CH}_0(X \times Y)_{\text{deg}=0} \rightarrow \text{Alb}_{X \times Y}(K)$.

Somekawa has defined a Galois symbol

$$(2) \quad K(K; A_1, \dots, A_m)/nK(K; A_1, \dots, A_m) \longrightarrow H^m(K, A_1[n] \otimes \dots \otimes A_m[n])$$

*This work was done during the second author stayed at Universität Bielefeld supported by SFB 701. He is grateful to the members there.

and conjectured that it is always injective. In this note we present a counterexample (see section 2). Let us describe it briefly. Let L/K be a cyclic extension of degree n and σ a generator of the Galois group $\text{Gal}(L/K)$. Let T be the kernel of the norm map $\text{Res}_{L/K} \mathbb{G}_m \rightarrow \mathbb{G}_m$. We show that the norm $K(L; T, T) \rightarrow K(K; T, T)$ induces an isomorphism $K_2(L; T, T)/(1 - \sigma) \rightarrow K_2(K; T, T)$. On the other hand, the corresponding map of Galois cohomology groups $H^2(L, T[n] \otimes T[n])/(1 - \sigma) \rightarrow H^2(K, T[n] \otimes T[n])$ is neither injective nor surjective (for a suitable choice of L/K). Note that, since T is split over L , the Galois symbol $K_2(L; T, T)/n \rightarrow H^2(L, T[n] \otimes T[n])$ is bijective. Consequently, $K_2(K; T, T)/n \rightarrow H^2(K, T[n] \otimes T[n])$ is in general not injective.

In the section 3 we show that the motive $M(T \times T)$ gives a counterexample to another generalization of the Milnor-Bloch-Kato conjecture (proposed by Beilinson).

We would like to thank Bruno Kahn for his comments on the first version of this note. In particular he pointed out to us that our counterexample to Somekawa's conjecture should also provide a counterexample to Beilinson's conjecture.

2 Counterexample to Somekawa's conjecture

Algebraic groups as Mackey-functors Let K be a field. For a finite field extension L/K and commutative algebraic groups G over K and H over L we denote by G_L the base change of G to L and by $\text{Res}_{L/K} H$ the Weil restriction of H . The functor $G \mapsto G_L$ is left and right adjoint to $H \mapsto \text{Res}_{L/K} H$. In particular there are adjunction homomorphisms $\iota_{L/K} : G \rightarrow \text{Res}_{L/K} G_L$ and $N_{L/K} : \text{Res}_{L/K} G_L \rightarrow G$. When L/K is a Galois extension, the Galois group $\text{Gal}(L/K)$ acts canonically on $\text{Res}_{L/K} G_L$. The following simple result, whose proof will be left to the reader, will be used later.

Lemma 1 *Let L/K be a cyclic Galois extension of degree n , σ a generator of $\text{Gal}(L/K)$ and let G be a commutative algebraic group over K . Let G' be the kernel of $N_{L/K} : \text{Res}_{L/K} G_L \rightarrow G$ so that $G'_L \cong G_L^{n-1}$. Then the map*

$$\text{Res}_{L/K}(G_L)^{n-1} \cong \text{Res}_{L/K} G'_L \xrightarrow{N_{L/K}} G' \hookrightarrow \text{Res}_{L/K} G_L$$

is given on the i -th summand by $1 - \sigma^i$.

We denote by \mathcal{C}_K the category of finite reduced K -schemes. Thus each object of \mathcal{C}_K is isomorphic to $\text{Spec}(E_1 \times \dots \times E_r)$ where $E_1, \dots, E_r/K$ are finite field extensions. A commutative algebraic group G over K defines

a *Mackey-functor*, i.e. a co- and contravariant functor $G : \mathcal{C}_K \rightarrow \text{Mod}_{\mathbb{Z}}$ satisfying (i), (ii) below. If $f : X \rightarrow Y$ is a morphism we denote by $f_* : G(X) \rightarrow G(Y)$ and $f^* : G(Y) \rightarrow G(X)$ the homomorphisms induced by co- and contravariant functoriality respectively.

(i) If $X = X_1 \coprod X_2 \in \text{Obj}(\mathcal{C}_K)$ then $G(X) = G(X_1) \oplus G(X_2)$.

(ii) If

$$\begin{array}{ccc} X' & \longrightarrow & Y' \\ & \searrow f' & \\ \downarrow g' & & \downarrow g \\ X & \longrightarrow & Y \\ & \searrow f & \end{array}$$

is a cartesian square in \mathcal{C}_K then $g^* f_* = (f')_*(g')^*$.

If $K \subseteq E_1 \subseteq E_2$ are two finite field extensions and $f : \text{Spec } E_2 \rightarrow \text{Spec } E_1$ the corresponding map in \mathcal{C}_K then f^* (resp. f_*) is given by $\iota_{E_2/E_1} : G(E_1) \rightarrow G(E_2)$ (resp. $N_{E_2/E_1} : G(E_2) \rightarrow G(E_1)$).

Local symbols We recall also the notion of a *local symbol* ([Se] and [So]) for G . Let $X \rightarrow \text{Spec } K$ is a proper non-singular algebraic curve (note that we do not assume that X is connected). Let $K(X)$ denote the ring of rational functions on X and $|X|$ the set of closed points of X . For $P \in |X|$ we denote by K_P the quotient field of the completion $\widehat{\mathcal{O}}_{X,P}$ of $\mathcal{O}_{X,P}$, by $v_P : K_P \rightarrow \mathbb{Z} \cup \{\infty\}$ the normalized valuation and by $K(P)$ the residue field of K_P . The local symbol at P is a homomorphism $\partial_P : (K_P)^* \otimes G(K_P) \rightarrow G(K(P))$. It is characterized by the following properties:

- (i) If $f \in (K_P)^*$ and $g \in G(\widehat{\mathcal{O}}_{X,P})$ then $\partial_P(f \otimes g) = v_P(f)g(P)$. Here $g(P)$ is the image of g under the canonical map $G(\widehat{\mathcal{O}}_{X,P}) \rightarrow G(K(P))$.
- (ii) For $f \in K(X)^*$ and $g \in G(K(X))$ we have $\sum_{P \in |X|} N_{K(P)/K}(\partial_P(f \otimes g)) = 0$.

Milnor K -groups attached to commutative algebraic groups Let G_1, \dots, G_m be commutative algebraic groups over K . In [So] Somekawa has introduced the Milnor K -group $K(K; G_1, \dots, G_m)$ (actually Somekawa considered only the case of semiabelian varieties though his construction works for arbitrary commutative algebraic groups). It is given as

$$K(K; G_1, \dots, G_m) = \left(\bigoplus_X G_1(X) \otimes \dots \otimes G_m(X) \right) / R$$

where X runs through all objects of \mathcal{C}_K and where the subgroup R is generated by the following elements:

(R1) If $f : X \rightarrow Y$ is a morphism in \mathcal{C}_K and if $x_{i_0} \in G_{i_0}(Y)$ for some i_0 and $x_i \in G_i(X)$ for $i \neq i_0$, then

$$x_1 \otimes \dots \otimes f_*(x_{i_0}) \otimes \dots \otimes x_m - f^*(x_1) \otimes \dots \otimes x_{i_0} \otimes \dots \otimes f^*(x_m) \in R.$$

(R2) Let $X \rightarrow \text{Spec } K$ be a proper non-singular curve, $f \in K(X)^*$ and $g_i \in G_i(K(X))$. Assume that for each $P \in |X|$ there exists $i(P)$ such that $g_i \in G_i(\hat{\mathcal{O}}_{X,P})$ for all $i \neq i(P)$. Then

$$\sum_{P \in |X|} g_1(P) \otimes \dots \otimes \partial_P(f \otimes g_{i(P)}) \otimes \dots \otimes g_m(P) \in R.$$

For $X \in \mathcal{C}_K$ and $x_i \in G_i(X)$ for $i = 1, \dots, m$ we write $\{x_1, \dots, x_m\}_{X/K}$ for the image of $x_1 \otimes \dots \otimes x_m$ in $K(K; G_1, \dots, G_m)$ (elements of this form will be referred to as *symbols*).

A sequence of algebraic groups $G' \rightarrow G \rightarrow G''$ over K will be called *Zariski exact* if $G'(E) \rightarrow G(E) \rightarrow G''(E)$ is exact for every extension E/K . The proof of the following result is straightforward; hence will be omitted.

Lemma 2 *Let m be a positive integer and let $i \in \{1, \dots, m\}$. Let G_1, \dots, G_m be commutative algebraic groups over K and let $G'_i \rightarrow G_i \rightarrow G''_i \rightarrow 1$ be a Zariski exact sequence of commutative algebraic groups over K . Then the sequence*

$$K(K; G_1, \dots, G'_i, \dots) \rightarrow K(K; G_1, \dots, G_i, \dots) \rightarrow K(K; G_1, \dots, G''_i, \dots) \rightarrow 0$$

is exact as well.

The norm map Let G_1, \dots, G_m be commutative algebraic groups over K and let L/K be a finite extension. Set $K(L; G_1, \dots, G_m) := K(L; (G_1)_L, \dots, (G_m)_L)$. Then we have the norm map [So]

$$(3) \quad N_{L/K} : K(L; G_1, \dots, G_m) \longrightarrow K(K; G_1, \dots, G_m)$$

defined on symbols by $N_{L/K}(\{x_1, \dots, x_m\}_{X/L}) = \{x_1, \dots, x_m\}_{X/K}$ for any $X \in \mathcal{C}_L$ and $x_i \in G_i(X)$ ($i = 1, \dots, m$). We give another interpretation of (3) below when L/K is separable. It is based on the following result.

Lemma 3 *Let L/K be a finite separable extension and let i, m be positive integers with $i \leq m$. Let $G_1, \dots, G_{i-1}, G_{i+1}, \dots, G_m$ be commutative algebraic groups over K and let G_i be a commutative algebraic group over L . Then, we have an isomorphism*

$$K(K; G_1, \dots, \text{Res}_{L/K} G_i, \dots, G_m) \cong K(L; (G_1)_L, \dots, G_i, \dots, (G_m)_L).$$

Proof. To simplify the notation we assume that $i = m$. We denote by $\pi^{-1} : \mathcal{C}_K \rightarrow \mathcal{C}_L$ and $\pi : \mathcal{C}_L \rightarrow \mathcal{C}_K$ the functors

$$\pi^{-1}(X \rightarrow \text{Spec } K) := (X \otimes_K L \rightarrow \text{Spec } L),$$

$$\pi(Y \rightarrow \text{Spec } L) := (Y \rightarrow \text{Spec } L \rightarrow \text{Spec } K).$$

π is left adjoint to π^{-1} . For $X \in \mathcal{C}_K$ and $Y \in \mathcal{C}_L$ let

$$p_X : X \otimes_K L \longrightarrow X, \quad \iota_Y : Y \longrightarrow Y \otimes_K L.$$

be the adjunction morphisms. We define homomorphisms

$$\phi : K(K; G_1, \dots, G_{m-1}, \text{Res}_{L/K} G_m) \longrightarrow K(K; (G_1)_L, \dots, (G_{m-1})_L, G_m),$$

$$\psi : K(L; (G_1)_L, \dots, (G_{m-1})_L, G_m) \longrightarrow K(K; G_1, \dots, G_{m-1}, \text{Res}_{L/K} G_m).$$

as follows. For $X \in \mathcal{C}_K$, $x_1 \in G_1(X), \dots, x_{m-1} \in G_{m-1}(X)$ and $x_m \in G_m(X \otimes_K L)$ we put

$$\phi(\{x_1, \dots, x_m\}_{X/K}) = \{p^*(x_1), \dots, p^*(x_{m-1}), x_m\}_{(X \otimes_K L)/L}.$$

Conversely, for $Y \in \mathcal{C}_L$ and $y_1 \in G_1(Y), \dots, y_m \in G_m(Y)$ let

$$\psi(\{y_1, \dots, y_{m-1}, y_m\}_{Y/L}) = \{y_1, \dots, y_{m-1}, \iota_*(y_m)\}_{Y/K}.$$

One can easily verify that these maps are well-defined and mutually inverse to each other. \square

Let G_1, \dots, G_m be commutative algebraic groups over K and let L/K be a finite separable extension. Take any $i \in \{1, \dots, m\}$. The map $N_{L/K} : \text{Res}_{L/K}(G_i)_L \rightarrow G_i$ induces a map $K(K; G_1, \dots, \text{Res}_{L/K}(G_i)_L, \dots, G_m) \longrightarrow K(K; G_1, \dots, G_m)$, and the composition of it with the isomorphism ψ above coincides with the norm map (3). When L/K is a Galois extension, the action of $\text{Gal}(L/K)$ on $\text{Res}_{L/K}(G_i)_L$ induces its action on

$$K(L; G_1, \dots, G_m) \cong K(K; G_1, \dots, \text{Res}_{L/K}(G_i)_L, \dots, G_m)$$

and we have $N_{L/K} \circ \sigma = N_{L/K}$ for all $\sigma \in \text{Gal}(L/K)$. This action does not depend on the choice of i .

Lemma 4 *Let L/K be a cyclic Galois extension and let $\sigma \in \text{Gal}(L/K)$ be a generator. Suppose that for two different $i \in \{1, \dots, m\}$ the sequence*

$$(4) \quad \text{Res}_{L/K}(G_i)_K \xrightarrow{N_{L/K}} G_i \longrightarrow 1$$

is Zariski exact. Then the sequence of abelian groups

$$K(L; G_1, \dots, G_m) \xrightarrow{1-\sigma} K(L; G_1, \dots, G_m) \xrightarrow{N_{L/K}} K(K; G_1, \dots, G_m) \longrightarrow 0$$

is exact.

Proof. Suppose that (4) is exact for $i = m - 1, m$. Let $G'_m := \text{Ker}(N_{L/K} : \text{Res}_{L/K}(G_m)_L \rightarrow G_m)$. By Lemmas 2 and 3 there are exact sequences

$$(5) \quad K(K; G_1, \dots, G'_m) \rightarrow K(L; G_1, \dots, G_m) \xrightarrow{N_{L/K}} K(K; G_1, \dots, G_m) \rightarrow 0$$

$$(6) \quad K(L; G_1, \dots, G_{m-1}, G'_m) \xrightarrow{N_{L/K}} K(K; G_1, \dots, G_{m-1}, G'_m) \rightarrow 0.$$

Since $(G'_m)_L \cong (G_m)_L^{n-1}$ ($n := [L : K]$) we can replace the first group of (6) by $K(L; G_1, \dots, G_m)^{n-1}$. By Lemma 1 the composite

$$K(L; G_1, \dots, G_m)^{n-1} \rightarrow K(K; G_1, \dots, G_{m-1}, G'_m) \rightarrow K(L; G_1, \dots, G_m)$$

is given on the i -th summand by $1 - \sigma^i$. The assertion follows. \square

Galois symbol Let G_1, \dots, G_m be connected commutative algebraic groups over K , and let n be an integer prime to the characteristic of K . For any finite extension L/K , we have a homomorphism [So]

$$(7) \quad h_L : K(L; G_1, \dots, G_m)/n \longrightarrow H^m(L, G_1[n] \otimes \dots \otimes G_m[n])$$

called the *Galois symbol*. This is characterized by the following properties.

- (i) If $x_i \in G_i(L)$ for $i = 1, \dots, m$, then $h_L(\{x_1, \dots, x_m\}_{L/L}) = (x_1) \cup \dots \cup (x_m)$. Here we write by (x_i) for the image of x_i in $H^1(L, G_i[n])$ by the connecting homomorphism associated to the exact sequence $1 \rightarrow G_i[n] \rightarrow G_i \xrightarrow{n} G_i \rightarrow 1$.
- (ii) If $M/L/K$ is a tower of finite extensions and if M/L is separable (resp. purely inseparable), then the diagram

$$\begin{array}{ccc} K(M; G_1, \dots, G_m)/n & \xrightarrow{h_M} & H^m(M, G_1[n] \otimes \dots \otimes G_m[n]) \\ \downarrow N_{M/L} & & \downarrow \\ K(L; G_1, \dots, G_m)/n & \xrightarrow{h_L} & H^m(L, G_1[n] \otimes \dots \otimes G_m[n]) \end{array}$$

is commutative, where the right vertical map is the corestriction (resp. the multiplication by $[M : L]$ under the identification $H^m(M, G_1[n] \otimes \dots \otimes G_m[n]) \cong H^m(L, G_1[n] \otimes \dots \otimes G_m[n])$).

Property (i) implies in particular that (7) coincides with the usual Galois symbol (1) in the case $G_1 = \dots = G_m = \mathbb{G}_m$. In [So] Remark 1.7, Somekawa conjectured that the Galois symbol associated to semiabelian varieties should be injective.

Galois cohomology of cyclic extensions Let L/K be a cyclic Galois extension of degree n and let σ be a generator of $G := \text{Gal}(L/K)$. For a discrete G_K -module M , tensoring the short exact sequence of G -modules

$$(8) \quad 0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Z}[G] \xrightarrow{1-\sigma} \mathbb{Z}[G] \longrightarrow \mathbb{Z} \longrightarrow 0$$

with M yields a distinguished triangle

$$(9) \quad M[1] \xrightarrow{\alpha} C^\cdot(M) \xrightarrow{\beta} M \xrightarrow{\gamma} M[2]$$

in the derived category $D(G_K)$. Here we denote by $C^\cdot(M)$ the complex

$$\text{Res}_{L/K} M \xrightarrow{1-\sigma} \text{Res}_{L/K} M$$

concentrated in degree -1 and 0 . The spectral sequence

$$E_1^{p,q} = H^q(K, C^p(M)) \implies E^{p+q} = H^{p+q}(K, C^\cdot(M))$$

induces short exact sequences

$$(10) \quad 0 \rightarrow H^q(L, M)_G \rightarrow H^q(K, C^\cdot(M)) \rightarrow H^{q+1}(L, M)^G \rightarrow 0.$$

It is easy to see that the composite

$$H^{q+1}(K, M) \xrightarrow{\alpha} H^q(K, C^\cdot(M)) \rightarrow H^{q+1}(L, M)^G$$

is the restriction and

$$H^q(L, M)_G \rightarrow H^q(K, C^\cdot(M)) \xrightarrow{\beta} H^q(K, M)$$

is induced by the corestriction. In particular we have $\gamma(H^q(K, M)) \subseteq \text{Ker}(\text{res} : H^{q+2}(K, M) \rightarrow H^{q+2}(L, M))$ hence

$$(11) \quad n\gamma(H^q(K, M)) = 0.$$

For an integer m prime to $\text{char } K$ and $r \in \mathbb{N}$ we write $\mathbb{Z}/m\mathbb{Z}(r) := \mu_m^{\otimes r}$ and

$$H^3(L/K, \mathbb{Z}/m\mathbb{Z}(2)) := \text{Ker}(H^3(K, \mathbb{Z}/m\mathbb{Z}(2)) \xrightarrow{\text{res}} H^3(L, \mathbb{Z}/m\mathbb{Z}(2))).$$

By restricting $\alpha : H^3(K, \mathbb{Z}/m\mathbb{Z}(2)) \rightarrow H^2(K, C^\cdot(\mathbb{Z}/m\mathbb{Z}(2)))$ to the subgroup $H^3(L/K, \mathbb{Z}/m\mathbb{Z}(2))$ and composing it with the inverse of the first map in (10) we obtain a map

$$(12) \quad H^3(L/K, \mathbb{Z}/m\mathbb{Z}(2)) \rightarrow \text{Ker}(H^2(L, \mathbb{Z}/m\mathbb{Z}(2))_G \xrightarrow{\text{cor}} H^2(K, \mathbb{Z}/m\mathbb{Z}(2))).$$

Lemma 5 *Assume that n is prime to $\text{char } K$ and $\mu_{n^2}(\bar{K}) \subset K$. Then the homomorphism (12) is injective for $m = n$.*

Proof. It is enough to show that $\gamma : H^1(K, \mathbb{Z}/n\mathbb{Z}(2)) \rightarrow H^3(K, \mathbb{Z}/n\mathbb{Z}(2))$ is zero. Consider the commutative diagram

$$\begin{array}{ccc} H^1(K, \mathbb{Z}/n\mathbb{Z}(2)) & \longrightarrow & H^1(K, \mathbb{Z}/n^2\mathbb{Z}(2)) \\ \downarrow \gamma & & \downarrow \gamma \\ H^3(K, \mathbb{Z}/n\mathbb{Z}(2)) & \longrightarrow & H^3(K, \mathbb{Z}/n^2\mathbb{Z}(2)) \end{array}$$

induced by the canonical injection $\mathbb{Z}/n\mathbb{Z}(2) \rightarrow \mathbb{Z}/n^2\mathbb{Z}(2)$. The assumption $\mu_{n^2}(\overline{K}) \subset K$ implies that the upper horizontal map can be identified with

$$K^*/(K^*)^n \longrightarrow K^*/(K^*)^{n^2}, x(K^*)^n \mapsto x^n(K^*)^{n^2}.$$

In particular the image is contained in $nH^1(K, \mathbb{Z}/n^2\mathbb{Z}(2))$. By (11) it is mapped under γ to $n\gamma(H^1(K, \mathbb{Z}/n^2\mathbb{Z}(2))) = 0$. On the other hand it is a simple consequence of the Merkurjev-Suslin theorem [MS] that the lower horizontal map is injective. Hence $\gamma(H^1(K, \mathbb{Z}/n\mathbb{Z}(2))) = 0$. \square

The counterexample Let L/K be as in the last section and let $T := \text{Ker}(N_{L/K} : \text{Res}_{L/K} \mathbb{G}_m \rightarrow \mathbb{G}_m)$. We make the following assumptions

- (13) n is prime to $\text{char } K$ and $\mu_{n^2}(\overline{K}) \subset K$,
- (14) $H^3(L/K, \mathbb{Z}/n\mathbb{Z}(2)) \neq 0$.

Proposition 6 *The Galois symbol $K(K; T, T)/n \rightarrow H^2(K, T[n] \otimes T[n])$ is not injective.*

Proof. Let σ be a generator of $G := \text{Gal}(L/K)$. The exact sequence

$$1 \longrightarrow \mathbb{G}_m \longrightarrow \text{Res}_{L/K} \mathbb{G}_m \xrightarrow{1-\sigma} \text{Res}_{L/K} \mathbb{G}_m \xrightarrow{N_{L/K}} \mathbb{G}_m \longrightarrow 1$$

yields two short exact sequences

- (15) $1 \longrightarrow \mathbb{G}_m \longrightarrow \text{Res}_{L/K} \mathbb{G}_m \longrightarrow T \longrightarrow 1$,
- (16) $1 \longrightarrow T \longrightarrow \text{Res}_{L/K} \mathbb{G}_m \longrightarrow \mathbb{G}_m \longrightarrow 1$.

Correspondingly, (8) induces two short exact sequences

$$(17) \quad 0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}[G] \rightarrow X \rightarrow 0, \quad 0 \rightarrow X \rightarrow \mathbb{Z}[G] \rightarrow \mathbb{Z} \rightarrow 0$$

where X denotes the cocharacter group of T . Note that the sequence (15) is Zariski exact by Hilbert 90. Since the map $\text{Res}_{L/K} \mathbb{G}_m \rightarrow T$ factors through

$\text{Res}_{L/K} \mathbb{G}_m \rightarrow \text{Res}_{L/K} T \rightarrow T$ the sequence $\text{Res}_{L/K} T \rightarrow T \rightarrow 1$ is Zariski exact as well. By Lemma 4 the upper horizontal map in the diagram

$$\begin{array}{ccc} (K(L; T, T)/n)_G & \xrightarrow{N_{L/K}} & K(K; T, T)/n \\ \downarrow & & \downarrow \\ H^2(L, T[n] \otimes T[n])_G & \xrightarrow{\text{cor}} & H^2(K, T[n] \otimes T[n]) \end{array}$$

is an isomorphism. The vertical maps are Galois symbols. Since T_L is a split torus the left vertical map is an isomorphism by the Merkurjev-Suslin theorem [MS]. Thus to finish the proof it remains to show that the lower vertical arrow is not injective. Note that $T[n] \cong \mathbb{Z}/n\mathbb{Z}(1) \otimes X$. Hence the assertion follows from Lemma 5 and Lemma 7 below.

Lemma 7 *There exists homomorphisms of G -modules $e : \mathbb{Z} \rightarrow X \otimes_{\mathbb{Z}} X$ and $f : X \otimes_{\mathbb{Z}} X \rightarrow \mathbb{Z}$ such that $f \circ e : \mathbb{Z} \rightarrow \mathbb{Z}$ is multiplication by $n - 1$.*

Proof. For a G -module M we write M^\vee for the G -module $\text{Hom}(M, \mathbb{Z})$. Let $(,) : \mathbb{Z}[G] \otimes_{\mathbb{Z}} \mathbb{Z}[G] \rightarrow \mathbb{Z}$ be the symmetric pairing given by

$$(18) \quad (g, g') = \begin{cases} 1 & \text{if } g = g', \\ 0 & \text{if } g \neq g'. \end{cases}$$

It yields an isomorphism $\mathbb{Z}[G] \rightarrow \mathbb{Z}[G]^\vee$. For a submodule $M \subseteq \mathbb{Z}[G]$ let

$$M^\perp = \{x \in \mathbb{Z}[G] \mid (x, m) = 0 \ \forall m \in M\}.$$

Then we have $X^\perp = \mathbb{Z}S$ and $(\mathbb{Z}S)^\perp = X$ where $S = \sum_{i=0}^{n-1} \sigma^i$. Thus (18) yields an isomorphism $X \cong (\mathbb{Z}[G]/\mathbb{Z}S)^\vee$. By (17) we have $\mathbb{Z}[G]/\mathbb{Z}S \cong X$, hence

$$X \otimes_{\mathbb{Z}} X \cong X \otimes_{\mathbb{Z}} X^\vee \cong \text{Hom}(X, X)$$

Thus it suffices to prove the assertion for $\text{Hom}(X, X)$. Obviously, for the two maps $e : \mathbb{Z} \rightarrow \text{Hom}(X, X), m \mapsto m \text{id}_X$ and $f : \text{Hom}(X, X) \rightarrow \mathbb{Z}, \tau \mapsto \text{Tr}(\tau)$ we have $f \circ e = \text{rank}(X) = n - 1$. \square

Remark 8 It is easy to construct examples where the assumptions (13) and (14) above are satisfied. For instance if K is a 2-local field satisfying property (13) and L/K is any cyclic extension of degree n then (14) holds by [Ka].

3 Counterexample to a conjecture of Beilinson

We first introduce some notation and recall a few facts from [Vo1] and [MVW]. Let K be a field of characteristic zero. Let Cor_K denote the additive category of finite correspondences ([MVW], 1.1). The objects of Cor_K

are smooth separated K -schemes of finite type and for $X, Y \in \text{Obj}(\text{Cor}_K)$ the group of morphisms $\text{Cor}_K(X, Y)$ is the free abelian group generated by integral closed subschemes W of $X \times Y$ which are finite and surjective over X . Let $D^-(\text{Shv}_{\text{Nis}}(\text{Cor}_K))$ (resp. $D^-(\text{Shv}_{\text{et}}(\text{Cor}_K))$) denote the derived category of complexes of Nisnevich (resp. étale) sheaves with transfer bounded from above.

The category of effective motivic complexes $\text{DM}_{\text{Nis}}^{\text{eff}, -}(K)$ (resp. étale effective motivic complexes $\text{DM}_{\text{et}}^{\text{eff}, -}(K)$) is the full subcategory of $D^-(\text{Shv}_{\text{Nis}}(\text{Cor}_K))$ (resp. $D^-(\text{Shv}_{\text{et}}(\text{Cor}_K))$) which consists of complexes C^* with homotopy invariant cohomology sheaves $H^i(C^*)$ for all i (see [Vo1], 3.1 or [MVW], 14.1, resp. 9.2). $\text{DM}_{\text{Nis}}^{\text{eff}, -}(K)$ and $\text{DM}_{\text{et}}^{\text{eff}, -}(K)$ are triangulated tensor categories. They are equipped with the t-structure induced from the standard t-structure on $D^-(\text{Shv}_{\text{Nis}}(\text{Cor}_K))$ (resp. $D^-(\text{Shv}_{\text{et}}(\text{Cor}_K))$). There is a covariant functor $M : \text{Cor}_K \rightarrow \text{DM}_{\text{Nis}}^{\text{eff}, -}(K)$, $X \mapsto M(X)$ and we have $M(X \times Y) = M(X) \otimes M(Y)$. There is also the "change of topology" functor $\alpha^* : \text{DM}_{\text{Nis}}^{\text{eff}, -}(K) \rightarrow \text{DM}_{\text{et}}^{\text{eff}, -}(K)$. It is a tensor functor which admits a right adjoint $R\alpha_* : \text{DM}_{\text{et}}^{\text{eff}, -}(K) \rightarrow \text{DM}_{\text{Nis}}^{\text{eff}, -}(K)$.

Beilinson [Be] has proposed the following generalization of the Milnor-Bloch-Kato conjecture: For any smooth affine K -scheme X the adjunction morphism $M(X) \rightarrow R\alpha_*\alpha^*M(X)$ induces an isomorphism on cohomology in degrees ≤ 0 , i.e. the map

$$(19) \quad a_X : M(X) \longrightarrow t_{\leq 0}R\alpha_*\alpha^*M(X)$$

is an isomorphism in $\text{DM}_{\text{Nis}}^{\text{eff}, -}(K)$.

If $X = (\mathbb{G}_m)^d = \mathbb{G}_m \times \dots \times \mathbb{G}_m$ (d -fold product of \mathbb{G}_m) we have $M(X) \cong (\mathbb{Z} \oplus \mathbb{Z}(1)[1])^d$. Thus a_X is an isomorphism if and only if

$$(20) \quad \mathbb{Z}(n) \longrightarrow t_{\leq n}R\alpha_*\alpha^*\mathbb{Z}(n)$$

is an isomorphism for all $n \leq d$. It is known (compare [SV]) that the Milnor-Bloch-Kato conjecture is equivalent to the assertion that (20) is an isomorphism for all $n \geq 0$.

Let L/K be a separable quadratic extension and let $T := \text{Ker}(N_{L/K} : \text{Res}_{L/K} \mathbb{G}_m \rightarrow \mathbb{G}_m)$. We shall show that (19) is in general not an isomorphism for $X = T^n$ for $n \geq 2$. By ([HK], 7.3) there exists a canonical decomposition $M(T) = \mathbb{Z} \oplus \mathbb{Z}(L/K, 1)[1]$ where $\mathbb{Z}(L/K, 1)$ is the cone of the morphism $\mathbb{Z}(1) \rightarrow \text{Res}_{L/K} \mathbb{Z}(1)$.

Remarks 9 (a) Here is a more explicit description of the motive $\mathbb{Z}(L/K, 1)$. The torus T defines a homotopy invariant étale (hence Nisnevich) sheaf with transfer and therefore an element of $\text{DM}_{\text{Nis}}^{\text{eff}, -}(K)$. We have

$$\mathbb{Z}(L/K, 1) \cong T[-1].$$

This can be deduced from the corresponding statement for \mathbb{G}_m ([MVW], 4.1) and the exactness of (15) (as a sequence in $Shv_{\text{Nis}}(\text{Cor}_K)$).

(b)¹ Let A_1, \dots, A_n be semi-abelian varieties over K . It should be possible to identify the generalized Milnor K -group $K(K; A_1, \dots, A_n)$ with a Hom-group in $\text{DM}_{\text{Nis}}^{\text{eff}, -}(K)$. For that we view A_1, \dots, A_n again as elements in $Shv_{\text{Nis}}(\text{Cor}_K)$. Then we expect that

$$K(K; A_1, \dots, A_n) \cong \text{Hom}_{\text{DM}_{\text{Nis}}^{\text{eff}, -}(K)}(\mathbb{Z}, A_1 \otimes \dots \otimes A_n).$$

If $A_1 = \dots = A_n = \mathbb{G}_m$ this is proved in ([MVW], lecture 5) and it is likely that the proof given there can be adapted to the case of arbitrary semi-abelian varieties.

For $p, q \geq 0$ and $n = p + q$ we define

$$\mathbb{Z}(L/K, p, q) := \mathbb{Z}(L/K, 1)^{\otimes p} \otimes \mathbb{Z}(q)$$

and denote by $C(p, q)$ the cone of $\mathbb{Z}(L/K, p, q) \rightarrow t_{\leq n} R\alpha_* \alpha^* \mathbb{Z}(L/K, p, q)$. Note that $\mathbb{Z}(L/K, p, q)[n]$ is a direct summand of $M(T^p \times (\mathbb{G}_m)^q)$. We also put $C(n) := C(0, n)$. We have

$$(21) \quad C(n) \cong (t_{\geq n+1} R\alpha_* \mathbb{Q}/\mathbb{Z}(n))[-1]$$

This follows from the Milnor-Bloch-Kato conjecture (in fact for our purpose we need (21) only after localization at the prime 2 where it follows from the Milnor conjecture [Vo2]).

Tensoring $\mathbb{Z}(1) \rightarrow \text{Res}_{L/K} \mathbb{Z}(1) \rightarrow \mathbb{Z}(L/K, 1) \rightarrow \mathbb{Z}(1)[1]$ with $\mathbb{Z}(L/K, p-1, q)$ (for $p \geq 1, q \geq 0$) yields a distinguished triangle

$$\mathbb{Z}(L/K, p-1, q+1) \rightarrow \text{Res}_{L/K} \mathbb{Z}(n) \rightarrow \mathbb{Z}(L/K, p, q) \rightarrow \mathbb{Z}(L/K, p-1, q+1)[1]$$

hence also a triangle

$$(22) \quad C(p-1, q+1) \rightarrow \text{Res}_{L/K} C(n) \rightarrow C(p, q) \rightarrow C(p-1, q+1)[1].$$

The following Lemma follows easily by induction on q using (21) and (22).

Lemma 10 *Let $p \geq 1, q \geq 0$ and $n = p + q$. Then we have $H^k(C(p, q)) = 0$ for $k < q + 2$ and*

$$H^{q+2}(C(p, q))(K) \cong H^{n+1}(L/K, \mathbb{Q}/\mathbb{Z}(n))$$

where $H^{n+1}(L/K, \mathbb{Q}/\mathbb{Z}(n)) := \text{Ker}(H^{n+1}(K, \mathbb{Q}/\mathbb{Z}(n)) \xrightarrow{\text{res}} H^{n+1}(L, \mathbb{Q}/\mathbb{Z}(n)))$.

¹This remark has been communicated to us by B. Kahn.

Since $[L : K] = 2$ we have

$$\begin{aligned} H^{n+1}(L/K, \mathbb{Q}/\mathbb{Z}(n)) &\cong H^{n+1}(L/K, \mathbb{Q}_2/\mathbb{Z}_2(n)) \cong H^{n+1}(L/K, \mathbb{Z}/2\mathbb{Z}(n)) \\ &\cong H^{n+1}(L/K, \mathbb{Z}/2\mathbb{Z}) \end{aligned}$$

(the second isomorphism is a consequence of the Milnor conjecture). Now the following Proposition follows by applying Lemma 10 for $(p, q) = (2, 0)$ and $(n, 0)$.

Proposition 11 (a) *There exists a short exact sequence*

$$0 \longrightarrow H^0(M(T \times T))(K) \longrightarrow R^0 \alpha_* \alpha^* M(T \times T)(K) \longrightarrow H^3(L/K, \mathbb{Z}/2\mathbb{Z}) \longrightarrow 0$$

In particular if $H^3(L/K, \mathbb{Z}/2\mathbb{Z}) \neq 0$ then (19) is not an isomorphism for $X = T \times T$.

(b) *More generally let n be an integer ≥ 2 and assume that $H^{n+1}(L/K, \mathbb{Z}/2\mathbb{Z}) \neq 0$. Then the map (19) is not an isomorphism for $X = T^n$. More precisely either the map*

$$H^{2-n}(M(X)) \longrightarrow R^{2-n} \alpha_* \alpha^* M(X)$$

is not surjective or

$$H^{3-n}(M(X)) \longrightarrow R^{3-n} \alpha_* \alpha^* M(X)$$

is not injective.

An n -local field K of characteristic 0 provides an example where the above assumption holds. In fact by [Ka] we have $H^{n+1}(L/K, \mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ for such fields.

References

- [Be] A.A. Beilinson, Talk at Fields Institute at Toronto on 21 March 2007.
- [HK] A. Huber, B. Kahn, *The slice filtration and mixed Tate motives*. Compos. Math. **142** (2006), 907–936.
- [Ka] K. Kato, *A generalization of local class field theory by using K -groups I*, J. Fac. Sci. Univ. Tokyo **26** (1979), 303–376.
- [MVW] C. Mazza, V. Voevodsky and C. Weibel, *Lecture notes on motivic cohomology*. Clay Mathematics Monographs **2**. American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2006.

- [MS] A.S. Merkurjev, A.A. Suslin, *K-cohomology of Severi-Brauer varieties and the norm residue homomorphism*. Math. USSR Izvestiya **21** (1983), 307–340.
- [Se] J.-P. Serre, *Algebraic groups and class fields*. Graduate Texts in Mathematics **117**. Springer Verlag 1988.
- [So] M. Somekawa, *On Milnor K-groups attached to semi-abelian varieties*. K-theory **4** (1990), 105–119.
- [SV] A.A. Suslin, V. Voevodsky, *Bloch-Kato conjecture and motivic cohomology with finite coefficients*. In: The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), 117–189, NATO Sci. Ser. C Math. Phys. Sci., 548, Kluwer Acad. Publ., Dordrecht, 2000.
- [Vo1] V. Voevodsky, *Triangulated categories of motives over a field*. In: V. Voevodsky, A. Suslin, E.M. Friedlander: Cycles, Transfers, and Motivic Homology Theories. Annals of Math. Studies **143**, Princeton University Press, 2000, 188–238.
- [Vo2] V. Voevodsky, *Motivic cohomology with $\mathbb{Z}/2$ -coefficients*. Publ. Math., Inst. Hautes Étud. Sci. **98**, 59–104 (2003).

Michael Spieß
 Fakultät für Mathematik
 Universität Bielefeld
 Postfach 100131
 D-33501 Bielefeld, Germany
 mspiess@math.uni-bielefeld.de

Takao Yamazaki
 Mathematical Institute
 Tohoku University
 Aoba
 Sendai 980-8578, Japan
 ytakao@math.tohoku.ac.jp