

Differential Geometry: Problem set 7

November 14, 2006

Due Wednesday, November 29

1. Consider the following vector field and 1-form on \mathbb{R}^3

$$\xi = (x^2 + x^3) \frac{\partial}{\partial x^1} + (x^1 + x^3) \frac{\partial}{\partial x^2} + (x^1 + x^2) \frac{\partial}{\partial x^3} \in \mathfrak{X}(\mathbb{R}^3)$$

$$\alpha = (x^1)^2 dx^1 + (x^2)^2 dx^2 + (x^3)^2 dx^3 \in \Omega^1(\mathbb{R}^3)$$

Compute

$$L_\xi \alpha$$

2. The 1-ball $\mathbb{B}_1^n \subset \mathbb{R}^n$ is a smooth manifold with boundary. Check that the top form on S^{n-1} given by problem 6 of homework 6 defines the boundary orientation of $S^{n-1} = \partial \mathbb{B}_1^n$.

Compute

$$\int_{S^{n-1}} \omega$$

[Hint: reduce it to an integral you know or can find somewhere....]

3. Let $f : M \rightarrow N$ be a smooth map. Check that

$$f^* : \Omega^* N \rightarrow \Omega^* M$$

descends to a homomorphism

$$f^* : H_{dR}^*(N) \rightarrow H_{dR}^*(M)$$

4. (a) Check that

$$\wedge : H_{dR}^p(M) \times H_{dR}^{n-p}(M) \rightarrow H_{dR}^n(M)$$

given by

$$[\alpha] \wedge [\beta] = [\alpha \wedge \beta]$$

is well-defined and bilinear.

- (b) If M is closed (i.e. compact without boundary) and orientable prove that

$$\dim(H_{dR}^n(M)) \geq 1$$

- (c) If M is closed oriented, then check that

$$\int \wedge : H_{dR}^p(M) \times H_{dR}^{n-p}(M) \rightarrow \mathbb{R}$$

given by

$$[\alpha], [\beta] \mapsto \int_M \alpha \wedge \beta$$

is a well-defined bilinear map.

5. Give S^n the induced Riemannian metric from the (standard) embedding into \mathbb{R}^{n+1} and compute in stereographic coordinates from the north pole $(0, 0, \dots, 0, 1)$ the metric g_{ij} .
6. One model of **hyperbolic n -space** is the Riemannian n -manifold

$$\mathbb{H}^n = \{(x^1, \dots, x^n) \mid x^n > 0\}$$

with the Riemannian metric

$$g_{ij} = \frac{\delta_{ij}}{(x^n)^2}$$

That is, the Euclidean inner product scaled by the reciprocal of the height squared.

- (a) Given $(t^1, \dots, t^{n-1}) \in \mathbb{R}^{n-1}$, check that the map

$$f(x^1, \dots, x^n) = (x^1 + t^1, \dots, x^{n-1} + t^{n-1}, x^n)$$

is an isometry.

- (b) Given $\lambda \in \mathbb{R}^+$ (the positive real numbers) check that

$$g(x^1, \dots, x^n) = (\lambda x^1, \dots, \lambda x^n)$$

is an isometry.