

MATH 242 QUIZ 4

NAME (Print your name): Type B

SECTION:

You need to show all your work to get a full credit.

Problem 1 Let $\vec{r}(t) = \cos 3t\vec{i} + \sin 3t\vec{j} + 4t\vec{k}$.

a)(8 points) Find the unit tangent and the unit normal vector.

solution) $\vec{r}'(t) = -3\sin 3t\vec{i} + 3\cos 3t\vec{j} + 4\vec{k}$

$$|\vec{r}'(t)| = \sqrt{16 + 9} = 5$$

$$\vec{T}(t) = \frac{\vec{r}'(t)}{|\vec{r}'(t)|} = \frac{1}{5}(-3\sin 3t\vec{i} + 3\cos 3t\vec{j} + 4\vec{k})$$

$$\vec{T}'(t) = \frac{1}{5}(-9\cos 3t\vec{i} - 9\sin 3t\vec{j})$$

$$|\vec{T}'(t)| = \frac{9}{5}$$

$$\text{So, } \vec{N}(t) = \frac{\vec{T}'(t)}{|\vec{T}'(t)|} = -\cos 3t\vec{i} - \sin 3t\vec{j}.$$

b)(4 points) Find the normal vector to the osculating plane at $t = \frac{\pi}{12}$.

solution) The normal vector to the osculating plane at $t = \frac{\pi}{12}$ is

$$\begin{aligned}\vec{B}\left(\frac{\pi}{12}\right) &= \vec{T}\left(\frac{\pi}{12}\right) \times \vec{N}\left(\frac{\pi}{12}\right) \\ &= \left\langle -\frac{3\sqrt{2}}{10}, \frac{3\sqrt{2}}{10}, \frac{4}{5} \right\rangle \times \left\langle -\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}, 0 \right\rangle \\ &= \frac{2\sqrt{2}}{5}\vec{i} - \frac{2\sqrt{2}}{5}\vec{j} + \frac{3}{5}\vec{k}.\end{aligned}$$

Problem 2(8 points) A particle has a velocity $\vec{v}(t) = \langle 2e^t, 2e^{-t}, t \rangle$ and its initial position is at $\vec{r}(0) = \langle 0, 1, 0 \rangle$. Find its position vector.

solution)

$$\vec{r}(t) = \int \vec{v}(t) dt = \langle 2e^t + C_1, -2e^{-t} + C_2, \frac{t^2}{2} + C_3 \rangle$$

$$\vec{r}(0) = \langle 2 + C_1, -2 + C_2, C_3 \rangle = \langle 0, 1, 0 \rangle$$

$$C_1 = -2, C_2 = 3, C_3 = 0$$

$$\text{So, } \vec{r}(t) = \langle 2e^t - 2, -2e^{-t} + 3, \frac{t^2}{2} \rangle.$$