

# A NOTE ON THE STRONG DIRECT SUMMAND CONJECTURE

JASON MCCULLOUGH

ABSTRACT. We prove special cases of the strong direct summand conjecture and the related strong monomial conjecture.

## 1. INTRODUCTION

In [HH95], Hochster and Huneke formulated the following conjecture.

**Conjecture 1.1** (Vanishing Maps of Tor Conjecture(VMTC)). *Let  $R \rightarrow A \rightarrow S$  be homomorphisms of Noetherian rings such that  $R$  and  $S$  are regular and such that  $A$  is module-finite over  $R$ . Let  $M$  be any  $R$ -module. Then the maps*

$$\mathrm{Tor}_i^R(M, A) \rightarrow \mathrm{Tor}_i^R(M, S)$$

*are zero for all  $i \geq 1$ .*

They show that the VMTC is a consequence of the existence of weakly functorial big Cohen-Macaulay algebras and, therefore, a theorem in the equicharacteristic case. The VMTC also holds in dimension at most 3 by Hochster's extension [Hoc02] of Heitmann's theorem in [Hei02]. They proved similar results previously in [HH90] and [HH93] using tight closure theory. Ranganathan proved some special cases in [Ran00]. The study of this conjecture is motivated by the fact that the VMTC implies that direct summands of regular rings are Cohen-Macaulay and that the VMTC implies Hochster's direct summand conjecture (see [HH90], [HH93], [HH95]).

**Conjecture 1.2** (Direct Summand Conjecture(DSC)). *Let  $R$  be a regular ring and let  $A$  be a module-finite extension of  $R$ . Then  $R$  is a direct summand of  $A$  as an  $R$ -module.*

The DSC first appeared in [Hoc73]. The DSC and its equivalent forms (including the monomial conjecture (MC), the canonical element conjecture (CEC) and the improved new intersection conjecture) have been studied by many people. We refer the reader to REFS. Like the other homological conjectures, the VMTC is still an open question in the general mixed characteristic case.

In [Ran00], Ranganathan proved that the VMTC is actually equivalent to the following statement.

**Conjecture 1.3** (Strong Direct Summand Conjecture(SDSC)). *Let  $(R, \mathfrak{m})$  be a regular local ring and let  $A$  be a module-finite extension of  $R$ . Then for every minimal generator  $x$  of  $\mathfrak{m}$  and for every height one prime ideal  $Q$  of  $A$  lying over  $xR$ , we have that  $xR$  is a direct summand of  $Q$  as an  $R$ -module.*

---

2000 *Mathematics Subject Classification*. Primary: 13D22; Secondary: 13D45, 13D07, 13H05.

*Key words and phrases*. Strong direct summand conjecture, strong monomial conjecture, vanishing maps of tor.

By the equivalence, it follows that the SDSC is a theorem in the equicharacteristic case and in dimension at most three. The equivalence is proved through a number of reductions and equivalences. See [Hoc07] for a nice summary. As the name suggests, the SDSC implies direct summand conjecture. Both conjectures reduce easily to the case where  $R$  and  $A$  are complete local domains. So it is natural to formulate a version of the monomial conjecture that implies the SDSC. Ranganathan formulated the following strong monomial conjecture in [Ran00]:

**Conjecture 1.4** (Strong Monomial Conjecture(SMC)). *Let  $A$  be a local domain with system of parameters  $x_1, \dots, x_d$ . Let  $Q$  be a height one prime of  $A$  containing  $x_1$ . Then for all  $t > 0$ ,*

$$x_1(x_1 \cdots x_d)^t \notin (x_1^{t+1}, \dots, x_d^{t+1})Q.$$

The SMC is one of the main focuses of this paper. Moreover, we extend the notion of the SMC to the following setting. Let  $A$  be a local domain with system of parameters  $\mathbf{x} = x_1, \dots, x_d$ . Let  $Q$  be a height one prime of  $A$  and let  $z \in Q - \mathbf{x}Q$ . Then we say the SMC holds for  $A$ ,  $Q$ ,  $\mathbf{x}$  and  $z$  if for all  $t > 0$

$$z(x_1 \cdots x_d)^t \notin (x_1^{t+1}, \dots, x_d^{t+1})Q.$$

Hence the usual SMC is the case where  $z = x_1$ . The generality introduced is necessary later in the statement of Theorem 2.5.

We extend results of Dutta [Dut98] and Strooker and Stuckrad [SS93] to prove the following

**Proposition 2.1.** *Let  $(A, \mathfrak{m})$  be a complete local domain and let  $\mathbf{x} = x_1, \dots, x_d$  a system of parameters for  $A$ . Let  $Q$  a height one prime of  $A$ . Let  $z \in Q - \mathbf{x}Q$  with  $z \notin \mathbf{x}Q$ . Let  $(S, \mathfrak{n}, k)$  be a local complete intersection ring with  $A = S/\mathfrak{p}$ ,  $\dim(S) = \dim(A)$  and  $\mathfrak{p}S_{\mathfrak{p}} = 0$ . Then for all  $t > 0$ ,*

$$z(x_1 \cdots x_d)^t \notin (x_1^{t+1}, \dots, x_d^{t+1})Q$$

*if and only if  $\text{Im} \left( \text{Hom}_S(Q, S) \xrightarrow{z} \text{Hom}_S(S, S) = S \right)$  is not contained in  $\mathbf{x}S$ . In particular, the strong monomial conjecture holds if and only if*

$$\text{Im} \left( \text{Hom}_S(Q, S) \xrightarrow{x_1} \text{Hom}_S(S, S) \right) \not\subset \mathbf{x}S.$$

Here  $\text{Hom}_S(Q, S) \xrightarrow{z} \text{Hom}_S(S, S)$  denotes the map induced by applying  $\text{Hom}_S(-, S)$  to the map  $S \rightarrow Q$  sending  $1 \mapsto z$ . Using the previous result, we obtain the following corollary.

**Corollary 2.2.** *The Strong Monomial Conjecture holds for all local rings if and only if it holds for all local almost complete intersection rings.*

Combining with a result from [Dut98], we also prove the following characterization of the SMC in terms of the length of certain Koszul homology modules. Here  $S$  denotes a complete intersection ring mapping onto  $A$  as in Proposition 2.1.

**Proposition 2.3.** *Set  $J = \text{Im}(\text{Hom}_S(Q, S) \xrightarrow{z} \text{Hom}_S(S, S)) \subset S$ . Then the Strong Monomial Conjecture holds for  $Q$ ,  $z$  and  $\mathbf{x}$  if and only if  $\ell(J/\mathbf{x}J) > \ell(H_1(\mathbf{x}; S/J))$ .*

Finally we show that under the mild hypothesis that  $A/\mathfrak{n}$  is infinite, where  $\mathfrak{n}$  is the maximal ideal of  $A$ , we may use Samuel's theory of superficial elements and the previous result to show the following:

**Theorem 2.5.** *Suppose the residue field of  $A$  is infinite. Let  $x_1, \dots, x_d$  be a s.o.p. Let  $Q$  be a height one prime in  $A$ . Let  $z \in Q$ . Finally let  $J = \text{Im}(\text{Hom}_S(Q, S) \xrightarrow{z} \text{Hom}_S(S, S))$ . Then there exist  $y_1, \dots, y_{d-1} \in A$  such that  $(x_1, \dots, x_d) = (y_1, \dots, y_{d-1}, x_d)$  and such that SMC holds for  $A, Q, y_1, \dots, y_{d-1}, x_d^t$  and  $x_1$  for  $t \gg 0$ .*

Note that  $x_1, \dots, x_d$  satisfies the SMC if and only if  $y_1, \dots, y_{d-1}, x_d$  as above do also. The above results are extensions of results of Dutta on the monomial conjecture in [Dut98] and use similar methods.

Finally we show that the SDSC is true if one assumes the direct summand conjecture in lower dimension that  $H_{\mathfrak{m}}^{d-1}(A) = 0$  and where  $d = \dim(A)$ . We note that this case includes rings  $A$  that are Cohen-Macaulay, local almost complete intersection domains, and complete local normal domains with  $S_3$  canonical module. (See [Dut97].)

## 2. THE STRONG MONOMIAL CONJECTURE

In [Dut98] and [SS93], the monomial conjecture is characterized by certain ideals of nonzero divisors not being contained in any parameter ideal in a Gorenstein or complete intersection ring surjecting onto  $A$ . We extend those results to the strong monomial conjecture below.

For this section, let us fix some notation. Let  $(R, \mathfrak{m})$  denote a regular local ring with  $\mathfrak{m}$  its unique maximal ideal. Let  $A$  denote a module-finite extension of  $R$ . Let  $d = \dim(A) = \dim(R)$ . For the purposes of the SDSC and SMC, we may assume that  $R$  and  $A$  are complete local domains. We focus on the SMC for most of this section.

We also make use of the following setup. Let  $R$  and  $A$  be as above. By mapping a Cohen ring onto  $A$  and modding out by an appropriate regular sequence, one can construct a complete intersection  $S$  surjecting onto  $A$  such that  $A = S/\mathfrak{p}$  for a prime  $\mathfrak{p} \subset S$  with  $\mathfrak{p}S_{\mathfrak{p}} = (0)$  and  $\mathfrak{p} \in \text{Ass}_S(S)$ . See Proposition 1 in [SS93]. For  $Q$  a height one ideal in  $A$  and  $z \in Q$ , we denote by  $\text{Hom}_S(Q, S) \xrightarrow{z} \text{Hom}_S(S, S)$  the map obtained by applying  $\text{Hom}_S(-, S)$  to the map  $S \rightarrow Q$  sending  $1 \mapsto z$ . Finally we abbreviate system of parameters by s.o.p.

**Proposition 2.1.** *Let  $A$  be a complete local domain and let  $\mathbf{x} = x_1, \dots, x_d$  a s.o.p. for  $A$ . Let  $Q$  a height one prime of  $A$ . Let  $z \in Q - \mathbf{x}Q$  with  $z \notin \mathbf{x}Q$ . Let  $(S, \mathfrak{n})$  be a local complete intersection ring with  $A = S/\mathfrak{p}$ ,  $\dim(S) = \dim(A)$  and  $\mathfrak{p}S_{\mathfrak{p}} = 0$ . Then for all  $t > 0$ ,*

$$z(x_1 \cdots x_d)^t \notin (x_1^{t+1}, \dots, x_d^{t+1})Q \quad (\star)$$

*if and only if  $\text{Im}(\text{Hom}_S(Q, S) \xrightarrow{z} \text{Hom}_S(S, S) = S)$  is not contained in  $\mathbf{x}S$ . In particular, the strong monomial conjecture holds if and only if*

$$\text{Im}(\text{Hom}_S(Q, S) \xrightarrow{x_1} \text{Hom}_S(S, S)) \not\subset \mathbf{x}S.$$

*Proof.* We have the sequence of maps

$$S \twoheadrightarrow A \xrightarrow{z} Q.$$

Set  $E = E_S(k) \simeq H_n^d(S)$ , where  $E_S(k)$  is the injective envelope of the residue field of  $S$ . Applying  $H_n^d(-)$  and  $- \otimes_S S/\mathbf{x}S$  we get the following diagram:

$$\begin{array}{ccc}
 & \text{Ann}_E(\mathbf{x}) & \\
 & \nearrow \simeq & \nwarrow \\
 S/\mathbf{x}S & \xrightarrow{\mu_S^{\mathbf{x}}} & H_n^d(S) = E \\
 \downarrow & \dashrightarrow \psi & \downarrow \\
 A/\mathbf{x}A & \xrightarrow{\mu_A^{\mathbf{x}}} & H_n^d(A) \\
 \downarrow z & \searrow \varphi & \downarrow z \\
 Q/\mathbf{x}Q & \xrightarrow{\mu_Q^{\mathbf{x}}} & H_n^d(Q)
 \end{array}$$

In this diagram,  $\mu_M^{\mathbf{x}} : M \rightarrow H_n^d(M)$  denotes the direct limit map

$$\mu_M^{\mathbf{x}} : M/\mathbf{x}M \rightarrow \varinjlim_t M/\mathbf{x}^t M \simeq H_n^d(M).$$

Note that  $\mu_S^{\mathbf{x}}$  is injective because  $S$  is Cohen-Macaulay and that  $\mu_S^{\mathbf{x}}$  factors through  $\text{Ann}_E(\mathbf{x})$ . Also note that the  $(\star)$  holds if and only if  $\varphi \neq 0$ . Further since  $S/\mathbf{x}S \rightarrow A/\mathbf{x}A$  is surjective,  $\varphi \neq 0$  if and only if  $\psi \neq 0$ .

Let  $(-)^{\vee} = \text{Hom}_S(-, E)$  denote the Matlis dual. Applying  $(-)^{\vee}$  to the above diagram yields the following commutative diagram:

$$\begin{array}{ccc}
 & S/\mathbf{x}S & \\
 & \swarrow \simeq & \nwarrow \\
 (S/\mathbf{x}S)^{\vee} & \longleftarrow & \text{Hom}_S(S, S) \simeq S \\
 \uparrow & \dashrightarrow \psi^{\vee} & \uparrow \\
 (A/\mathbf{x}A)^{\vee} & \longleftarrow & \text{Hom}_S(A, S) \simeq \Omega_A \\
 \uparrow & \searrow \varphi^{\vee} & \uparrow z \\
 (Q/\mathbf{x}Q)^{\vee} & \longleftarrow & \text{Hom}_S(Q, S)
 \end{array}$$

Since  $(-)^{\vee}$  is a faithful, exact functor, we have that  $\psi \neq 0$  if and only if  $\psi^{\vee} \neq 0$ . Since  $\text{Hom}_S(Q, S)$  injects into  $S$ , we see that this is equivalent to  $J := \text{Im}(\text{Hom}_S(Q, S)) \not\subseteq \mathbf{x}S$ .  $\square$

As noted in the introduction, the freedom in separating  $z$  from  $x_1$  in the previous proposition will be used to prove an analog of the SMC later in Theorem 2.5.

We follow the development by Dutta [Dut98] extending results about the monomial conjecture to the SMC. First, we obtain the following reduction of the SMC using the previous result.

**Corollary 2.2.** *The Strong Monomial Conjecture holds for all local rings if and only if it holds for all local almost complete intersection rings.*

*Proof.* Let  $A$  be a local ring and  $S$  a complete intersection ring of equal dimension with  $A = S/\mathfrak{p}$  and  $\mathfrak{p}S_{\mathfrak{p}} = 0$ . Let  $Q$  be a height one prime of  $A$ . Pick  $\lambda \in \mathfrak{p} -$

$\bigcup\{\mathfrak{q} \in \text{Ass}_S(S), \mathfrak{q} \neq \mathfrak{p}\}$ . Write  $Q = \tilde{Q}/\mathfrak{p}$  for a height one prime  $\tilde{Q}$  of  $S$ . Now take  $\mathfrak{q} \in \text{Ass}_S(S)$ . If  $\mathfrak{q} = \mathfrak{p}$ , then since  $\mathfrak{p}S_{\mathfrak{p}} = 0$ , we have  $\mathfrak{q} \notin \text{Supp}_S(\mathfrak{p}/\lambda S)$ . If  $\mathfrak{q} \neq \mathfrak{p}$  then  $\mathfrak{q} \notin \text{Supp}_S(\mathfrak{p}/\lambda S)$  by the choice of  $\lambda$ . Hence  $\text{Ass}_S(\text{Hom}_S(\mathfrak{p}/\lambda S, S)) = \text{Supp}_S(\mathfrak{p}/\lambda S) \cap \text{Ass}_S(S) = \emptyset$  and so  $\text{Hom}_S(\mathfrak{p}/\lambda S, S) = 0$ . Therefore the injection  $\varphi$  from the exact sequence

$$0 \rightarrow \text{Hom}_S(Q, S) \xrightarrow{\varphi} \text{Hom}_S(\tilde{Q}/\lambda S, S) \rightarrow \text{Hom}_S(\mathfrak{p}/\lambda S, S)$$

is actually an isomorphism. Hence the image of  $\text{Hom}_S(Q, S)$  in  $S$  is the same as the image of  $\text{Hom}_S(\tilde{Q}/\lambda S, S)$ . The claim follows from the previous proposition.  $\square$

In the following result,  $H_i(\mathbf{x}; M)$  denotes the  $i$ th Koszul homology of the  $A$ -module  $M$  with respect to a list of element  $\mathbf{x}$ .

**Proposition 2.3.** *Let  $A$  be a complete local domain and let  $S$  be a complete intersection surjecting onto  $A$  as in Proposition 2.1. Let  $Q$  be a height one prime of  $A$ . Set  $J = \text{Im}(\text{Hom}_S(Q, S) \xrightarrow{z} \text{Hom}_S(S, S)) \subset S$ . Then the Strong Monomial Conjecture holds for  $Q$ ,  $z$  and  $\mathbf{x}$  if and only if  $\ell(J/\mathbf{x}J) > \ell(H_1(\mathbf{x}; S/J))$ .*

*Proof.* Consider the short exact sequence

$$0 \rightarrow J \rightarrow S \rightarrow S/J \rightarrow 0.$$

Applying  $-\otimes_S S/\mathbf{x}S$  we get the exact sequence

$$0 \rightarrow H_1(\mathbf{x}; S/J) \rightarrow J/\mathbf{x}J \rightarrow S/\mathbf{x}S \rightarrow S/(J + \mathbf{x}S) \rightarrow 0.$$

By the previous lemma, the SMC holds if and only if  $J \not\subset \mathbf{x}S$ , which happens if and only if  $\ell(S/\mathbf{x}S) > \ell(S/(J + \mathbf{x}S))$ . (Here  $\ell(M)$  denotes the length of  $M$ .) Since length is additive, we have  $\ell(J/\mathbf{x}J) > \ell(H_1(\mathbf{x}; S/J))$ , and conversely.  $\square$

We will make use of the following result of Dutta, which appears as Proposition 1.4 in [Dut98]. Here  $e(x_1, \dots, x_n; A)$  denotes the Hilbert multiplicity of  $A$  with respect to a s.o.p.  $x_1, \dots, x_n$ .

**Proposition 2.4** (Dutta). *Let  $x_1, \dots, x_n$  be a s.o.p. of a local ring  $A$  such that*

$$e(x_1, \dots, x_n; A) = e(x_2, \dots, x_n; A/x_1A) = \dots = e(x_n; A/(x_1, \dots, x_{n-1}A)).$$

*Then  $H_i(x_1, \dots, x_n; A)$  is a module of finite length for every  $i > 0$ .*

By extending by a faithfully flat extension, we may assume that the residue field of  $A$  is infinite. Recall that this does not change the issue of splitting. So we may apply Samuel's theory of superficial elements. We refer the reader to [ZS93] for background material on superficial elements.

**Theorem 2.5.** *Suppose the residue field of  $A$  is infinite. Let  $x_1, \dots, x_d$  be a s.o.p. Let  $Q$  be a height one prime in  $A$ . Let  $z \in Q$ . Then there exist  $y_1, \dots, y_{d-1} \in A$  such that  $(x_1, \dots, x_{d-1}) = (y_1, \dots, y_{d-1})$  and such that the SMC holds for  $A$ ,  $Q$ , and  $y_1, \dots, y_{d-1}, x_d^t$  and  $x_1$  for  $t \gg 0$ .*

*Proof.* Lift the system of parameters  $x_1, \dots, x_d$  to a system of parameters in  $S$ . (By abuse of notation, write them the same.) Thus by Remark to Theorem VIII-22 of [ZS93], we may choose  $y_1, \dots, y_{d-1}$  in such a way so that each  $y_i$  is of the form

$$y_i = a_{1j}x_1 + \dots + \lambda_i x_i + \dots + a_{dj}x_d$$

with  $\lambda_i$  a unit and  $a_{ij} \in A$  for all  $i$  and  $j$ . (Note that  $(x_1, \dots, x_d) = (y_1, \dots, y_{d-1}, x_d)$  in  $S$  and hence also in  $A$ .) Further, we may choose the  $y_i$  to be superficial

elements with respect to  $\mathbf{x}$ . The same is true of their images in  $S/J$ , where  $J = J = \text{Im}(\text{Hom}_S(Q, S) \xrightarrow{z} \text{Hom}_S(S, S))$  as above. We have by Proposition 2.4 that  $\ell(H_1(y_1, \dots, y_{d-1}; S/J)) < \infty$ .

Set  $\mathbf{y} = y_1, \dots, y_{d-1}$ . Now consider the exact sequence

$$0 \rightarrow \frac{H_1(\mathbf{y}; S/J)}{x_d^t H_1(\mathbf{y}; S/J)} \rightarrow H_1(y_1, \dots, y_{d-1}, x_d^t; S/J) \rightarrow (0 : x_d^t)A/\mathbf{y} \rightarrow 0.$$

Since  $\ell(H_1(\mathbf{y}; S/J)) < \infty$ , we have  $x_n^t H_1(\mathbf{y}; A) = 0$  for  $t \gg 0$ . Also  $\ell((0 : x_n^t)A/\mathbf{y})$  is constant for  $t \gg 0$ . Therefore  $\ell(H_1(y_1, \dots, y_{d-1}, x_d^t; S/J))$  is constant for  $t \gg 0$ .

Finally observe that since  $\text{depth}(S) = d$ , we have  $\dim(J) = d$ . Thus

$$\lim_{t \rightarrow \infty} J/(y_1, \dots, y_{d-1}, x_d^t)J = \infty.$$

It follows then that for  $t$  sufficiently large,

$$\ell(J/(y_1, \dots, y_{d-1}, x_d^t)J) > \ell(H_1(y_1, \dots, y_{d-1}, x_d^t; S/J)).$$

By the previous Proposition, the SMC holds for  $z$ ,  $Q$  and the system of parameters  $y_1, \dots, y_{d-1}, x_d^t$ .  $\square$

Now we return to the strong direct summand conjecture. In Theorem 2.1 of [Dut01], Dutta proved that the direct summand conjecture holds when  $H_{\mathfrak{m}}^{d-1}(A) = 0$ . (To be precise, it is shown that the stronger canonical element conjecture holds for  $A$ .) We use this result to show that the SDSC also holds in this case if one assumes the general direct summand conjecture in lower dimension.

**Theorem 2.6.** *Let  $R$  and  $A$  be complete local rings such that  $R$  is regular and such that  $A$  is module-finite over  $R$ . Suppose  $H_{\mathfrak{m}}^{d-1}(A) = 0$  and that the direct summand conjecture holds in dimension  $d - 1$ . Then the strong direct summand conjecture holds for  $A$ .*

*Proof.* Let  $Q$  be a height one prime of  $A$  lying over  $xR$  where  $x$  is a regular parameter of  $R$ . Note that by [Dut97], since  $H_{\mathfrak{m}}^{d-1}(A) = 0$ , the direct summand conjecture holds for  $A$ . This is equivalent to saying that the map  $\text{Hom}_R(A, R) \rightarrow \text{Hom}_R(R, R)$  is onto. By local duality, this is equivalent to the map  $H_{\mathfrak{m}}^d(R) \rightarrow H_{\mathfrak{m}}^d(A)$  being injective. Similarly, we have that  $H_{\mathfrak{m}}^{d-1}(R/xR) \rightarrow H_{\mathfrak{m}}^{d-1}(A/Q)$  and we must prove that  $H_{\mathfrak{m}}^d(xR) \rightarrow H_{\mathfrak{m}}^d(Q)$  is injective.

We have the following commutative diagram with exact rows.

$$\begin{array}{ccccccccc} 0 = H_{\mathfrak{m}}^{d-1}(A) & \longrightarrow & H_{\mathfrak{m}}^{d-1}(A/Q) & \longrightarrow & H_{\mathfrak{m}}^d(Q) & \longrightarrow & H_{\mathfrak{m}}^d(A) & \longrightarrow & 0 \\ & & \uparrow & & \uparrow & & \uparrow & & \\ & & 0 & \longrightarrow & H_{\mathfrak{m}}^{d-1}(R/xR) & \longrightarrow & H_{\mathfrak{m}}^d(xR) & \longrightarrow & H_{\mathfrak{m}}^d(R) & \longrightarrow & 0 \end{array}$$

As in the previous proof, the maps  $H_{\mathfrak{m}}^{d-1}(R/xR) \rightarrow H_{\mathfrak{m}}^{d-1}(A/Q)$  and  $H_{\mathfrak{m}}^{d-1}(R) \rightarrow H_{\mathfrak{m}}^d(A)$  are injective. Since  $H_{\mathfrak{m}}^{d-1}(A) = 0$ , it follows that  $H_{\mathfrak{m}}^d(xR) \rightarrow H_{\mathfrak{m}}^d(Q)$  is injective as well. Therefore  $xR \hookrightarrow Q$  splits.  $\square$

Finally we recall two results of Dutta that illuminate special cases of the previous theorem. The first appears as Lemma 2.1 in [Dut97]. The second is a rewording of Theorem 2.6 in [Dut97].

**Theorem 2.7** (Dutta). *Let  $A$  be a complete local almost complete intersection domain. Then  $H_{\mathfrak{m}}^{d-1}(A) = 0$ .*

**Theorem 2.8** (Dutta). *Let  $A$  be a complete local normal domain such that its canonical module  $\Omega_A$  is  $S_3$ . Then  $H_{\mathfrak{m}}^{d-1}(A) = 0$ .*

**Corollary 2.9.** *Let  $A$  be a complete local domain of dimension  $d$ . Suppose the DSC holds for rings of dimension  $d - 1$ . Then the SDSC holds in the following cases:*

- (1)  *$A$  is Cohen-Macaulay.*
- (2)  *$A$  is an almost complete intersection domain.*
- (3)  *$A$  is normal and  $\Omega_A$  is  $S_3$ , where  $\Omega_A$  denotes the canonical module of  $A$ .*

*Proof.* In all three cases,  $H_{\mathfrak{m}}^{d-1}(A) = 0$ . The result follows from Theorem 2.6.  $\square$

### 3. ACKNOWLEDGEMENTS

The author was supported in part by a University of Illinois Research Board grant and NSA grant number H98230-06-1-0026, both awarded to Prof. Sankar Dutta. The author would like to thank Sankar Dutta for his direction and support in numerous discussions.

### REFERENCES

- [Dut97] S. P. Dutta, *Dualizing complex and the canonical element conjecture II*, The Journal of the London Mathematical Society, Second Series **56** (1997), no. 1, 49–63.
- [Dut98] ———, *A note on the monomial conjecture*, Trans. Amer. Math. Soc. **350** (1998), no. 7, 2871–2878.
- [Dut01] ———, *Splitting of local cohomology of syzygies of the residue field*, Journal of Algebra **244** (2001), 168–185.
- [Hei02] Raymond C. Heitmann, *The direct summand conjecture in dimension three*, Ann. of Math. **156** (2002), no. 2, 695–712.
- [HH90] Melvin Hochster and Craig Huneke, *Tight closure, invariant theory, and the Brionçon-Skoda theorem*, Journal of the American Mathematical Society **3** (1990), no. 1, 31–116.
- [HH93] ———, *Phantom homology*, vol. 103, Memoirs of the AMS, no. 490, American Mathematical Society, 1993.
- [HH95] ———, *Applications of the existence of Big Cohen-Macaulay algebras*, Advances in Mathematics **113** (1995), 45–117.
- [Hoc73] Melvin Hochster, *Contracted ideals from integral extensions of regular rings*, Nagoya Mathematical Journal **51** (1973), 25–43.
- [Hoc02] ———, *Big Cohen-Macaulay algebras in dimension three via Heithmann’s theorem*, Journal of Algebra **254** (2002), 395–408.
- [Hoc07] ———, *Homological conjectures, old and new*, Illinois Journal of Math **51** (2007), no. 1, 151 – 169.
- [Ran00] Nandini Ranganathan, *Splitting in module-finite extension rings and the vanishing conjecture for maps of tor*, Ph.D. thesis, University of Michigan, 2000.
- [SS93] Jan R. Strooker and Jurgen Stückrad, *Monomial conjecture and complete intersections*, Manuscripta Math. **79** (1993), no. 2, 153–159.
- [ZS93] Oscar Zariski and Pierre Samuel, *Commutative algebra*, vol. II, Graduate Texts in Mathematics, no. 29, Springer-Verlag, 1993.

DEPARTMENT OF MATHEMATICS, UNIVERISTY OF ILLINOIS AT URBANA-CHAMPAIGN, 273 ALT-GELD HALL, MC-382, 1209 W. GREEN ST., URBANA, IL 61801-2975

*E-mail address:* jmcullo@math.uiuc.edu