

RAMSEY'S THEOREM AND CONE AVOIDANCE

DAMIR D. DZHAFAROV AND CARL G. JOCKUSCH, JR.

ABSTRACT. It was shown by Cholak, Jockusch, and Slaman that every computable 2-coloring of pairs admits an infinite low_2 homogeneous set H . We answer a question of the same authors by showing that H may be chosen to satisfy in addition $C \not\leq_T H$, where C is a given noncomputable set. This is shown by analyzing a new and simplified proof of Seetapun's cone avoidance theorem for Ramsey's theorem. We then extend the result to show that every computable 2-coloring of pairs admits a pair of low_2 infinite homogeneous sets whose degrees form a minimal pair.

1. INTRODUCTION

In this paper we show that the infinite homogeneous sets asserted to exist by Ramsey's theorem for pairs can be chosen to have low information content in a sense to be made precise below.

Definition 1.1. Fix an infinite set $S \subseteq \omega$ and $n, k \in \omega$.

- (i) We denote by $[S]^n$ the collection of all sets $T \subseteq S$ of cardinality n .
- (ii) A k -coloring f of $[S]^n$ is a map $[S]^n \rightarrow \{0, 1, \dots, k-1\}$. If $S = \omega$ and $n = 2$, we call f a k -coloring of *pairs*.
- (iii) We call a k -coloring f of $[S]^2$ *stable* if for each $s \in S$ there exist $i < k$ and $t_0 \in \omega$ such that $f(\{s, t\}) = i$ for all $t \in S$ with $t > t_0$.
- (iv) A set $H \subseteq S$ is said to be *homogeneous* for a k -coloring f of $[S]^n$ if f is constant on $[H]^n$.

Ramsey's theorem for k -colorings of exponent n (denoted RT_k^n) asserts that for every k -coloring of $[\omega]^n$ there exists an infinite homogeneous set. The effective content and logical strength of RT_k^n has been studied in [6], [9], and [1], among other papers.

Definition 1.2. Given a sequence C_0, C_1, \dots of sets, we call a set S *cone avoiding* (for this sequence) provided $C_i \not\leq_T S$ for all $i \in \omega$.

We begin by recalling the following two results, both answering questions of Jockusch [6], pages 274 and 277.

Theorem 1.3 (Seetapun [9], Theorem 2.1). *Given a sequence C_0, C_1, \dots of non-computable sets, every computable 2-coloring of pairs admits an infinite cone avoiding homogeneous set.*

We are grateful to Denis Hirschfeldt for many insightful discussions throughout the preparation of this article, and in particular for first bringing to our attention the possibility of using our proof of Theorem 3.4 to prove Theorem 3.5. We thank Antonín Kučera and Robert Soare for helpful comments on some of the material in Section 2. We also thank Joseph Mileti for a careful reading of an early draft of this article. The second author was partially supported by NSFC Grand International Joint Project 'New Directions in Theory and Applications of Models of Computation', No. 60310213.

This theorem was used in [9], Theorem 3.1, to show that RT_2^2 is strictly weaker than ACA_0 over RCA_0 . For information on the systems ACA_0 and RCA_0 , and Reverse Mathematics in general, see Simpson [11].

Theorem 1.4 (Cholak, Jockusch, and Slaman [1], Theorem 3.1). *Every computable 2-coloring of $[\omega]^2$ admits an infinite homogeneous set H which is low_2 , i.e. $H'' \leq_T 0''$.*

Cholak, Jockusch, and Slaman gave two different proofs of the above theorem. The first proof (in Section 4 of [1]) featured control of the first jump of H . It established the stronger result ([1], Corollary 12.6) that every computable 2-coloring of pairs admits an infinite homogeneous set whose jump has degree at most \mathbf{d} , where \mathbf{d} is a given degree $\gg 0'$. (See Definition 2.6 and the subsequent discussion. Also see the remarks after the proof of Theorem 3.4 for an explanation of a missing point in the proof of this stronger result.) The other proof of Theorem 1.4 (in Section 5 of [1]) featured direct control of the second jump of H . It was adapted in [1], Theorem 10.2, to models of second order arithmetic to show that every Π_1^1 sentence provable from $\text{RCA}_0 + \Sigma_2^0\text{-IND} + \text{RT}_2^2$ is provable from just $\text{RCA}_0 + \Sigma_2^0\text{-IND}$, where $\Sigma_2^0\text{-IND}$ is the induction scheme for Σ_2^0 formulas. Seetapun and Slaman [9], page 580, asked whether Theorem 1.3, for the special case where $C_i \equiv_T 0'$ for all $i \in \omega$, could be effectivized to produce a non-high cone avoiding infinite homogeneous set. A positive answer (for arbitrary noncomputable C_i) was given by Cholak, Jockusch, and Slaman [1], Theorem 12.2. In view of Theorem 1.4, they in turn raised the following question:

Question 1.5 (Cholak, Jockusch, and Slaman [1], Question 13.11). *Given a non-computable set C , does every computable 2-coloring of pairs admit an infinite low_2 homogeneous set H with $C \not\leq_T H$?*

In this paper, we present a new proof of Seetapun's theorem (Theorem 1.3) which is considerably more straightforward than both the original (cf. [9], Section 2) as well as the simplification of Hummel and Jockusch (cf. [3], Theorem 2.1). This proof is based in part on the method used by Cholak, Jockusch, and Slaman in [1], Section 4, to prove Theorem 1.4 and benefited from discussions with Steffen Lempp and a subsequent suggestion of Slaman for simplifying it. The proof proceeds by first reducing the result to the stable case, and this reduction is straightforward. The crucial new element in the proof is Lemma 4.7, which handles cone avoidance in the stable case. By keeping track of the effective content in this new proof, we are able to give an affirmative answer to Question 1.5. We extend this result to show that every computable 2-coloring of pairs admits a pair of low_2 infinite homogeneous sets whose degrees form a minimal pair. The proof of this uses an extension of the technique of [1], Section 5, together with a new basis theorem for Π_1^0 classes. Along the way, we extend Theorem 2.5 of Jockusch and Stephan [5] to show that there is a minimal pair of low_2 cohesive degrees.

2. BASIS THEOREMS

The low basis theorem of Jockusch and Soare [7], Theorem 2.1, played a crucial role in the proof of Theorem 1.4. Hence, it is not surprising that a cone avoiding version of the Low Basis Theorem will play a central role in answering Question 1.5. The needed result is due to Linda Lawton (cf. [2], Theorem 4.1).

Theorem 2.1 (Lawton). *Let $T \subseteq 2^{<\omega}$ be an infinite computable tree, let C_0, C_1, \dots be a sequence of noncomputable sets, and let $C = \bigoplus_{i \in \omega} C_i$. Then T has an infinite cone avoiding path f such that $f' \leq_T 0' \oplus C$. Furthermore, an index for f' as a $(0' \oplus C)$ -computable set can be found $(0' \oplus C)$ -effectively from an index for T .*

The theorem is proved by forcing with Π_1^0 classes, alternating between forcing the jump and applying the following lemma.

Lemma 2.2 (Lawton). *Let $T \subseteq 2^{<\omega}$ be an infinite computable tree, let C be any noncomputable set, and let $e \in \omega$. Then T has an infinite computable subtree \tilde{T} such that $\Phi_e^f \neq C$ for all $f \in [\tilde{T}]$, and an index for \tilde{T} can be found $(0' \oplus C)$ -effectively from e and an index for T .*

Proof. For each $n \in \omega$, define the computable tree

$$U_n = \{\sigma \in T : \neg(\Phi_{e,|\sigma|}^\sigma(n) \downarrow = C(n))\},$$

and notice that U_n must be infinite for some n . Indeed, if each U_n were finite, then to compute $C(n)$ for a given n we would need only to search for $m, k \in \omega$ such that $\Phi_{e,|\sigma|}^\sigma(n) = k$ for all $\sigma \in T$ of length m , from which we conclude that $C(n) = k$. This contradicts the assumption that C is noncomputable. Clearly, an index for U_n as a computable tree can be found C -effectively from e and an index for T , and $0'$ can determine whether or not U_n is infinite. (For the latter, use that $U_n \subseteq 2^{<\omega}$.) Thus, $0' \oplus C$ can find the least n such that U_n is infinite, and we set $\tilde{T} = U_n$. \square

Note, in particular, that if the sequence C_0, C_1, \dots is taken uniformly Δ_2^0 , Theorem 2.1 asserts the existence, in any non-empty Π_1^0 class of sets, of a low member which avoids the upper cone of each C_i . The following variation on this result, in the case where the Π_1^0 class in question contains no computable members, is included here for general interest. As a special case, we get that any non-empty Π_1^0 class of sets with no computable members contains a low set Turing incomparable with a given noncomputable low set.

Theorem 2.3. *Let $T \subseteq 2^{<\omega}$ be an infinite computable tree with no computable paths, let C_0, C_1, \dots be a sequence of noncomputable sets, and let $D = \bigoplus_{i \in \omega} C_i$. Then T contains an infinite path f such that $f \upharpoonright_T C_i$ for all $i \in \omega$, and $f' \leq_T D$. Further, an index for f' as a D -computable function can be found D -effectively from an index for T .*

We can obtain the theorem from the next lemma, by interspersing the construction used in its proof into that of Theorem 2.1.

Lemma 2.4. *Let $T \subseteq 2^{<\omega}$ be an infinite computable tree with no computable paths, let C be any set, and let $e \in \omega$. Then T has an infinite computable subtree \tilde{T} such that $\Phi_e^C \neq f$ for all $f \in [\tilde{T}]$, and an index for \tilde{T} can be found C' -effectively from e and an index for T .*

Proof. Since $T \subseteq 2^{<\omega}$ is infinite and computable and has no computable paths, T has at least two distinct paths, and thus there exist incompatible strings σ_1 and σ_2 such that each σ_i is extendible to a path through T . Such σ_1 and σ_2 can be found $0'$ -effectively since the set of extendible nodes of T is $0'$ -computable. Fix n such that $\sigma_1(n) \downarrow \neq \sigma_2(n) \downarrow$. Use C' to determine whether $\Phi_e^C(n) \downarrow$. If not, let $\tilde{T} = T$. If so, choose $j \in \{1, 2\}$ such that $\Phi_e^C(n) \downarrow \neq \sigma_j(n) \downarrow$, and let \tilde{T} be the set of all strings $\sigma \in T$ which are compatible with σ_j . \square

Finally, we present the following “minimal pair basis theorem” which will be central to our work in Section 6. Recall that two degrees \mathbf{a} and \mathbf{b} are said to form a *minimal pair* if $\mathbf{a} \cap \mathbf{b} = \mathbf{0}$ (we do not require that \mathbf{a} and \mathbf{b} be non-zero).

Theorem 2.5. *Let $T \subseteq 2^{<\omega}$ be an infinite computable tree, and let B be any set. Then T has an infinite path f with $f'' \leq_T 0'' \oplus B'$ whose degree makes a minimal pair with that of B . Furthermore, an index for f'' as a $(0'' \oplus B')$ -computable set can be found $(0'' \oplus B')$ -effectively from an index for T .*

Proof. We force with Π_1^0 classes. Set $T_0 = T$ and suppose that for some $e \geq 0$, $T_e \subseteq 2^{<\omega}$ is an infinite computable tree of known index. For each $n \in \omega$, define

$$U_n = \{\sigma \in T_e : \Phi_{e,|\sigma|}^\sigma(n) \uparrow\},$$

the index of which as a computable tree can be found effectively from an index for T_e . First, ask $0''$ whether U_n is infinite for some n , and if so, search $0'$ -effectively for the least such n and set $T_{e+1} = U_n$. In this case, Φ_e^f is not total for any $f \in [T_{e+1}]$. Otherwise, we know that Φ_e^f is total for all $f \in [T_e]$. In this case, we ask $0''$ if there exist extendible nodes $\sigma, \tau \in T_e$ and an $n \in \omega$ such that $\Phi_{e,|\sigma|}^\sigma(n) \downarrow \neq \Phi_{e,|\tau|}^\tau(n) \downarrow$. If there do, find the least such σ, τ , and n in some fixed effective listing and inquire of B' whether $\Phi_e^B(n)$ converges. If it does, let T_{e+1} consist of all nodes in T_e comparable with ρ , where ρ is the lexicographically smaller of σ and τ such that $\Phi_e^\rho(n) \neq \Phi_e^B(n)$. Clearly $\Phi_e^f \neq \Phi_e^B$ for all $f \in [T_{e+1}]$ in this case. If $\Phi_e^B(n)$ diverges, or if there do not exist any such σ, τ , and n , set $T_{e+1} = T_e$. Suppose now that Φ_e^B and Φ_e^f are equal and total, where $f \in [T_{e+1}]$. Then each U_n is finite and σ, τ and n as above did not exist. Given n , $\Phi_e^B(n)$ can now be computed as follows. First, search for the least k such that $\Phi_{e,|\sigma|}^\sigma(n) \downarrow$ for all $\sigma \in T$ of length k , and then search for the least $\ell \geq k$ and $m \in \omega$ such that $\Phi_{e,|\sigma|}^\sigma(n) = m$ for all $\sigma \in T$ of length k having an extension on T of length ℓ . The second search succeeds because for any extendible σ on T of length k it must be that $\Phi_{e,|\sigma|}^\sigma(n) = \Phi_e^B(n)$, so m must be equal to $\Phi_e^B(n)$.

As usual, we iterate the construction to build a sequence $2^{<\omega} = T_0 \supseteq T_1 \supseteq \dots$ of infinite computable trees, and we take $f \in \bigcap_{e \in \omega} T_e$. Since the construction was $(0'' \oplus B')$ -effective, and since it decided for each e the totality of Φ_e^f , it follows that $f'' \leq_T 0'' \oplus B'$. \square

We will apply this result when B satisfies $B' \leq_T 0''$, in which case of course the path f of the previous theorem is low₂.

We conclude this section with the following definition due to Simpson [10], page 648:

Definition 2.6. Let \mathbf{a} and \mathbf{b} be Turing degrees. Then we write $\mathbf{a} \gg \mathbf{b}$ to mean that every infinite \mathbf{b} -computable tree $T \subseteq 2^{<\omega}$ has an \mathbf{a} -computable path.

It is known that $\mathbf{a} \gg \mathbf{b}$ if and only if every \mathbf{b} -computable $\{0, 1\}$ -valued partial function has an \mathbf{a} -computable total extension. From this it follows that there is a nonempty Π_1^0 class \mathcal{P} such that every element of \mathcal{P} has degree $\gg \mathbf{0}$. Namely, consider the class \mathcal{P} of all $\{0, 1\}$ -valued (total) functions g such that $g(\langle e, i \rangle) = \Phi_e(i)$ whenever $\Phi_e(i) \downarrow \leq 1$. The low basis theorem applied to \mathcal{P} consequently implies that there exists a low degree $\mathbf{a} \gg \mathbf{0}$. Theorem 2.1, on the other hand, gives for each sequence C_0, C_1, \dots of noncomputable sets the existence of a cone avoiding degree $\mathbf{a} \gg \mathbf{0}$ whose jump is below $\mathbf{0}' \cup \text{deg}(\bigoplus_{i \in \omega} C_i)$. And similarly Theorem 2.5

implies that for each degree \mathbf{b} there is a degree $\mathbf{a} \gg \mathbf{0}$ with $\mathbf{a}'' \leq \mathbf{0}'' \cup \mathbf{b}'$ and (\mathbf{a}, \mathbf{b}) a minimal pair.

3. CONE AVOIDANCE AND FIRST JUMP CONTROL

Recall that, given a sequence $\vec{R} = \{R_0, R_1, \dots\}$ of sets, an infinite set S is called \vec{R} -cohesive if for each $i \in \omega$ either $S \subseteq^* R_i$ or $S \subseteq^* \overline{R_i}$. If \vec{R} contains exactly the computable sets then an \vec{R} -cohesive set S is called r -cohesive, while if \vec{R} contains exactly the computably enumerable sets then S is called *cohesive*. It was shown by Jockusch and Stephan in [5], Corollary 2.4, that the r -cohesive and cohesive degrees coincide. (A degree is *cohesive* if it contains a cohesive set and r -cohesive if it contains an r -cohesive set.)

Theorem 1.4 was proved by using a low₂ r -cohesive set to reduce it to the stable case, and then proving the stable case by transforming it to a statement about Δ_2^0 sets. We proceed analogously here, but adding cone avoidance. We state two key lemmas, delaying their proofs until Section 5 below, and derive from them our main results. For the remainder of this section, let a degree $\mathbf{d} \gg \mathbf{0}'$ be fixed. Additionally, except where otherwise stated, let C_0, C_1, \dots be a fixed sequence of noncomputable sets, and let $C = \oplus_{i \in \omega} C_i$. ‘‘Cone avoiding’’ below refers to this sequence.

Lemma 3.1. *There exists a cone avoiding r -cohesive set X , and if $C \leq_T \mathbf{0}'$ then X can be chosen with jump of degree at most \mathbf{d} .*

Lemma 3.2. (i) *For $k \geq 2$, if A_1, \dots, A_k is a sequence of disjoint sets whose union is ω , then for some $i \leq k$, A_i contains an infinite cone avoiding subset.*

(ii) *If $A_1, \dots, A_k, C \leq_T \mathbf{0}'$, then for some $i \leq k$, A_i contains an infinite cone avoiding subset with jump of degree at most \mathbf{d} .*

We will, in addition, need the following well-known result:

Lemma 3.3 (Cholak, Jockusch, and Slaman [1], page 9). (i) *The restriction of any computable k -coloring of pairs to an r -cohesive set is stable.*

(ii) *If f is a computable stable k -coloring of pairs, then there exist pairwise disjoint Δ_2^0 sets A_1, \dots, A_k covering ω , any infinite subset of any of which computes an infinite homogeneous set for f .*

For clarity, we now restate Seetapun’s theorem, and show how it follows from Lemmas 3.1 and 3.2. The effectiveness of Lemmas 3.1 and 3.2 allows us to add an effectiveness result here for the case where the sequence C_0, C_1, \dots is uniformly Δ_2^0 . The latter is already a partial answer to Question 1.5.

Theorem 3.4. (i) (Seetapun [9], Theorem 2.1) *Every computable 2-coloring of pairs admits an infinite cone avoiding homogeneous set H .*

(ii) *If $C \leq_T \mathbf{0}'$, then H can be chosen so that $\deg(H)' \leq \mathbf{d}$.*

Proof. (i) Let f be a computable 2-coloring of pairs. By Lemma 3.1, fix a cone avoiding r -cohesive set X , and let $p : \omega \rightarrow X$ be its principal function. Define an X -computable 2-coloring g of pairs by $g(\{s, t\}) = f(\{p(s), p(t)\})$, for all $\{s, t\} \in [\omega]^2$, which must be stable by Lemma 3.3(i). By part (ii) of the same lemma, relativized to X , there exists a $\Delta_2^{0, X}$ set A , such that for any infinite subset S of A or \overline{A} , $X \oplus S$ computes an infinite homogeneous set for f . On the other hand, by Lemma 3.2(i), relativized to X , there is an infinite subset S of either A

of \bar{A} whose join with X is cone avoiding. Hence, g admits an infinite homogeneous set $\tilde{H} \leq_T X \oplus S$, and so by definition of g , $H = p(\tilde{H})$ is homogeneous for f . Now we have

$$H \leq_T X \oplus \tilde{H} \leq_T X \oplus S,$$

implying that H is cone avoiding.

(ii) Assume $C \leq_T 0'$. By Theorem 6.5 of Simpson [10], there exists a degree \mathbf{e} satisfying $\mathbf{d} \gg \mathbf{e} \gg 0'$, so by Lemma 3.1 we may choose X to satisfy $\deg(X)' \leq \mathbf{e}$. In particular, $\mathbf{d} \gg \deg(X)'$, and so we may use part (ii) of Lemma 3.2 above, relativized to X , in place of part (i) to find S with $X \oplus S$ not only cone avoiding but also with jump of degree at most \mathbf{d} . Defining H as before, we thus have $\deg(H)' \leq \deg(X \oplus S)' \leq \mathbf{d}$, as desired. \square

We remark that using density of the degrees under \gg in the proof of part (ii) above is also necessary to obtain Theorem 12.6 of [1] mentioned in the introduction, and is missing from the exposition in [1]. This was pointed out by Joseph Mileti.

An affirmative answer to Question 1.5 now easily follows.

Theorem 3.5. *Let C be noncomputable. Every computable 2-coloring f of pairs admits an infinite low₂ homogeneous set H such that $C \not\leq_T H$.*

Proof. We consider several cases:

Case 1: $C \not\leq_T 0''$. By Theorem 1.4, f admits an infinite low₂ homogeneous set H , meaning C cannot be computable in H'' , and hence also not in H .

Case 2: $C \leq_T 0'$. Fix \mathbf{a} satisfying $\mathbf{a} \gg 0'$ and $\mathbf{a}' \leq 0''$, as in the discussion at the end of Section 2. Now the desired result follows by Theorem 3.4(ii) using \mathbf{a} in place of \mathbf{d} .

Case 3: $C \not\leq_T 0'$ and $C \leq_T 0''$. By Lemma 2.1, relativized to $0'$, there exists a cone avoiding degree $\mathbf{a} \gg 0'$ with $\mathbf{a}' \leq 0'' \cup \deg(C) = 0''$. By Theorem 1.4 (or rather, by the extension of this theorem mentioned following its statement above), f admits an infinite homogeneous set H such that $\deg(H)' \leq \mathbf{a}$, so H is low₂. And since $\deg(C) \not\leq \mathbf{a}$, it follows that C is not computable in H . \square

We do not know whether Theorem 3.5 can be strengthened to assert that for every $C >_T 0$ every computable 2-coloring of pairs admits an infinite homogeneous set $H \not\leq_T C$ with $\deg(H)' \leq \mathbf{d}$. That is, we do not know whether the condition in Theorem 3.4(ii) that C be Δ_2^0 can be omitted in the situation where ‘‘cone avoiding’’ means simply $H \not\leq_T C$.

4. STRATEGIES FOR MEETING REQUIREMENTS

Lemmas 3.1 and 3.2 will be proved using Mathias forcing with suitable restrictions on the infinite part of the conditions. In this section we show that each requirement for these lemmas is dense for the relevant version of Mathias forcing, and we calculate the oracle needed to find the extensions needed to witness this density. We then complete the proofs of these lemmas in the next section. Throughout this section, we assume fixed a degree $\mathbf{d} \gg 0'$. Lemma 4.4 below is a variation on the proof in Section 4.1 of [1] of Theorem 2.5 of Jockusch and Stephan [5]. Lemmas 4.6, 4.7, and 4.8 are all variations of Lemma 4.6 of Cholak, Jockusch, and Slaman [1]. The last of these was suggested to us by Denis Hirschfeldt.

We begin by recalling and fixing some terminology.

Definition 4.1. (i) A *(Mathias) precondition* is a pair of sets (D, L) with D finite and $\max(D) < \min(L)$. A *(Mathias) condition* is a precondition (D, L) such that L is infinite.

(ii) A set G *satisfies* a precondition (D, L) if $D \subseteq G \subseteq D \cup L$.

(iii) A precondition (\tilde{D}, \tilde{L}) *extends* a precondition (D, L) , written $(\tilde{D}, \tilde{L}) \prec (D, L)$ if $\tilde{L} \subseteq L$ and \tilde{D} satisfies (D, L) . We call such an extension a *finite extension* if $L - \tilde{L}$ is finite.

(iv) We say a precondition (D, L) is *cone avoiding*, *computable*, or *low* if L is, respectively, cone avoiding, computable, or low. If (D, L) is computable or low, an *index* for this precondition is a pair of numbers (d, ℓ) such that d is the canonical index of D and ℓ is, respectively, a Δ_1^0 index for L or a $\Delta_1^{0,K}$ index for L' .

Definition 4.2. Fix $e \in \omega$ and a precondition (D, L) .

(i) We say (D, L) *forces agreement on e* if there is no $x \in \omega$, and there are no finite sets F_0, F_1 satisfying (D, L) such that $\Phi_e^{F_0}(x) \downarrow \neq \Phi_e^{F_1}(x) \downarrow$.

(ii) We say (D, L) *forces e out of the jump* if there is no finite set F satisfying (D, L) such that $e \in F'$.

4.1. Strategies for Proving Lemma 3.1.

Lemma 4.3 (Cholak, Jockusch, and Slaman [1], Lemma 4.4). *Let R be any computable set, (D, L) a computable condition, and $e \in \omega$.*

(i) *There is a computable extension of (D, L) which is satisfied only by sets G with $|G| \geq e$, and an index for it can be found effectively from e and an index for (D, L) .*

(ii) *There is a computable extension of (D, L) which is satisfied only by sets G with $G \subseteq^* R$ or $G \subseteq^* \bar{R}$, and an index for it can be found \mathbf{d} -effectively from e and an index for (D, L) .*

(iii) *There is a computable extension of (D, L) such that $G'(e)$ is the same for all sets G satisfying it, and an index for it can be found $0'$ -effectively from e and an index for (D, L) .*

Lemma 4.4. *Let C be any set, (D, L) a condition, and $e \in \omega$. Then there is a finite extension of (D, L) such that, for all sets G satisfying it, either $\Phi_e^G \neq C$, or Φ_e^G is not total, or else Φ_e^G is L -computable. Furthermore, there is a fixed uniform procedure for finding canonical indices of finite sets H_0, H_1 from e , the canonical index of D , and an oracle for $L' \oplus C$ such that the desired finite extension of (D, L) has the form $(D \cup H_0, L \setminus H_1)$.*

Proof. We consider two cases, the distinction between which is uniformly L' -effective in e and the canonical index of D .

Case 1: (D, L) *does not force agreement on e .* Then we search, L' -effectively from e and the canonical index of D , for $x \in \omega$, and for finite sets F_0, F_1 satisfying (D, L) such that $\Phi_e^{F_0}(x) \downarrow \neq \Phi_e^{F_1}(x) \downarrow$. One of the two computations, say $\Phi_e^{F_j}(x)$, must differ from $C(x)$, and we can, uniformly C' -effectively from i, x, F_0 , and F_1 , find such a j . We then let $\tilde{D} = F_j$, and let \tilde{L} consist of the elements of L greater than $\max(F_j)$ and use $(\Phi_e^{F_j}(x))$. Any set G satisfying (\tilde{D}, \tilde{L}) will agree with F_j below use $(\Phi_e^{F_j}(x))$ and hence will satisfy $\Phi_e^G(x) \downarrow = \Phi_e^{F_j}(x) \downarrow \neq C(x)$.

Case 2: (D, L) *forces agreement on e .* We then set $(\tilde{D}, \tilde{L}) = (D, L)$, and claim that if G satisfies (\tilde{D}, \tilde{L}) then either Φ_e^G is not total or else it is L -computable. Indeed,

if Φ_e^G is total, then to compute its value on input x we have only to L -effectively search through the canonical indices of finite sets satisfying (D, L) until we find one, call it F , such that $\Phi_e^F(x) \downarrow$. Such an F clearly exists since, for example, any sufficiently long initial segment of G will do, and we have $\Phi_e^F(x) = \Phi_e^G(x)$ for any such F , else the previous case would have applied.

In either case, (\tilde{D}, \tilde{L}) is a finite extension of (D, L) with the desired properties, and the desired uniformity follows easily from the construction. \square

4.2. Strategies for Proving Lemma 3.2. Let C_0, C_1, \dots be a sequence of non-computable sets, and let $C = \bigoplus_{i \in \omega} C_i$.

Lemma 4.5 (Cholak, Jockusch, and Slaman [1], Lemma 4.6). *Let A be any set, (D, L) a condition with $L \cap A$ and $L \cap \bar{A}$ infinite, and $e, i \in \omega$. There is a finite extension (\tilde{D}, \tilde{L}) of (D, L) such that, for all sets G satisfying this extension, $|G \cap A| \geq e$ and $|G \cap \bar{A}| \geq i$ (so in particular, (\tilde{D}, \tilde{L}) is low or cone avoiding if (D, L) is). Moreover, an index for this extension can be found $(0' \oplus A)$ -effectively from e, i , and an index for (D, L) .*

The following lemma adds cone avoidance to Lemma 4.6 of [1].

Lemma 4.6. *Let A be any set, (D, L) a cone avoiding condition, and $e, i \in \omega$. There is a cone avoiding extension (\tilde{D}, \tilde{L}) of (D, L) such that either $(G \cap A)'(e)$ is the same for all sets G satisfying this extension, or $(G \cap \bar{A})'(i)$ is the same for all such G . If, additionally, \mathbf{d} is a degree with $\mathbf{d} \gg 0'$, $A, C \leq_T 0'$, and (D, L) is low, then (\tilde{D}, \tilde{L}) can be chosen low, and its index can be found \mathbf{d} -effectively from e, i , and an index for (D, L) . Furthermore, \mathbf{d} can compute whether (\tilde{D}, \tilde{L}) forces the value of $(G \cap A)'(e)$ or $(G \cap \bar{A})'(i)$ and what value is forced.*

Proof. Define the $\Pi_1^{0,L}$ class \mathcal{P} to consist of all (possibly finite) sets $S \subseteq L$ such that $(D \cap A, S)$ forces e out of the jump and $(D \cap \bar{A}, L - S)$ forces i out of the jump. We consider two cases.

Case 1: $\mathcal{P} \neq \emptyset$. By Theorem 2.1, relativized to L , there exists an $S \in \mathcal{P}$ such that $L \oplus S$ is cone avoiding, and hence the same is true of S and $L - S$. Fix such an S . At least one of the following subcases must apply since L is infinite.

Subcase 1a: S is infinite. We then set $(\tilde{D}, \tilde{L}) = (D, S)$, and observe that if G is any set satisfying (\tilde{D}, \tilde{L}) , then $G \cap A$ satisfies $(D \cap A, S)$. Hence, $e \notin (G \cap A)'$, else any initial segment F of G sufficiently long that $D \cap A \subseteq F \cap A$ and $\Phi_e^{F \cap A}(e) \downarrow$ would contradict that $(D \cap A, S)$ forces e out of the jump.

Subcase 1b: $L - S$ is infinite. We set $(\tilde{D}, \tilde{L}) = (D, L - S)$, and it follows by a similar argument that $i \notin (G \cap \bar{A})'$ for all G satisfying (\tilde{D}, \tilde{L}) .

Case 2: $\mathcal{P} = \emptyset$. Then in particular $L \cap A \notin \mathcal{P}$, so at least one of the following subcases must hold:

Subcase 2a: *Some finite set F with $\Phi_e^F(e) \downarrow$ satisfies $(D \cap A, L \cap A)$.* We let $\tilde{D} = D \cup F$ and let \tilde{L} consist of those elements of L greater than $\max(\tilde{D})$ and use $(\Phi_e^F(e))$. Since $\tilde{L} =^* L$, it follows that \tilde{L} is cone avoiding. Furthermore, for any set G satisfying (\tilde{D}, \tilde{L}) , $G \cap A$ and F agree below $\text{use}(\Phi_e^F(e))$. It follows that $\Phi_e^{G \cap A}(e) \downarrow = \Phi_e^F(e)$, and hence $e \in (G \cap A)'$.

Subcase 2b: *Some finite set F with $\Phi_i^F(i) \downarrow$ satisfies $(D \cap \bar{A}, L - A)$. We let $\tilde{D} = D \cup F$ and let \tilde{L} consist of those elements of L greater than $\max(\tilde{D})$ and use $(\Phi_i^F(i))$. It follows analogously that $i \in (G \cap \bar{A})'$.*

In either case, (\tilde{D}, \tilde{L}) extends (D, L) and clearly has the desired properties.

Now assume that A and C are computable in $0'$, and that (D, L) is low. Notice that an index for \mathcal{P} as a $\Pi_1^{0,L}$ class can be found L' -effectively, hence $0'$ -effectively, from e, i , and canonical indices for $D \cap A$ and $D \cap \bar{A}$, hence, since A is Δ_2^0 , $0'$ -effectively from an index for (D, L) . By the effectiveness of Theorem 2.1 relativized to L , we can, in Case 1 above, $(0' \oplus C)$ -effectively, hence $0'$ -effectively, from an index for \mathcal{P} , find S and a $\Delta_1^{0,K}$ index for $(L \oplus S)'$ so as to additionally satisfy

$$(L \oplus S)' \leq_T L' \oplus C \leq_T 0' \oplus C \equiv_T 0'.$$

It follows that S and $L - S$ are both low, and hence that the extension (\tilde{D}, \tilde{L}) is, in every case considered above, not only cone avoiding but also low.

The distinction between Cases 1 and 2 is L' -effective from an index for \mathcal{P} , and hence $0'$ -effective.

We now argue, as in [1], Lemma 4.6, that one can \mathbf{d} -effectively choose a true subcase (1a or 1b) of Case 1. Consider the two statements “ S is infinite” and “ $L - S$ is infinite”. These are $\Pi_2^{0,L \oplus S}$ statements, hence Π_2^0 statements since $L \oplus S$ is low, and indices for them as such can be found effectively from a $\Delta_1^{0,K}$ index for $(L \oplus S)'$, an index for (D, L) , and an index for \mathcal{P} . Since at least one of the statements is true, it follows by choice of $\mathbf{d} \gg \mathbf{0}'$ that we can \mathbf{d} -effectively choose one which is true. (This is true because there is a partial function ψ computable in $0'$ which given indices a, b of Σ_2^0 statements at least one of which is true outputs an index (a or b) of one which is true, and \mathbf{d} can compute a total function which extends ψ .)

Since A is Δ_2^0 , the distinction between Subcases 2a and 2b amounts to $0'$ -effectively searching through finite sets satisfying $(D \cap A, L \cap A)$ and $(D \cap \bar{A}, L - A)$ until we find one, call it F , such that either $\Phi_e^F(e) \downarrow$ or $\Phi_i^F(i) \downarrow$.

We conclude that, in any case, an index for (\tilde{D}, \tilde{L}) can be found \mathbf{d} -effectively from e, i , and an index for (D, L) , as desired. Furthermore, we can \mathbf{d} -effectively find a statement in the list $e \in (G \cap A)', e \notin (G \cap A)', i \in (G \cap \bar{A})', i \notin (G \cap \bar{A})'$ which is true of all G satisfying (\tilde{D}, \tilde{L}) . \square

Lemma 4.7. *Let A be any set, (D, L) a cone avoiding condition, and $e, i, s, t \in \omega$. There is a cone avoiding extension (\tilde{D}, \tilde{L}) of (D, L) such that, for all sets G satisfying this extension, either $\Phi_e^{G \cap A} \neq C_s$ or $\Phi_i^{G \cap \bar{A}} \neq C_t$. If, additionally, $A, C \leq_T 0'$ and (D, L) is low, then (\tilde{D}, \tilde{L}) can be chosen low, and its index can be found \mathbf{d} -effectively from e, i , and an index for (D, L) .*

Proof. Define \mathcal{P} to be the class of all $S \subseteq L$ such that $(D \cap A, S)$ forces agreement on e and $(D \cap \bar{A}, L - S)$ forces agreement on i . Then \mathcal{P} is a $\Pi_1^{0,L}$ class.

Case 1: $\mathcal{P} \neq \emptyset$. By Theorem 2.1, relativized to L , there exists $S \in \mathcal{P}$ such that $L \oplus S$ is cone avoiding. In particular, S and $L - S$ are cone avoiding as well.

Subcase 1a: *S is infinite.* We then set $(\tilde{D}, \tilde{L}) = (D, S)$, and claim that if G is any set satisfying (\tilde{D}, \tilde{L}) then $\Phi_e^{G \cap A} \neq C_s$. Indeed, if $\Phi_e^{G \cap A}$ is total then we can, S -effectively from e and the canonical index of $D \cap A$, compute $\Phi_e^{G \cap A}(x)$ for any x , since we have only to search through finite sets F with $D \cap A \subseteq F \subseteq S$ until we

find one such that $\Phi_e^F(x) \downarrow$. Since a set satisfies (\tilde{D}, \tilde{L}) only if its intersection with A satisfies $(D \cap A, S)$, and since the latter forces agreement on e , it follows that we will then have $\Phi_e^F(x) = \Phi_e^{G \cap A}(x)$. Thus, our claim follows by virtue of S being cone avoiding.

Subcase 1b: $L - S$ is infinite. We set $(\tilde{D}, \tilde{L}) = (D, L - S)$ and can argue as above to conclude that $\Phi_i^{G \cap \bar{A}} \neq C_t$ for any set G satisfying (\tilde{D}, \tilde{L}) .

Case 2: $\mathcal{P} = \emptyset$. Thus $L \cap A \notin \mathcal{P}$, so we have at least one of the following two possibilities:

Subcase 2a: $(D \cap A, L \cap A)$ does not force agreement on e . Then for some $x \in \omega$, and for some finite set F satisfying $(D \cap A, L \cap A)$, $\Phi_e^F(x) \neq C_s(x)$. We set $\tilde{D} = D \cup F$, and let \tilde{L} consist of all elements of L greater than $\max(\tilde{D})$ and use $(\Phi_e^F(x))$. Then $\tilde{L} =^* L$, hence \tilde{L} is cone avoiding, and clearly $\Phi_e^{G \cap A} \neq C_s$ for any set G satisfying (\tilde{D}, \tilde{L}) .

Subcase 2b: $(D \cap \bar{A}, L - A)$ does not force agreement on i . We can then analogously define (\tilde{D}, \tilde{L}) so that $\Phi_e^{G \cap \bar{A}} \neq C_t$ for any set G satisfying (\tilde{D}, \tilde{L}) .

In either case, (\tilde{D}, \tilde{L}) extends (D, L) and has the desired properties.

Now if A and C are computable in $0'$ and (D, L) is low, we can argue as in the proof of the previous lemma. Briefly, an index for \mathcal{P} can, since A is Δ_2^0 , be found $0'$ -effectively from e, i, s, t and an index for (D, L) , and from this index, S in Case 1 can be found $0'$ -effectively so that S and $L - S$ are both low. The distinction between Cases 1 and 2 is L' -effective, hence $0'$ -effective, from an index for \mathcal{P} , while a true subcase of Case 1 (Subcase 1a or 1b) can be chosen \mathbf{d} -effectively. To distinguish between Subcases 2a and 2b, we can, since $A, C_s, C_t \leq_T 0'$, search $0'$ -effectively through ω and all finite sets until we find some $x \in \omega$ and either a finite set F satisfying $(D \cap A, L \cap A)$ with $\Phi_e^F(x) \downarrow \neq C_s(x)$, or a finite set F satisfying $(D \cap \bar{A}, L - A)$ with $\Phi_i^F(x) \downarrow \neq C_t(x)$. \square

Lemma 4.8. *Let A be any set, (D, L) a cone avoiding condition, and $e, i, s, t \in \omega$.*

(i) *There is a cone avoiding extension of (D, L) such that either $(G \cap A)'(e)$ is the same for all sets G satisfying this extension, or $\Phi_i^{G \cap \bar{A}} \neq C_t$ for all such G .*

(ii) *There is a cone avoiding extension of (D, L) such that either $(G \cap \bar{A})'(i)$ is the same for all sets G satisfying this extension, or $\Phi_e^{G \cap A} \neq C_s$ for all such G .*

If, additionally, $A, C \leq_T 0'$ and (D, L) is low, then (\tilde{D}, \tilde{L}) can be chosen low, and its index can be found \mathbf{d} -effectively from e, i , and an index for (D, L) .

Proof. We prove (i), the proof of (ii) being similar. Let \mathcal{P} be the $\Pi_1^{0,L}$ class of all sets $S \subseteq L$ such that $(D \cap A, S)$ forces e out of the jump and $(D \cap \bar{A}, L - S)$ forces agreement on i .

Case 1: $\mathcal{P} \neq \emptyset$. By Theorem 2.1, relativized to L , there exists $S \in \mathcal{P}$ such that S and $L - S$ are cone avoiding.

Subcase 1a: S is infinite. Set $(\tilde{D}, \tilde{L}) = (D, S)$. If G satisfies (\tilde{D}, \tilde{L}) , then $e \notin (G \cap A)'$ as in Subcase 1a of the proof of Lemma 4.6.

Subcase 1b: $L - S$ is infinite. We set $(\tilde{D}, \tilde{L}) = (D, L - S)$, concluding that $\Phi_i^{G \cap \bar{A}} \neq C_t$ for all G satisfying (\tilde{D}, \tilde{L}) , as in Subcase 1b of the proof of Lemma 4.7.

Case 2: $\mathcal{P} = \emptyset$. Then $L \cap A \notin \mathcal{P}$, so at least one of the following must obtain:

Subcase 2a: *some finite set F with $\Phi_e^F(e) \downarrow$ satisfies $(D \cap A, L \cap A)$.* We set $\tilde{D} = D \cup F$, and let \tilde{L} consist of all elements of L greater than $\max(\tilde{D})$ and use $(\Phi_e^F(e))$. As in Subcase 2a of the proof of Lemma 4.6, it follows that $e \in (G \cap A)'$ for every set G satisfying (\tilde{D}, \tilde{L}) .

Subcase 2b: *$(D \cap \bar{A}, L - A)$ does not force agreement on i .* Then for some $x \in \omega$, and some finite set F satisfying $(D \cap \bar{A}, L - A)$, $\Phi_i^F(x) \neq C_t(x)$ for any set G satisfying (\tilde{D}, \tilde{L}) .

Now suppose A and C are computable in $0'$ and that (D, L) is low. An index for \mathcal{P} can, since A is Δ_2^0 , be found $0'$ -effectively from e, i, t and an index for (D, L) . In Case 1 then, this index can be used to find S , $0'$ -effectively, so that S and $L - S$ are both low, in addition to being cone avoiding. The distinction between Cases 1 and 2 is L' -effective, hence $0'$ -effective, from an index for \mathcal{P} , and one may \mathbf{d} -effectively choose a true subcase of Case 1. To distinguish Subcases 2a and 2b, we can, since $A, C_t \leq_T 0'$, search $0'$ -effectively through ω and all finite sets, until we either find a finite set F satisfying $(D \cap A, L \cap A)$ with $\Phi_e^F(e) \downarrow$, or else some $x \in \omega$ and some finite set F satisfying $(D \cap \bar{A}, L - A)$ with $\Phi_i^F(x) \downarrow \neq C_t(x)$. \square

5. COMBINING STRATEGIES

In this section, we combine the forcing strategies from the previous section to prove Lemmas 3.1 and 3.2. Again, let C_0, C_1, \dots be a fixed sequence of noncomputable sets with $C = \bigoplus_{i \in \omega} C_i$.

Lemma 5.1. *Fix a degree $\mathbf{d} \gg 0'$. For every uniformly computable sequence R_0, R_1, \dots of sets, there exists a cone avoiding set G which is cohesive for this sequence and whose jump is of degree at most $\mathbf{d} \cup \deg(C)$.*

Proof. We force with computable conditions to build a set G which, for all $e, i \in \omega$, meets the requirements

$$\begin{aligned} \text{Req}_{4e} & : |G| \geq e \\ \text{Req}_{4e+1} & : G \subseteq^* R_e \text{ or } G \subseteq^* \overline{R_e} \\ \text{Req}_{4e+2} & : G'(e) \text{ is decided in the construction} \\ \text{Req}_{4\langle e, s \rangle + 3} & : \Phi_e^G \neq C_s. \end{aligned}$$

Let a computable condition (D, L) and $n \in \omega$ be given. If $n = 4e$ we apply Lemma 4.3(i), if $n = 4e + 1$ we apply (ii) with $R = R_e$, and if $n = 4e + 2$ we apply (iii). If $n = 4\langle e, s \rangle + 3$ we apply Lemma 4.4 with $C = C_s$. By the effectiveness of Lemmas 4.3 and 4.4, we in any case obtain a computable condition (\tilde{D}, \tilde{L}) such that any set satisfying this extension meets Req_n , and an index for this extension can be found $(\mathbf{d} \cup \deg(C))$ -effectively from an index for (D, L) .

To complete the proof, let $(D_0, L_0) = (\emptyset, \omega)$, and iterate the preceding density construction to $(\mathbf{d} \cup \deg(C))$ -effectively build a chain

$$(D_0, L_0) \succ (D_1, L_1) \succ \dots$$

such that, for any $n \in \omega$, any set satisfying (D_{n+1}, L_{n+1}) meets Req_n . Clearly, $G = \bigcup_i D_i$ has the desired properties. \square

Proof of Lemma 3.1. In view of the closing remark in Section 2, we may fix a cone avoiding degree $\mathbf{a} \gg \mathbf{0}$. The $\{0, 1\}$ -valued partial computable function

$$\tilde{p}(x) = \begin{cases} \Phi_e(x) & \text{if } \Phi_e(x) \downarrow \leq 1 \\ \uparrow & \text{otherwise} \end{cases}$$

consequently has a total, $\{0, 1\}$ -valued \mathbf{a} -computable extension p . Hence, the sequence

$$\vec{R} = \{R_0, R_1, \dots\},$$

where for $i \in \omega$,

$$R_i = \{x \in \omega : p(e, x) = 1\},$$

is a uniformly \mathbf{a} -computable sequence containing all the computable sets. Since $\deg(C_i) \not\leq \mathbf{a}$ for all $i \in \omega$, we may relativize Lemma 5.1 to \mathbf{a} to consequently obtain a cone avoiding \mathbf{r} -cohesive set X , as desired.

Now suppose $C \leq_T 0'$. We can choose \mathbf{a} so that $\mathbf{a}' \leq \mathbf{0}' \cup \deg(C)$, meaning in fact that \mathbf{a} can be chosen low, and hence such that $\mathbf{d} \gg \mathbf{a}' = \mathbf{0}'$. It follows, by the effectiveness of Lemma 5.1, that X can be found so that $\deg(X)' \leq \mathbf{d} \cup \deg(C) = \mathbf{d}$. \square

We next present a slightly weaker form of Lemma 3.2, from which the latter will follow by an easy induction.

Lemma 5.2. *Fix a degree $\mathbf{d} \gg \mathbf{0}'$.*

- (i) *Any set A either contains or is disjoint from an infinite cone avoiding set.*
- (ii) *If $A, C \leq_T 0'$, then A contains or is disjoint from an infinite cone avoiding set whose jump has degree at most \mathbf{d} .*

Proof. (i) If there exists a cone avoiding set S such that either $S \cap A$ is finite or $S \cap \bar{A}$ is finite, then we are done. So assume not. We force with cone avoiding conditions (D, L) to construct a set G which meets, for all $e, i, s, t \in \omega$ the requirements

$$\begin{aligned} \text{Req}_{2e} & : |G \cap A| \geq e \text{ and } |G \cap \bar{A}| \geq e \\ \text{Req}_{2\langle e, i, s, t \rangle + 1} & : \Phi_e^{G \cap A} \neq C_s \text{ or } \Phi_i^{G \cap \bar{A}} \neq C_t. \end{aligned}$$

We claim that these requirements are dense, in the sense that every cone avoiding condition can be extended to another, with the property that any set satisfying the extension meets a prescribed requirement. So let (D, L) be a cone avoiding condition, and fix $n \in \omega$. If $n = 2e$, then by our opening assumption we may apply Lemma 4.5. If $n = 2\langle e, i, s, t \rangle + 1$ we apply Lemma 4.7. In any case, we obtain a cone avoiding condition $(\tilde{D}, \tilde{L}) \prec (D, L)$ such that any set satisfying this condition meets Req_n .

We conclude by setting $(D_0, L_0) = (\emptyset, \omega)$, and iterating the preceding argument to construct a chain

$$(D_0, L_0) \succ (D_1, L_1) \succ \dots$$

such that, for any n , any set satisfying (D_{n+1}, L_{n+1}) meets Req_n . Taking $G = \bigcup_i D_i$, we note that $G \cap A$ and $G \cap \bar{A}$ must both be infinite by virtue of our even-numbered requirements. Now for every $e, i, s, t \in \omega$, either $\Phi_e^{G \cap A} \neq C_s$ or $\Phi_i^{G \cap \bar{A}} \neq C_t$ must hold, by virtue of the odd-numbered requirements. It follows that either $\Phi_e^{G \cap A} \neq C_s$ for every $e, s \in \omega$ or else $\Phi_i^{G \cap \bar{A}} \neq C_t$ for every $i, t \in \omega$, and we let X be $G \cap A$ or $G \cap \bar{A}$, depending on which of these two possibilities is true. Thus, X is an infinite cone avoiding subset of A or \bar{A} .

(ii) Without loss of generality, assume that neither A nor \bar{A} contains a low cone avoiding subset, since otherwise the result is trivial. We proceed as in (i), except that we force with conditions that are low as well as cone avoiding. We renumber the requirements $2e$ and $2\langle e, i, s, t \rangle + 1$ by $5e$ and $5\langle e, i, s, t \rangle + 1$, respectively, and add the requirements

- $\text{Req}_{5\langle e, i \rangle + 2}$: $(G \cap A)'(e)$ or $(G \cap \bar{A})'(i)$ is determined in the construction
- $\text{Req}_{5\langle e, i, t \rangle + 3}$: $(G \cap A)'(e)$ is determined in the construction or $\Phi_i^{G \cap \bar{A}} \neq C_t$
- $\text{Req}_{5\langle e, i, s \rangle + 4}$: $(G \cap \bar{A})'(i)$ is determined in the construction or $\Phi_e^{G \cap A} \neq C_s$.

We claim that these expanded requirements are \mathbf{d} -effectively dense. Given a low cone avoiding condition (D, L) , and an $n \in \omega$, we proceed as above for $n = 5e$ and $n = 5\langle e, i, s, t \rangle + 1$. If $n = 5\langle e, i \rangle + 2$, we appeal to Lemma 4.6, while if $n = 5\langle e, i, t \rangle + 3$ or $n = 5\langle e, i, s \rangle + 4$, we appeal to Lemma 4.8(i) and (ii), respectively. In any case, therefore, we pass to an extension (\tilde{D}, \tilde{L}) which is satisfied only by sets meeting Req_n . Moreover, since A and C are computable in $0'$, the effectiveness of Lemmas 4.5, 4.6, 4.7, and 4.8 implies that we can find (\tilde{D}, \tilde{L}) low and cone avoiding, and that we can do so \mathbf{d} -effectively.

We set $(D_0, L_0) = (\emptyset, \omega)$, and iterate the density construction to \mathbf{d} -effectively obtain a chain

$$(D_0, L_0) \succ (D_1, L_1) \succ \dots$$

with the property that any set satisfying (D_{n+1}, L_{n+1}) meets Req_n . Let $G = \bigcup_i D_i$. Then by virtue of the requirements congruent to 0, modulo 5, $G \cap A$ and $G \cap \bar{A}$ are both infinite. One of the following cases must now obtain:

Case 1: $\Phi_e^{G \cap A} \neq C_s$ and $\Phi_i^{G \cap \bar{A}} \neq C_t$ for every $e, i, s, t \in \omega$, so both $G \cap A$ and $G \cap \bar{A}$ are cone avoiding. By virtue of the requirements congruent to 2, modulo 5, for every $e, i \in \omega$, either $(G \cap A)'(e)$ or $(G \cap \bar{A})'(i)$ is determined during the construction. Hence, either $(G \cap A)'(e)$ is determined for every $e \in \omega$, or else $(G \cap \bar{A})'(i)$ is determined for every $i \in \omega$. In the former case, $(G \cap A)'$ has degree at most \mathbf{d} , and so we let $X = G \cap A$. In the latter case, let $X = G \cap \bar{A}$.

Case 2: $\Phi_e^{G \cap A} = C_s$ for some $e, s \in \omega$. The requirements congruent to 1, modulo 5, imply that for every $e, i, s, t \in \omega$, either $\Phi_e^{G \cap A} \neq C_s$ or $\Phi_i^{G \cap \bar{A}} \neq C_t$ holds. Hence $G \cap \bar{A}$ is cone avoiding. On the other hand, the requirements congruent to 4, modulo 5, imply that for every $e, i, s \in \omega$, either $(G \cap \bar{A})'(i)$ is determined in the construction or else $\Phi_e^{G \cap A} \neq C_s$. It follows, therefore, that $\Phi_i^{G \cap \bar{A}} \neq C_t$ for all $i, t \in \omega$, and that $(G \cap \bar{A})'(i)$ is determined in the construction for all $i \in \omega$, so $(G \cap \bar{A})'$ has degree at most \mathbf{d} . We set $X = G \cap \bar{A}$.

Case 3: $\Phi_i^{G \cap \bar{A}} = C_t$ for some $i, t \in \omega$. We can argue as in the previous case, but appeal instead to the requirements congruent to 1 and 3, modulo 5, and set $X = G \cap A$.

In any case, X is an infinite cone avoiding subset of A or \bar{A} and X' is determined in the construction. Since the construction is \mathbf{d} -effective, this implies that $\text{deg}(X)' \leq \mathbf{d}$, which completes the proof. \square

Proof of Lemma 3.2. (i) We prove the lemma in relativized form by induction on k , the case $k = 2$ being simply a statement of Lemma 5.2(i). Since that lemma clearly relativizes, we may assume that the present result, along with all its relativizations,

holds for some $k \geq 2$. Let A_1, \dots, A_{k+1} be a sequence of disjoint sets covering ω . Applying Lemma 5.2(i) to A_{k+1} yields an infinite cone avoiding set \tilde{X} either contained in or disjoint from A_{k+1} . In the former case, we can take \tilde{X} itself for the desired, cone avoiding subset. In the latter, we have $\tilde{X} \subseteq A_1 \cup \dots \cup A_k$, and so $A_1 \cap \tilde{X}, \dots, A_k \cap \tilde{X}$ is a sequence of k disjoint sets whose union is \tilde{X} . Relativizing our inductive hypothesis to \tilde{X} yields an infinite cone avoiding subset X of $A_i \cap \tilde{X}$ for some $i \leq k$, whence since $A_i \cap \tilde{X} \subseteq A_i$, the induction is complete.

(ii) We again use induction, with the base case, this time, following from Lemma 5.2(ii) by taking \mathbf{d} there low over $\mathbf{0}'$. Thus, assume the result for some $k \geq 2$, and let A_1, \dots, A_{k+1} be a sequence of disjoint Δ_2^0 sets covering ω . By Lemma 5.2(ii), A_{k+1} contains or disjoint from an infinite cone avoiding set \tilde{X} whose jump is computable in \mathbf{d} , hence a set which is low₂ since $\mathbf{d}' \leq \mathbf{0}''$. If $\tilde{X} \subseteq A_{k+1}$, the proof is complete. Otherwise, we apply the induction hypothesis relative to \tilde{X} to the sequence $A_1 \cap \tilde{X}, \dots, A_k \cap \tilde{X}$, finding an infinite cone avoiding subset X of some $A_i \cap \tilde{X}$ such that X is low₂ over \tilde{X} , and hence low₂. \square

6. A MINIMAL PAIR OF LOW₂ HOMOGENEOUS SETS

The goal of this section is to prove the following extension of Theorem 3.5:

Corollary 6.1. *Every computable 2-coloring of pairs admits a pair of infinite low₂ homogeneous sets whose degrees form a minimal pair.*

Fix a set B satisfying $B' \leq_T 0''$. The corollary obviously follows from Theorem 1.4 and the next theorem, the proof of which will occupy most of the rest of this section.

Theorem 6.2. *Every computable 2-coloring of pairs admits an infinite low₂ homogeneous set whose degree forms a minimal pair with $\deg(B)$.*

In our proof of Theorem 3.4(ii) above, we constructed an infinite low₂ cone avoiding homogeneous set by controlling its first jump to lie below a fixed degree $\gg \mathbf{0}'$. We forced with low cone avoiding Mathias conditions and appealed to a variant of the low basis theorem obtained by adding cone avoidance, namely Theorem 2.1. It would seem natural to replace the requirement that our conditions (D, L) be cone avoiding by the requirement that $(\deg(L), \deg(B))$ form a minimal pair. (Indeed, this latter requirement is necessary if L is introreducible, i.e. computable from all of its infinite subsets.) However, it no longer seems possible to require that L be low. (For example, if $B = 0'$ and L is even Δ_2^0 , it would follow that L is computable, which is too much to ask, since in our proof we need to apply basis theorems for Π_1^0 classes to obtain L .) The same difficulty persists even if we insist that B be low₂, which would be sufficient to obtain Corollary 6.1. For let \mathcal{P} be the Π_1^0 class of all $\{0, 1\}$ -valued diagonally noncomputable functions. Each member of \mathcal{P} computes a fixed point free function by Lemma 4.1 of Jockusch, Lerman, Soare, and Solovay [4] (cf. Soare [12], Exercise V.5.9 and Remark VII.1.9). If we take for B any Δ_2^0 element of \mathcal{P} , then by Theorem 2 of Kučera [8] no Δ_2^0 , let alone low, element of this class can have degree forming a minimal pair with $\deg(B)$.

Thus we instead prove the above theorem by directly forcing the second jump of the constructed set to be computable in $0''$, which we do by suitably extending the techniques of [1], Section 5. As in that article, one main advantage of presenting the first proof, that is, of controlling only the first jump, is that it uses a considerably

simpler notion of forcing. In our case, of course, that proof also lent itself to both a new argument for Seetapun's theorem and an answer to Question 1.5, of which the following does only the latter.

Our proof of Theorem 6.2 can, as usual, be broken up into an r-cohesive and a stable part, from which the theorem then follows analogously to the way Theorem 3.4 did from Lemmas 3.1 and 3.2. For completeness, we show how this proof goes following the statements of the necessary lemmas.

Lemma 6.3. *There exists a low₂ r-cohesive set whose degree makes a minimal pair with B .*

Here we pause to note the following extension of Theorem 2.5 of Jockusch and Stephan [5] on the existence of a low₂ cohesive degree. It follows at once from Lemma 6.3, the result of Jockusch and Stephan just cited, and the fact mentioned above (cf [5], Corollary 2.4) that the cohesive and r-cohesive degrees coincide.

Corollary 6.4. *There exists a minimal pair of low₂ cohesive degrees.*

Lemma 6.5. *Any Δ_2^0 set A contains or is disjoint from an infinite low₂ set whose degree forms a minimal pair with B .*

Proof of Theorem 6.2. Fix a computable 2-coloring of pairs f , and by Lemma 6.3, fix also a low₂ r-cohesive set X with $(\deg(X), \deg(B))$ a minimal pair. Define a stable 2-coloring of pairs g by $g(\{s, t\}) = f(\{p(s), p(t)\})$ for all $\{s, t\} \in [\omega]^2$, where p is the principal function of X . By Lemma 3.3(ii), relative to X , there exists a $\Delta_2^{0,X}$ set A such that, if S is any infinite subset of A or \bar{A} , $X \oplus S$ computes an infinite homogeneous set for g . We have $B' \leq_T 0'' \leq_T X''$, so we may relativize Lemma 6.5 to X to conclude that either A or \bar{A} contains an infinite set S such that $(S \oplus X)'' \leq_T X'' \leq_T 0''$ and every set Turing reducible to both $S \oplus X$ and B is Turing reducible to X . Letting \tilde{H} be an infinite homogeneous set for g computed by S , we find that $H = p(\tilde{H})$ is an infinite low₂ homogeneous set for f . Then H is low₂ since $H \leq_T X \oplus S$ and $X \oplus S$ is low₂. Suppose now that Y satisfies $Y \leq_T H$ and $Y \leq_T B$. Then $Y \leq_T S \oplus X$ (as $H \leq_T S \oplus X$). It follows that $Y \leq_T X$ since Y is Turing reducible to both B and $S \oplus X$. Since Y is Turing reducible to both X and B , and the degrees of X and B form a minimal pair, it follows that Y is computable, as needed to complete the proof that $\deg(H)$ and $\deg(B)$ form a minimal pair. \square

The proofs of Lemmas 6.3 and 6.5 are obtained by modifying those in Cholak, Jockusch, and Slaman [1], Section 5, of Theorems 3.3 and 3.6 of the same article. In particular, we change the forcing conditions used, add minimal pair requirements, and modify some versions of "smallness" to take account of minimal pair requirements. The proofs we are modifying are quite involved, and we have not made a large number of changes. Hence, we proceed by assuming the reader is familiar with Section 5 of [1], and instead focus on the differences. Admittedly, this is at the cost of making the proof deceptively short. The interested reader may wish to consult [1], pages 14–26, for an in-depth introduction with a view towards the underlying motivation, and pages 16–22 for the technical details.

6.1. Proving Lemma 6.3. Let $\{\pi_e(G, \vec{x})\}$ be an effective enumeration of the Π_1^0 formulas in the displayed variables, and let $\{R_e\}$ be a listing of the computable sets

such that an index of the characteristic function of R_e is $0''$ -computable from e . We seek to construct a set G meeting, for each $e \in \omega$, the following requirements:

$$\begin{aligned} \text{Req}_{4e} & : |G| \geq e \\ \text{Req}_{4e+1} & : G \subseteq^* R_e \text{ or } G \subseteq^* \overline{R_e} \\ \text{Req}_{4e+2} & : (\exists \vec{x}) \pi_e(G, \vec{x}) \text{ is decided in the construction} \\ \text{Req}_{4e+3} & : \Phi_e^G = \Phi_e^B = f \text{ total} \implies f \text{ computable.} \end{aligned}$$

We proceed as in [1], Section 5.1, but instead of forcing with low conditions we will force with those in the following definition. Otherwise, our notation and terminology will be exactly as in [1], Definitions 5.1 and 5.2, and the accompanying discussions.

Definition 6.6. A condition (D, L) is *low₂ and minimal pair forming* if L is low₂ and of degree forming a minimal pair with $\deg(B)$. An *index* for such a condition is a pair of numbers (d, ℓ) in which d is the canonical index of D and ℓ is a $\Delta_1^{0, K'}$ index for L'' .

Our main task is to adapt Lemmas 5.3, 5.4, and 5.5 of [1] to the above conditions. The first of these carries over easily. For if S is a finite set of Σ_2^0 formulas with at most G free and $\psi(G)$ is a Π_1^0 instance of some formula in S , then the proof of this lemma actually shows that any S -large condition (D, L) can be L' -effectively, finitely extended to a condition forcing $\neg\psi(G)$. This extension will clearly be low₂ and minimal pair forming if (D, L) is, in which case it can thus be found $0''$ -effectively from an index for (D, L) and the Gödel number of ψ . Lemma 5.4 of [1] is adapted as follows:

Lemma 6.7. *Let a low₂, minimal pair forming condition (D, L) of known index be given, along with the canonical index of a finite set S of Σ_2^0 formulas free in at most G . There is a $0''$ -effective procedure by which to decide, from these indices, whether or not this condition is S -large. If so, there exists an $n \in \omega$ and a sequence $(D_i, L_i, k_i, \vec{w}_i : i < n)$ witnessing this fact such that each L_i is low₂ and of degree forming a minimal pair with $\deg(B)$. Moreover, this sequence, together with indices for the (D_i, L_i) , can be found $0''$ -effectively.*

Proof. In the proof of Lemma 5.4 of [1], the definition of (D, L) being S -small is written in the form $(\exists z)(\exists Z)P(z, Z, D, L, S)$, where P is a Π_1^0 predicate with parameters Z and L of known index (we are identifying the finite sets D and S with their canonical indices). As pointed out there, the predicate $(\exists Z)P(z, Z, D, L, S)$ is consequently $\Pi_1^{0, L}$. It follows that, as a predicate of D and S , $(\exists z)(\exists Z)P(z, Z, D, L, S)$ is $\Sigma_2^{0, L}$, and we may consequently find a Δ_3^0 index for it from a $\Delta_1^{0, K'}$ index for L'' . Hence, from the appropriate indices for (D, L) and S , $0''$ can decide whether or not (D, L) is S -small. If it is, we search $0''$ -effectively for a $z \in \omega$ satisfying $(\exists Z)P(z, Z, D, L, S)$, and then, since the degrees of L and B form a minimal pair, use Theorem 2.5 relative to L to find a solution Z to $P(z, Z, D, L, S)$ which is low₂ over L , hence low₂, and whose degree forms a minimal pair with $\deg(B)$. The proof now concludes much like that of Lemma 5.4 of [1], but appealing to the effectiveness of Theorem 2.5 instead of the low basis theorem. \square

The statement of Lemma 5.5 of [1] now carries over verbatim to low₂, minimal pair forming conditions, the proof being identical except of course that all references to Lemma 5.4 of [1] are replaced by references to Lemma 6.7 above.

Under these modifications, the construction of G can go ahead, *mutatis mutandis*, as it did in [1], Section 5.1.3. For each e , the requirement $s = 4e + 3$ is met by applying a modified version of Lemma 4.4 with C in that lemma replaced by Φ_e^B to the low_2 , minimal pair forming condition (D_{s-1}, L_{s-1}) given at stage s . (From the proof of Lemma 4.4, it is not necessary to assume that C (now called Φ_e^B) is total. The case divisions are as before, and an oracle for B' suffices to find j in Case 1.)

6.2. Proving Lemma 6.5. Let A be a Δ_2^0 set, and, in keeping with the notation of Cholak, Jockusch, and Slaman [1], page 19, define $A_0 = A$ and $A_1 = \bar{A}$. Since we are forcing with conditions that are merely low_2 and not necessarily low, it is not sufficient for our needs to assume that neither A_0 nor A_1 has an infinite low subset of degree making a minimal pair with $\text{deg}(B)$. Thus, we assume that neither A_0 nor A_1 has even a low_2 such subset, and will construct a subset $G \subseteq A_i$ for some $i < 2$ to contradict this assumption.

6.2.1. Constructing G inside of A_0 . Following [1], Section 5.2.1, we begin by restricting to low_2 , minimal pair forming conditions (D, L) with $D \subset A_0$. We define what it means for a such a condition to be S -small₀ or S -large₀ as in [1], page 19. However, in order to ensure that the set G we construct has degree forming a minimal pair with $\text{deg}(B)$, we additionally need the following secondary notion of largeness:

Definition 6.8. Let

$$S = \{(\exists \vec{x})\psi_0(G, \vec{x}), \dots, (\exists \vec{x})\psi_k(G, \vec{x})\}$$

be a finite set of Σ_2^0 formulas free in at most G , the ψ_i being Π_1^0 , and let $e \in \omega$. A condition (D, L) is (S, e) -small₀ if there exists an $n \in \omega$ and a sequence $(D_i, L_i, k_i, \vec{w}_i : i < n)$ such that the L_i partition L , each k_i is a number $\leq k$ and each \vec{w}_i a tuple of numbers, each (D_i, L_i) is a precondition extending (D, L) with $D_i \subset A_0$, and for each $i < n$, either $\max(L_i) \leq \max(\vec{w}_i)$, or (D_i, L_i) forces $\psi_{k_i}(G, \vec{w}_i)$, or (D_i, L_i) forces agreement on e . A condition is (S, e) -large₀ if it is not (S, e) -small₀. (This definition is the same as the definition of “ S -small” except for the final conjunct, which is included for the sake of the e th minimal pair requirement.)

Notice that if a condition is (S, e) -large₀ then it is S -large₀, and every finite extension of it is also (S, e) -large₀. We begin by assuming that whenever a condition (D, L) is S -large₀, then $(D, L \cap A_0)$ is (S, e) -large₀ for all $e \in \omega$, so that, in particular, whenever (D, L) is an S -large₀ condition, so is $(D, L \cap A_0)$. If this assumption holds, we will construct a suitable $G \subseteq A_0$, and otherwise we will construct $G \subseteq A_1$. By this assumption, Lemma 5.3 can be adapted to conditions (D, L) with $D \subset A_0$ just as in Section 5.2.1 of [1]. The following gives the necessary adaptation of Lemma 5.4 of [1] for our purposes:

Lemma 6.9. *Let an index for a low_2 , minimal pair forming condition (D, L) with $D \subset A_0$ be given, along with an $e \in \omega$, and the canonical index of a finite set S of Σ_2^0 formulas free in at most G . There are $0''$ -effective procedures by which to decide, from these indices, whether or not this condition is S -small₀, and, from e and these indices, whether or not it is (S, e) -small₀. If it is S -small₀ or (S, e) -small₀, there exists an $n \in \omega$ and a sequence $(D_i, L_i, k_i, \vec{w}_i : i < n)$ witnessing this fact such that each $D_i \subset A_0$, and each L_i is low_2 and of degree forming a minimal*

pair with $\deg(B)$. Moreover, this sequence, together with indices for the (D_i, L_i) , can be found $0''$ -effectively.

Proof. In Section 5.2.1 of [1] the definition of (D, L) being S -small₀ is put in the form $(\exists z)[R^A(z) \wedge (\exists Z)P(z, Z, D, L, S)]$, where R^A is a $\Delta_1^{0,A}$ predicate, and P is the same Π_1^0 predicate defined in the proof of Lemma 6.7. The definition of (D, L) being (S, e) -small₀ can similarly be put in the form $(\exists z)[\tilde{R}^A(z) \wedge (\exists Z)\tilde{P}(z, Z, e, D, L, S)]$, where \tilde{R}^A is $\Delta_2^{0,A}$ and \tilde{P} is Π_1^0 . Thus, (D, L) being S -small₀ and (D, L) being (S, e) -small₀ can each be defined by some $\Sigma_1^{0, A \oplus L'}$ predicate, and hence by one computable in $(A \oplus L)'$. But since $A \leq_T 0' \leq_T L'$, and since L is low₂, it follows that $(A \oplus L)'\leq_T 0''$, and hence that each of these predicates is still Δ_3^0 . The proof can now be completed just as that of Lemma 6.7. \square

Now Lemma 5.5 of [1] can be adapted in the obvious way to low₂, minimal pair forming conditions (D, L) with $D \subset A_0$. Its proof is the same, except that Lemma 6.9 above is used in place of Lemma 5.4 of [1].

Finally, we have the following lemma which will ensure that the set we end up constructing is of degree forming a minimal pair with $\deg(B)$:

Lemma 6.10. *Let (D, L) be a low₂, minimal pair forming condition which is S -large₀, and let $e \in \omega$. There exists a low₂, minimal, pair forming condition extending (D, L) which is S -large₀, and which is satisfied only by sets G such that Φ_e^G is computable whenever it is total and equal to Φ_e^B . Furthermore, an index for this extension can be found $0''$ -effectively from e and an index for (D, L) .*

Proof. By Lemma 6.9, we can $0''$ -effectively from e and an index for (D, L) determine whether or not this condition is (S, e) -small₀.

If it is, we can $0''$ -effectively find an $n \in \omega$ and a sequence $(D_i, L_i, k_i, \vec{w}_i : i < n)$ witnessing this in which each (D_i, L_i) is low₂ and minimal pair forming. There must clearly be some $i < n$ so that L_i is infinite and (D_i, L_i) forces agreement on e , else we could easily form a sequence of preconditions to witness that (D, L) is S -small₀. Furthermore, if each such (D_i, L_i) were S -small₀, then we could replace each one by the preconditions from the sequence witnessing this fact, thereby again obtaining that (D, L) is S -small₀. Thus, we may $0''$ -effectively choose i so that L_i is infinite and (D_i, L_i) is S -large₀ and forces agreement on e . This condition can serve as the desired extension, since if G satisfies it then Φ_e^G is clearly L -computable whenever it is total.

If (D, L) is (S, e) -large₀ then by assumption so is $(D, L \cap A_0)$, meaning in particular that it cannot force agreement on e . We can thus $0''$ -effectively find an $x \in \omega$ and a finite set F satisfying $(D, L \cap A_0)$ so that $\Phi^F(x) \downarrow \neq \Phi^B(x)$. Set $\tilde{D} = D \cup F$ and let \tilde{L} consist of all elements of F greater than $\max(F)$ and use $(\Phi^F(x))$. Clearly, (\tilde{D}, \tilde{L}) serves as the desired extension. \square

We now construct an infinite set $G \subseteq A_0$ which is low₂ and of degree forming a minimal pair with $\deg(B)$ analogously to the way we constructed an r -cohesive such set above, only forcing with large₀ conditions in place of large ones. Our assumption that A_1 has no low₂ subset of degree forming a minimal pair with $\deg(B)$ ensures that we meet requirement Req_{4e} of Section 6.1 above, our adaptations of Lemmas 5.3 and 5.5 of [1] ensure that we meet requirement Req_{4e+2} , and Lemma 6.10 ensures that we meet Req_{4e+3} .

6.2.2. *Constructing G inside of A_1 .* Assume next that there exists a low₂, minimal pair forming condition (\tilde{D}, \tilde{L}) , an $\tilde{e} \in \omega$, and a finite set

$$\tilde{S} = \{(\exists \vec{x})\tilde{\psi}_0(G, \vec{x}), \dots, (\exists \vec{x})\tilde{\psi}_\ell(G, \vec{x})\}$$

of Σ_2^0 formulas free in at most G , such that $\tilde{D} \subset A_0$, (\tilde{D}, \tilde{L}) is \tilde{S} -large₀, but $(\tilde{D}, \tilde{L} \cap A_0)$ is (\tilde{S}, \tilde{e}) -small₀. This is a slightly different assumption from that of [1], Section 5.2.2, resulting from the fact that we needed Definition 6.8 above in order to obtain Lemma 6.10. Nonetheless, essentially the same argument as there will work here, with only some minor technical alterations along the way. The basic idea is that we switch from trying to build the desired set G inside of A_0 to building it inside its complement, A_1 .

We need to suitably modify the notion from [1], page 20, of a condition being 1-acceptable, and then that from [1], Definition 5.6, of a condition being S -small₁. Here we remark in passing that the latter is misstated in [1], and should, for the purposes of Section 5.2.2 of that article, be defined as in part (iii) of the following definition, but with (5) below not obtaining for any $i < n$.

Definition 6.11. Let a condition (D, L) , an $e \in \omega$, and a finite set

$$S = \{(\exists \vec{x})\psi_0(G, \vec{x}), \dots, (\exists \vec{x})\psi_k(G, \vec{x})\}$$

of Σ_2^0 formulas free in at most G , be given.

- (i) We say (D, L) is 1-acceptable if $D \subset A_1$, $L \subseteq \tilde{L}$, and (\tilde{D}, L) is (\tilde{S}, \tilde{e}) -large₀.
- (ii) We say (D, L) is (S, e) -small₁ if it is 1-acceptable and there exists an $n \in \omega$ and a sequence $(D_i, L_i, k_i, \vec{w}_i : i < n)$ such that the L_i partition L , each k_i is a number $\leq \max(k, \ell)$ and each \vec{w}_i a tuple of numbers, each (D_i, L_i) is a precondition, and for each $i < n$, at least one of the following holds:
 - (1) $\max(L_i) \leq \max(\vec{w}_i)$,
 - (2) $D_i \subset A_1$, and (D_i, L_i) extends (D, L) and forces $\psi_{k_i}(G, \vec{w}_i)$,
 - (3) $D_i \subset A_1$, and (D_i, L_i) extends (D, L) and forces agreement on e ,
 - (4) $D_i \subset A_0$, and (D_i, L_i) extends (\tilde{D}, \tilde{L}) and forces $\tilde{\psi}_{k_i}(G, \vec{w}_i)$,
 - (5) $D_i \subset A_0$, and (D_i, L_i) extends (\tilde{D}, \tilde{L}) and forces agreement on \tilde{e} .
- (iii) If (D, L) is (S, e) -small₁ via a sequence for which (3) above does not obtain for any $i < n$, then we say (D, L) is S -small₁.
- (iv) We say (D, L) is (S, e) -large₁, respectively S -large₁, if it is not (S, e) -small₁, respectively if it is not S -small₁.

Any condition which is (S, e) -large₁ obviously enjoys analogs of the two basic properties mentioned following Definition 6.8 for (S, e) -large₀ conditions. It is also easily seen that if (D, L) is 1-acceptable and finitely extended by (\hat{D}, \hat{L}) where $\hat{D} \subset A_1$, then this extension is also 1-acceptable. The following analog of Lemma 5.7 of [1] now ensures that we will not at any future stage be forced to switch back to constructing G inside of A_0 .

Lemma 6.12. *Let S be a finite set of Σ_2^0 formulas free in at most G , and let $e \in \omega$. If (D, L) is S -large₁, then so is $(D, L \cap A_1)$. Likewise, if (D, E) is (S, e) -large₁, then so is $(D, L \cap A_1)$.*

Proof. By assumption, $(\tilde{D}, \tilde{L} \cap A_0)$ is (\tilde{S}, \tilde{e}) -small₀, so fix a sequence witnessing this fact. By comparing Definitions 6.8 and 6.11, it is easy to see that this sequence witnesses that $(D, L \cap A_0)$ is S -small₁ for any S , and hence (S, e) -small₁ for any

S and e . This sequence could consequently be combined with any sequence witnessing that $(D, L \cap A_1)$ is S -small₁ or (S, e) -small₁ to witness that (D, L) is itself, respectively, S -small₁ or (S, e) -small₁. \square

We can thus adapt Lemma 5.3 of [1] to 1-acceptable, low₂, minimal pair forming conditions (D, L) with $D \subset A_1$ almost exactly as we adapted it in Section 6.2.1 to conditions with $D \subset A_0$. Similarly, Lemma 6.9 carries over simply by replacing all instances of A_0 , small₀ and large₀ by A_1 , small₁ and large₁, respectively. This leaves Lemma 5.5 of [1] and Lemma 6.10, the proofs of which require a minor case analysis in order to work for 1-acceptable conditions. This is essentially what is required to adapt the former lemma in Section 5.2.2 of [1], but since it is not explicitly described there, we provide the details. The latter lemma can be handled similarly.

Lemma 6.13. *Let (D, L) be a low₂, minimal pair forming condition which is 1-acceptable, S -large₁, and $S \cup \{(\exists \vec{x})\psi(G, \vec{x})\}$ -small₁ for some Π_1^0 formula $\psi(G, \vec{x})$ free in at most G and \vec{x} . There exists a low₂, minimal pair forming extension of this condition which is 1-acceptable and S -large₁, and which forces a Π_1^0 instance of $\psi(G, \vec{x})$. An index of this extension can be found 0''-effectively from an index for (D, L) , the canonical index for S , and the Gödel number of ψ .*

Proof. Fix $n \in \omega$ and a sequence $(D_i, L_i, k_i, \vec{w}_i : i < n)$ with each (D_i, L_i) low₂ and minimal pair forming to witness that (D, L) is $S \cup \{(\exists \vec{x})\psi(G, \vec{x})\}$ -small₁. First, notice that if no (D_i, L_i) with L_i infinite had $D_i \subset A_1$ then our sequence would in fact witness that (D, L) is \hat{S} -small₁ for any set \hat{S} , and hence in particular that it is S -small₁. Now let I_0 consist of all $i < n$ such that L_i is infinite and (D_i, L_i) is 1-acceptable, noting that this must be non-empty else each condition (D_i, L_i) with L_i infinite and $D_i \subset A_1$ could be replaced in our sequence by preconditions from a sequence witnessing that (\tilde{D}, L_i) is (\tilde{S}, \tilde{e}) -small₀ to obtain a sequence witnessing that (\tilde{D}, L) is (\tilde{S}, \tilde{e}) -small₀ and hence that (D, L) is not 1-acceptable. Clearly, each (D_i, L_i) with $i \in I_0$ forces a Π_1^0 instance of some formula in $S \cup \{(\exists \vec{x})\psi(G, \vec{x})\}$. Let I_1 consist of all $i \in I_0$ with (D_i, L_i) S -large₁. If I_1 were empty, then, depending on whether i is or is not an element of I_0 , each (D_i, L_i) with $D_i \subset A_1$ could be replaced in our original sequence either by preconditions from a sequence witnessing that (\tilde{D}, L_i) is S -small₀, or else by preconditions from a sequence witnessing that (D_i, L_i) is S -small₁, yielding a sequence witnessing that (D, L) is S -small₁. By the effectiveness of Lemma 6.9, we may therefore 0''-effectively from the appropriate indices find an $i \in I_1$. Since (D_i, L_i) is S -large₁, it cannot force a Π_1^0 instance of any formula in S , and hence it must force $\psi(G, \vec{w}_i)$. \square

All the lemmas are now combined in the usual way to build the desired $G \subseteq A_1$, but forcing with low₂, minimal pair forming conditions which are 1-acceptable and large₁, starting with the \emptyset -large₁ condition (\emptyset, \tilde{L}) .

7. QUESTIONS

On a visit to Chicago during the preparation of this article, Stephen Simpson asked us the following interesting questions pertaining to Theorem 3.5 and Corollary 6.1.

Question 7.1 (Simpson). Given two noncomputable sets C_0 and C_1 , does every computable 2-coloring of pairs admit an infinite low_2 homogeneous set H that is cone avoiding for both?

Question 7.2 (Simpson). Does every computable 2-coloring of pairs admit a pair of infinite homogeneous sets whose degrees form a minimal pair and whose join is low_2 ?

In response to the first of these, we have the following strengthening of Theorem 3.5:

Theorem 7.3. *Let C_0, \dots, C_n be a finite sequence of noncomputable sets. Every computable 2-coloring f of pairs admits an infinite low_2 homogeneous set which is cone avoiding for this sequence.*

Proof. Write $\{0, \dots, n\}$ as a disjoint union $I \cup J \cup K$, where $C_i \leq_T 0'$ for all $i \in I$, $C_j \not\leq_T 0'$ and $C_j \leq_T 0''$ for $j \in J$, and $C_k \not\leq_T 0''$ for $k \in K$. By Theorem 2.1, relativized to $0'$ and applied to the $0'$ -computable tree all of whose paths have degree $\gg 0'$, we can find a degree $\mathbf{d} \gg 0'$ such that $\mathbf{d}' \leq 0''$ and $\deg(C_j) \not\leq \mathbf{d}$ for all $j \in J$ (if $J = \emptyset$, let \mathbf{d} be any degree $\gg 0'$ which is low over $0'$). Now by Theorem 3.4(ii), there exists an infinite homogeneous set H for f such that $\deg(H)' \leq \mathbf{d}$ and $C_i \not\leq_T H$ for all $i \in I$ (if $I = \emptyset$, let H be any infinite homogeneous set with jump of degree at most \mathbf{d}). Since H is low_2 we also have $C_k \not\leq_T H$ for all $k \in K$, as wanted. \square

We do not know the answer to Question 7.2. A natural attempt at obtaining an affirmative answer is to assume that the set B in Section 6 is low_2 , rather than merely that $B' \leq_T 0''$, and then to try to show that the homogeneous set asserted to exist by Theorem 6.2 can be made to satisfy $(H \oplus B)'' \leq_T 0''$. Our obstacle to doing this is that we do not know how to strengthen Theorem 2.5 from asserting the existence of a low_2 path to asserting the existence of one whose join with B is low_2 .

As already mentioned at the end of Section 3, we do not know the answer to the next question.

Question 7.4. Given a noncomputable set C , does every computable 2-coloring of pairs admit an infinite homogeneous set $H \not\leq_T C$ whose jump is of degree at most \mathbf{d} , where \mathbf{d} is a given degree $\gg 0'$?

In a similar vein, we can ask the following:

Question 7.5. Given a degree $\mathbf{d} \gg 0'$, does every computable 2-coloring of pairs admit a pair of infinite homogeneous sets whose degrees form a minimal pair, and whose jumps are each of degree at most \mathbf{d} ?

In Section 6 we control the second jump of the constructed set, and so it does not appear that this method would be useful in finding an affirmative answer to either of the above questions. But it does not appear that the methods of Sections 4 and 5 could be easily adapted to give affirmative answers either. If we tried to do so for the first question, we would certainly need a different case analysis in the proof of Theorem 3.5, but this would in turn demand a different proof of Theorem 2.1. As for the second question, we would seem to need a version of Theorem 2.5 which is effective in a given degree $\gg 0'$, but we do not know of a proof of this theorem which does not make essential use of a $0''$ oracle.

REFERENCES

- [1] Peter A. Cholak, Carl G. Jockusch, Jr., and Theodore A. Slaman, *On the strength of Ramsey's theorem for pairs*, J. Symbolic Logic **66** (2001), no. 1, 1–55.
- [2] Denis R. Hirschfeldt, Carl G. Jockusch, Jr., Bjørn Kjos-Hanssen, Steffen Lempp, and Theodore A. Slaman, *The strength of some combinatorial principles related to Ramsey's theorem for pairs*, pp. 143-161 in Computational Prospects of Infinity, Part II: Presented Talks, Lecture Notes Series, Institute for Mathematical Sciences, National University of Singapore, Vol. 15, edited by C.T. Chong, Q. Feng, T. Slaman, H. Woodin, and Y. Yang, published by World Scientific, New Jersey, London, Singapore, Beijing, Shanghai, Hong Kong, Taipei, Chennai, 2008.
- [3] Tamara Lakins Hummel, *Effective versions of Ramsey's theorem: avoiding the cone above $\mathbf{0}'$* , J. Symbolic Logic **59** (1994), no. 4, 1301–1325.
- [4] Carl G. Jockusch, Jr., Manuel Lerman, Robert I. Soare, and Robert M. Solovay, *Recursively enumerable sets modulo iterated jumps and extensions of Arslanov's completeness criterion*, J. Symbolic Logic **54** (1989), no. 4, 1288–1323.
- [5] Carl G. Jockusch, Jr. and Frank Stephan, *A cohesive set which is not high*, Math. Logic Quart. **39** (1993), no. 4, 515–530.
- [6] Carl G. Jockusch, Jr., *Ramsey's theorem and recursion theory*, J. Symbolic Logic **37** (1972), no. 2, 268–280.
- [7] Carl G. Jockusch, Jr. and Robert I. Soare, Π_1^0 *classes and degrees of theories*, Trans. Amer. Math. Soc. **173** (1972), 33–56.
- [8] Antonín Kučera, *An alternative, priority-free solution to Post's problem*, Mathematical foundations of computer science, 1986 (Bratislava, 1986), Lecture Notes in Comput. Sci., vol. 233, Springer, Berlin, 1986, pp. 493–500.
- [9] David Seetapun and Theodore A. Slaman, *On the strength of Ramsey's theorem*, Notre Dame J. Formal Logic **36** (1995), no. 4, 570–582, Special Issue: Models of arithmetic.
- [10] Stephen G. Simpson, *Degrees of unsolvability: a survey of results*, Handbook of Mathematical Logic (J. Barwise, ed.), North-Holland, Amsterdam, 1977, pp. 631–652.
- [11] Stephen G. Simpson, *Subsystems of second order arithmetic*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1999.
- [12] Robert I. Soare, *Recursively enumerable sets and degrees*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1987, A study of computable functions and computably generated sets.

UNIVERSITY OF CHICAGO

E-mail address: damir@math.uchicago.edu

UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN

E-mail address: jockusch@math.uiuc.edu