

# **Chapter 3: Higher-Order Linear Diff. Equations**

## **Section 3.5: Non-Homogeneous Equations and Undermined Coefficients**

# Theorem

- Theorem: Let  $y_p$  be a particular solution of the nonhomogeneous equation

$$y^{(n)} + p_1(x)y^{(n-1)} + \cdots + p_{n-1}(x)y' + p_n(x)y = f$$

on an open interval  $I$  where each  $p_i$  and  $f$  are continuous. Let  $y_1, y_2, \dots, y_n$  be  $n$  linear independent solutions of the homogeneous  $n$ -th -order linear equation  $y^{(n)} + p_1(x)y^{(n-1)} + \cdots + p_{n-1}(x)y' + p_n(x)y = 0$ . If  $Y$  is any solution whatsoever of above non-homogeneous equation on  $I$ , then there exist numbers  $c_1, c_2, \dots, c_n$  such that

$$Y(x) = c_1y_1(x) + c_2y_2(x) + \cdots + c_ny_n(x) + y_p(x)$$

for all  $x$  in  $I$ .

# Rule 1

$$Ly = f$$

Suppose that no term appearing either in  $f(x)$  or in any of its derivatives satisfies the associated homogeneous equation  $Ly = 0$ . Then take as a trial solution for  $y_p$  a linear combination of all linearly independent such terms and their derivatives. Then determine the coefficients by substitution of this trial solution into the nonhomogeneous equation

$$Ly = f(x).$$

Example:  $y'' + 3y' + 4y = 3x + 2$

Example:  $y'' - 4y = 2e^{3x}$

Example:  $y'' + 4y = 3x^3$

Example:  $y'' - 3y' + 2y = 3e^{-x} - 10 \cos 3x$  with  $y(0) = 1$  and  $y'(0) = 2$

Example:  $y^{(3)} + 9y' = x \sin x + x^2 e^{2x}$

# Rule 2

$$Ly = f$$

If the function  $f(x)$  is of either form

$$P_m(x)e^{rx} \cos kx \quad \text{or} \quad P_m(x)e^{rx} \sin kx,$$

take as the trial solution

$$y_p(x) = x^s [(A_0 + A_1x + A_2x^2 + \cdots + A_mx^m)e^{rx} \cos kx \\ + (B_0 + B_1x + B_2x^2 + \cdots + B_mx^m)e^{rx} \sin kx]$$

where  $s$  is the smallest nonnegative integer such that no term in  $y_p$  duplicates a term in the complementary function  $f_c$ . Then determine the coefficients in above equation by substituting  $y_p$  into the nonhomogeneous equation.

terms in  $f(x)$  and their derivatives

$y_p(x)$

$$P_m = \mathbf{a}_0 + \mathbf{a}_1x + \mathbf{a}_2x^2 + \cdots + \mathbf{a}_mx^m$$

$$x^s (A_0 + A_1x + A_2x^2 + \cdots + A_mx^m)$$

$$\mathbf{a} \cos kx + \mathbf{b} \sin kx$$

$$x^s (A \cos kx + B \sin kx)$$

$$e^{rx}(\mathbf{a} \cos kx + \mathbf{b} \sin kx)$$

$$x^s e^{rx} (A \cos kx + B \sin kx)$$

$$P_m e^{rx}$$

$$x^s (A_0 + A_1x + A_2x^2 + \cdots + A_mx^m) e^{rx}$$

$$P_m (\mathbf{a} \cos kx + \mathbf{b} \sin kx)$$

$$x^s [(A_0 + A_1x + A_2x^2 + \cdots + A_mx^m) \cos kx \\ + (B_0 + B_1x + B_2x^2 + \cdots + B_mx^m) \sin kx]$$

$$P_m (\mathbf{a} \cos kx + \mathbf{b} \sin kx) e^{rx}$$

$$x^s [(A_0 + A_1x + A_2x^2 + \cdots + A_mx^m) e^{rx} \cos kx \\ + (B_0 + B_1x + B_2x^2 + \cdots + B_mx^m) e^{rx} \sin kx]$$

terms in  $f(x)$  and their derivatives

$y_p(x)$

$$P_m = \mathbf{a}_0 + \mathbf{a}_1x + \mathbf{a}_2x^2 + \cdots + \mathbf{a}_mx^m$$

$$x^s (A_0 + A_1x + A_2x^2 + \cdots + A_mx^m)$$

$$\mathbf{a} \cos kx + \mathbf{b} \sin kx$$

$$x^s (A \cos kx + B \sin kx)$$

$$e^{rx}(\mathbf{a} \cos kx + \mathbf{b} \sin kx)$$

$$x^s e^{rx} (A \cos kx + B \sin kx)$$

$$P_m e^{rx}$$

$$x^s (A_0 + A_1x + A_2x^2 + \cdots + A_mx^m) e^{rx}$$

$$P_m (\mathbf{a} \cos kx + \mathbf{b} \sin kx)$$

$$x^s [(A_0 + A_1x + A_2x^2 + \cdots + A_mx^m) \cos kx \\ + (B_0 + B_1x + B_2x^2 + \cdots + B_mx^m) \sin kx]$$

$$P_m (\mathbf{a} \cos kx + \mathbf{b} \sin kx) e^{rx}$$

$$x^s [(A_0 + A_1x + A_2x^2 + \cdots + A_mx^m) e^{rx} \cos kx \\ + (B_0 + B_1x + B_2x^2 + \cdots + B_mx^m) e^{rx} \sin kx]$$

# Examples

Determine the appropriate form for a particular solution of

Example:  $y^{(3)} + y'' = 3e^x + 4x^2$

Example:  $y'' + 6y' + 13y = e^{-3x} \cos 2x$

Example:  $y'' - 6y' + 13y = xe^{3x} \sin 2x$

Example:  $(D - 1)^3(D^2 - 4)y = xe^x + e^{2x} + e^{-2x}$

# Variation of Parameters

Consider

$$Ly = f$$

where  $L$  is a differential operator with constant coefficients. Let  $y_1, y_2, \dots, y_n$  be a linearly independent solution of  $Ly = 0$ .

Question: Whether it is possible to choose a particular solution in such a way that the combination

$$y_p(x) = u_1(x)y_1(x) + u_2(x)y_2(x) + \dots + u_n(x)y_n(x)$$

is a particular solution of  $Ly = f$ .

Example:  $y'' + y = \tan x$

# Variation of Parameters

Theorem: If the nonhomogeneous equation

$$y'' + P(x)y' + Q(x)y = f(x)$$

has complementary function  $y_c(x) = c_1y_1(x) + c_2y_2(x)$ , then a particular solution is given by

$$y_p(x) = -y_1(x) \int \frac{y_2(x)f(x)}{W(x)} dx + y_2(x) \int \frac{y_1(x)f(x)}{W(x)} dx$$

where  $W = W(y_1, y_2)$  is the Wronskian of the two independent solutions  $y_1$  and  $y_2$  of the associated homogeneous equation.

Example:  $y'' + y' - 2y = 4e^x$

Example:  $y'' + y = \tan x$

Example:  $y'' + y = \csc^2 x$