

## Sixth Homework Set — Solutions

### Chapter 4

Problem 55

$$\begin{aligned}
 P(\text{no errors}) &= P(\text{no errors}|\text{first typist})P(\text{first typist}) \\
 &\quad + P(\text{no errors}|\text{second typist})P(\text{second typist}) \\
 &= \frac{1}{2} \left( \frac{3^0}{0!}e^{-3} + \frac{4 \cdot 2^0}{0!}e^{-4.2} \right) \\
 &= \frac{1}{2} (e^{-3} + e^{-4.2}).
 \end{aligned}$$

Problem 57  $X$  is Poisson with parameter  $\lambda = 3$ .

$$\begin{aligned}
 \text{(a)} \quad P\{X \geq 3\} &= 1 - P\{0\} - P\{1\} - P\{2\} = 1 - e^{-3} \left( 1 + 3 + \frac{9}{2} \right) = 0.5768. \\
 \text{(b)} \quad P\{X \geq 3|X \geq 1\} &= \frac{P\{X \geq 3\}}{P\{X \geq 1\}} = \frac{P\{X \geq 3\}}{1 - e^{-3}} = 0.6070.
 \end{aligned}$$

Problem 59 Let  $X$  be the number of times you win a prize. Then  $X$  is binomial with  $n = 50$  and  $p = \frac{1}{100}$ , i.e., we can use the Poisson approximation with  $\lambda = 50 \cdot \frac{1}{100} = \frac{1}{2}$ .

$$\begin{aligned}
 \text{(a)} \quad P\{X \geq 1\} &= 1 - P\{X = 0\} = 1 - e^{-\frac{1}{2}} = 0.3935 \\
 \text{(b)} \quad P\{X = 1\} &= \frac{1}{2}e^{-\frac{1}{2}} = 0.3033 \\
 \text{(c)} \quad P\{X \geq 2\} &= 1 - P\{X = 0\} - P\{X = 1\} = 1 - e^{-\frac{1}{2}} \left( 1 + \frac{1}{2} \right) = 0.0902
 \end{aligned}$$

Problem 61 Let  $X$  be Poisson with parameter  $\lambda = 1000 \cdot 0.0014 = 1.4$ . Then  $P\{X \geq 2\} = 1 - P\{X = 0\} - P\{X = 1\} = 1 - e^{-1.4}(1 + 1.4) = 0.4082$ .

Problem 63 Let  $X$  be a Poisson random variable with parameter  $\lambda = \frac{5}{2}$ . Then  $X$  gives a reasonable description of the number of people entering the casino between 12 and 12:05.

$$\begin{aligned}
 \text{(a)} \quad P\{X = 0\} &= e^{-\frac{5}{2}} = 0.0821 \\
 \text{(b)} \quad P\{X \geq 4\} &= 1 - e^{-\frac{5}{2}} \left( 1 + \frac{5}{2} + \frac{25}{8} + \frac{125}{48} \right) = 0.2424
 \end{aligned}$$

Problem 72 Let  $A$  be the stronger team.  $P(A \text{ wins in } i \text{ games}) = \binom{i-1}{i-4} 0.6^i 0.4^{i-4}$ , for  $i = 4, \dots, 7$ . Hence

$$P(A \text{ wins best-of-seven series}) = \sum_{i=4}^7 \binom{i-1}{i-4} 0.6^4 0.4^{i-4} = 0.7102.$$

Similarly,

$$P(A \text{ wins best-of-three series}) = \sum_{i=2}^3 \binom{i-1}{i-2} 0.6^4 0.4^{i-2} = 0.6480.$$

Problem 73 Let  $X$  be the number of games played in a match. Then  $P\{X = i\} = 2 \binom{i-1}{i-4} \left(\frac{1}{2}\right)^i$  for  $i = 4, \dots, 7$ . Hence,  $E[X] = 2 \sum_{i=4}^7 i \binom{i-1}{i-4} \left(\frac{1}{2}\right)^i = 5.8125$ .

Problem 77 Let  $E$  be the event that right-hand box is emptied while the left-hand box still contains  $k$  matches. Then, using a negative binomial random variable with  $p = \frac{1}{2}$ ,  $r = N$ , and  $n = 2N - k$ , we see that  $P(E) = \binom{2N-k-1}{N-1} \left(\frac{1}{2}\right)^{2N-k}$ . Now the desired probability is  $2P(E)$ .

Problem 78 Let  $E$  be the event that a single drawing results in two white and two black balls. Then  $P(E) = \frac{\binom{4}{2} \binom{4}{2}}{\binom{8}{4}} = \frac{18}{35}$ .

Let  $X$  be the number of selections until  $E$  occurs. Then

$$P\{X = n\} = \frac{17^{n-1} \cdot 18}{35^n}.$$

Problem 79 (a)  $P\{X = 0\} = \frac{\binom{94}{10}}{\binom{100}{10}} = 0.5223$

(b)

$$\begin{aligned} P\{X > 2\} &= 1 - P\{X = 0\} - P\{X = 1\} - P\{X = 2\} \\ &= \frac{\binom{100}{10} - \binom{94}{10} - \binom{6}{1} \binom{94}{9} - \binom{6}{2} \binom{94}{8}}{\binom{100}{10}} = 0.0126 \end{aligned}$$