

# Hyperbolic and quasisymmetric structure of hyperspaces

Leonid V. Kovalev and Jeremy T. Tyson

ABSTRACT. A hyperspace is a space of nonempty closed sets equipped with the Hausdorff metric. Among the subjects considered in this paper are Gromov hyperbolicity, quasisymmetric equivalence and bi-Lipschitz embeddings of hyperspaces.

## 1. Introduction

Let  $(X, d)$  be a metric space. We denote by  $\text{CL}(X)$  the hyperspace of nonempty closed subsets of a metric space  $X$ , equipped with the Hausdorff distance

$$D(A, B) = \inf\{\epsilon > 0 : A \subset N_\epsilon(B) \text{ and } B \subset N_\epsilon(A)\},$$

where  $N_\epsilon(A) = \{x \in X : \text{dist}(x, A) \leq \epsilon\}$  is the closed  $\epsilon$ -neighborhood of  $A \subset X$ . If there is no such  $\epsilon$ , the distance between  $A$  and  $B$  is infinite. Notice that if  $D(A, B) < \infty$ , then the above infimum is attained. The distance  $D$  is a metric on  $\text{CL}(X)$  if and only if  $X$  is bounded. For unbounded  $X$  the set  $\text{CL}(X)$  naturally splits into an infinite collection of disjoint metric spaces. The most notable of these is  $\mathcal{H}(X)$ , the space of nonempty bounded closed subsets of  $X$ . However, we are also interested in other components of  $\text{CL}(X)$ . Each of them has a form  $\mathcal{H}(X; C) := \{A \in \text{CL}(X) : D(A, C) < \infty\}$  for some  $C \in \text{CL}(X)$ . Note that  $\mathcal{H}(X; C) = \mathcal{H}(X)$  if and only if  $C$  is bounded. Also, let  $\mathcal{H}^c(X) = \{A \in \mathcal{H}(X) : A \text{ is geodesically convex}\}$  denote the hyperspace of bounded closed convex sets. An open ball with center  $x$  and radius  $r$  will be denoted by  $B(x, r)$ .

Most of the existing research on hyperspaces equipped with the Hausdorff distance is focused on  $\mathcal{H}(X)$  and its subsets such as  $\mathcal{H}^c(X)$ . However, the spaces  $\mathcal{H}(X; C)$  with  $C$  unbounded do arise naturally. As an example, consider a Lipschitz quotient mapping  $f : X \rightarrow Y$ , where  $X$  and  $Y$  are metric spaces. Recall that  $f$  is called a Lipschitz quotient [4] if there exist two constants  $0 < l \leq L < \infty$  such that

$$B(f(x), lr) \subset f(B(x, r)) \subset B(f(x), Lr), \quad x \in X, r > 0.$$

A Lipschitz quotient  $f : X \rightarrow Y$  induces a map  $G_f : \text{CL}(Y) \rightarrow \text{CL}(X)$  such that  $G_f(A) = f^{-1}(A)$ . It is easy to check that  $G_f$  is bi-Lipschitz on each component of  $\text{CL}(Y)$ . However, in many interesting cases  $G_f$  does not map  $\mathcal{H}(Y)$  into  $\mathcal{H}(X)$ ;

---

2000 *Mathematics Subject Classification*. Primary 54E35; Secondary 54E40, 51F99.

Research of J. T. T. supported by the National Science Foundation under grant DMS 0228807.

instead  $G_f(\mathcal{H}(Y))$  is contained in  $\mathcal{H}(X; C)$  for some unbounded set  $C$ . Even the simplest elements of  $\mathcal{H}(Y)$ , namely singletons  $\{y\}$ , can be mapped by  $G_f$  into sets with rather complicated structure [11, 26]. It remains an open question whether  $G_f(\{y\})$  is always a discrete set when  $X = Y = \mathbb{R}^n$ ,  $n > 2$ . See [4, 19, 22] for more on this problem.

The spaces  $\mathcal{H}(X; C)$  with  $C$  unbounded are typically larger and more complex than  $\mathcal{H}(X)$ . For instance,  $\mathcal{H}(X)$  is separable whenever  $X$  is *proper* (i.e. such that all of its closed bounded subsets are compact). By contrast,  $\mathcal{H}(X; C)$  is non-separable if  $X$  is connected and  $C$  is unbounded (Theorem 5.1). Furthermore, large classes of metric spaces admit isometric or bi-Lipschitz embeddings into  $\mathcal{H}(X; C)$ .

Section 2 is concerned with hyperspaces of special classes of metric spaces: Gromov hyperbolic, CAT(0) and length spaces. We attempt to determine the extent to which such properties of  $X$  are inherited by  $\mathcal{H}(X)$  or other components of  $\text{CL}(X)$ . Sections 3 and 4 are focused on mappings between hyperspaces. The following class of mappings has been of much interest recently [15].

**DEFINITION 1.1.** Let  $X$  and  $Y$  be metric spaces. An injective mapping  $f : X \rightarrow Y$  is called *quasisymmetric*, or  $\eta$ -quasisymmetric, if there is a homeomorphism  $\eta : [0, \infty) \rightarrow [0, \infty)$  such that

$$\frac{d_Y(f(x_1), f(x_3))}{d_Y(f(x_2), f(x_3))} \leq \eta \left( \frac{d_X(x_1, x_3)}{d_X(x_2, x_3)} \right)$$

for any distinct points  $x_1, x_2, x_3 \in X$ .

In general, a quasisymmetric mapping  $f : X \rightarrow Y$  does not lift to a quasisymmetric mapping of  $\mathcal{H}(X)$ . We call  $f$  *hyperquasisymmetric* if it does. Theorem 3.4 provides two characterizations of such maps, which arise naturally in the context of Gromov hyperbolicity and bi-Lipschitz homogeneity. In §4 we consider a notion of dimension of metric spaces which is invariant under hyperquasisymmetric mappings.

## 2. Geodesics in hyperspaces

Throughout the paper all rectifiable curves are assumed to be parametrized proportional to the arclength, unless stated otherwise. A metric space  $(X, d)$  is called a *length space* if for any  $x, y \in X$  and any  $\epsilon > 0$  there is a curve  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$ ,  $\gamma(1) = y$  and the length of  $\gamma$  is at most  $d(x, y) + \epsilon$ . If such  $\gamma$  exists even for  $\epsilon = 0$ , then  $X$  is called a *geodesic space*. Every proper length space is geodesic [14, 27].

The hyperspace  $\mathcal{H}(X)$  of a geodesic space  $X$  need not be geodesic itself, as is demonstrated by the following example from [10]. Let  $X$  be the Banach space  $c_0$ , namely, the space of all real-valued sequences converging to 0. Let

$$A = \{x \in c_0 : x_n = 1 + 1/n \text{ for an odd number of } n\text{'s, and } x_n = 0 \text{ otherwise}\};$$

$$B = \{x \in c_0 : x_n = 1 + 1/n \text{ for an even number of } n\text{'s, and } x_n = 0 \text{ otherwise}\}.$$

One can see that  $A, B \in \mathcal{H}(X)$  and  $D(A, B) = 1$ , but  $N_{1/2}(A) \cap N_{1/2}(B) = \emptyset$ , hence there is no geodesic connecting  $A$  to  $B$ . If  $X$  is assumed to be both geodesic and compact, then  $\mathcal{H}(X)$  is geodesic as well [10], however, the assumption of compactness is somewhat restrictive.

**THEOREM 2.1.** *Let  $X = (X, d)$  be a length space. Then  $\mathcal{H}(X; C)$  is a length space for all  $C \in \text{CL}(X)$ .*

PROOF. Choose a number  $\sigma > 1$ . For each  $x, y \in X$ , let  $\gamma_{xy} : [0, 1] \rightarrow X$  be a curve of length at most  $\sigma d(x, y)$  such that  $\gamma_{xy}(0) = x$  and  $\gamma_{xy}(1) = y$ . By our convention,  $\gamma_{xy}$  is parametrized proportional to its arclength, which implies  $d(\gamma_{xy}(s), \gamma_{xy}(t)) \leq \sigma |s - t| d(x, y)$  for all  $0 \leq s < t \leq 1$ .

Let  $A, B \in \mathcal{H}(X; C)$  and let  $E = \{(a, b) \in A \times B : d(a, b) \leq \sigma D(A, B)\}$ . By the definition of the Hausdorff metric,

$$(2.1) \quad \pi_A(E) = A \quad \text{and} \quad \pi_B(E) = B,$$

where  $\pi_A, \pi_B$  are the coordinate projections on  $A \times B$ .

Define a function  $\Gamma_{AB}$  from  $[0, 1]$  to the power set of  $X$  as follows:

$$\Gamma_{AB}(t) = \overline{\{\gamma_{ab}(t) : (a, b) \in E\}}.$$

Then  $\Gamma_{AB}(t)$  is closed, and contained in the closed  $\sigma^2 D(A, B)$ -neighborhood of  $A$ . Thus  $\Gamma_{AB} : [0, 1] \rightarrow \mathcal{H}(X; C)$ . By (2.1),  $\Gamma_{AB}(0) = A$  and  $\Gamma_{AB}(1) = B$ . We claim that  $\Gamma_{AB}$  is a rectifiable curve of length at most  $\sigma^2 D(A, B)$ , although it need not be parametrized proportional to the arclength. Let  $s$  and  $t$  be given,  $0 \leq s < t \leq 1$ . For each  $(a, b) \in E$

$$d(\gamma_{ab}(s), \gamma_{ab}(t)) \leq \sigma |s - t| d(a, b) \leq \sigma^2 |s - t| D(A, B),$$

which implies

$$(2.2) \quad D(\Gamma_{AB}(s), \Gamma_{AB}(t)) \leq \sigma^2 |s - t| D(A, B).$$

Thus the length of  $\Gamma_{AB}$  does not exceed  $\sigma^2 D(A, B)$ . Since  $\sigma > 1$  is arbitrary,  $\mathcal{H}(X; C)$  is a length space.  $\square$

COROLLARY 2.2. *Let  $X$  be a proper geodesic space. Then  $\mathcal{H}(X; C)$  is a geodesic space for all  $C \in \text{CL}(X)$ .*

PROOF. Since  $X$  is proper, for any  $a \in X$  and any  $B \in \text{CL}(X)$  there exists  $b \in B$  such that  $d(a, b) = \text{dist}(a, B)$ . This and the fact that  $X$  is a geodesic space allow us to carry out the proof of Theorem 2.1 with  $\sigma = 1$ . Then (2.2) takes form

$$D(\Gamma_{AB}(s), \Gamma_{AB}(t)) \leq |s - t| D(A, B).$$

Since  $\Gamma_{AB}(0) = A$  and  $\Gamma_{AB}(1) = B$ ,

$$\begin{aligned} D(A, B) &\leq D(\Gamma_{AB}(0), \Gamma_{AB}(s)) + D(\Gamma_{AB}(s), \Gamma_{AB}(t)) + D(\Gamma_{AB}(t), \Gamma_{AB}(1)) \\ &\leq (s + |s - t| + 1 - t) D(A, B) \\ &= D(A, B). \end{aligned}$$

Thus equality holds throughout, and we conclude that

$$D(\Gamma_{AB}(s), \Gamma_{AB}(t)) = |s - t| D(A, B)$$

for all  $A, B \in \mathcal{H}(X; C)$  and all  $0 \leq s < t \leq 1$ .  $\square$

REMARK 2.3. In general, geodesics in  $\mathcal{H}(X; C)$  are not unique, even if they are unique in  $X$ . However, if  $\Gamma : [0, 1] \rightarrow \mathcal{H}(X; C)$  is a geodesic between  $A$  and  $B$ , then for every  $t \in [0, 1]$  the inequalities  $D(\Gamma(t), A) \leq tD$  and  $D(\Gamma(t), B) \leq (1 - t)D$  imply

$$\Gamma(t) \subset N_{tD}(A) \cap N_{(1-t)D}(B),$$

where  $D = D(A, B)$ . If the sets  $N_{tD}(A) \cap N_{(1-t)D}(B)$  are not too large, this observation can be used to prove the uniqueness of a geodesic between  $A$  and  $B$ .

It is interesting to determine which properties of metric spaces are inherited by their hyperspaces. The following proposition provides a negative result of this kind. We refer to [9] for definitions of Gromov hyperbolic spaces and CAT(0) spaces.

**PROPOSITION 2.4.** *Let  $X$  be a proper geodesic space. The hyperspace  $\mathcal{H}(X; C)$  is Gromov hyperbolic if and only if  $X$  is bounded.*

**PROOF.** If  $X$  is bounded, then so is  $\mathcal{H}(X; C)$ , and bounded spaces are trivially Gromov hyperbolic.

Suppose that  $X$  is unbounded. Pick two points  $a, b \in X$  at distance  $L$  from each other. Let  $C' = C \setminus B(a, 3L)$  and let  $\gamma_{ab} : [0, 1] \rightarrow X$  be a geodesic connecting  $a$  to  $b$ . The Hausdorff distance between any two of the sets  $C' \cup \{a\}$ ,  $C' \cup \{b\}$  and  $G = C' \cup \gamma_{ab}([0, 1])$  is equal to  $L$ . For  $0 \leq t \leq 1$  let  $\Gamma_{\{a\}\{b\}}(t) = C' \cup \{\gamma_{ab}(tL)\}$ ,  $\Gamma_{\{a\}G}(t) = C' \cup \gamma_{ab}([0, tL])$ , and  $\Gamma_{G\{b\}}(t) = C' \cup \gamma_{ab}([tL, 1])$ . These are geodesics in  $\mathcal{H}(X; C)$ . For every  $t$  the sets  $C' \cup \Gamma_{\{a\}G}(t)$  and  $C' \cup \Gamma_{G\{b\}}(t)$  contain at least one of the points  $a$  and  $b$ . It follows that the distance in  $\mathcal{H}(X; C)$  from  $C' \cup \Gamma_{\{a\}\{b\}}(1/2)$  to  $C' \cup \Gamma_{\{a\}G}([0, 1]) \cup \Gamma_{G\{b\}}([0, 1])$  is equal to  $L/2$ . Since  $L$  can be arbitrarily large,  $\mathcal{H}(X; C)$  is not Gromov hyperbolic.  $\square$

Despite Proposition 2.4, one can find large (unbounded) subsets of  $\mathcal{H}(X)$  that are Gromov hyperbolic, provided that  $X$  is CAT(0) and Gromov hyperbolic. To this end, we introduce *uniformly bounded hyperspaces*

$$\mathcal{H}_L(X) = \{A \in \mathcal{H}(X) : \text{diam } A \leq L\}.$$

Unfortunately, the analogue of Corollary 2.2 is false for  $\mathcal{H}_L(X)$ . Indeed, let  $S(R) = \{x \in \mathbb{R}^n : |x| = R\}$  be a sphere equipped with the intrinsic metric. Choose  $h \in (0, R)$  and let  $A = \{x \in S(R) : x_n = h\}$ ,  $B = \{x \in S(R) : x_n = -h\}$ . Clearly  $D(A, B) = 2R \sin^{-1}(h/R)$ . Using Remark 2.3 we find that there is only one geodesic connecting  $A$  to  $B$  in  $\mathcal{H}(X)$ , namely

$$\Gamma_{AB}(t) = \{x \in S(R) : x_n = R \sin(\sin^{-1}(h/R)(1 - 2t))\}.$$

Since  $\text{diam } \Gamma_{AB}(1/2) = \pi R$ , it follows that  $\mathcal{H}_L(S(R))$  is not geodesic for any  $L < \pi R$ .

**PROPOSITION 2.5.** *If  $X$  is a proper CAT(0) space, then  $\mathcal{H}_L(X)$  is a geodesic space for all  $L \geq 0$ .*

**PROOF.** Given  $A, B \in \mathcal{H}_L(X)$ , let  $\Gamma_{AB}$  be as in the proof of Theorem 2.1 (with  $\sigma = 1$ ). Let  $\gamma = \gamma_{ab}$  and  $\gamma' = \gamma_{a'b'}$ , where  $a, a' \in A$  and  $b, b' \in B$ . Since in CAT(0) spaces the distance function is convex [9, p. 176], we have

$$d(\gamma(t), \gamma'(t)) \leq (1 - t)d(\gamma(0), \gamma'(0)) + td(\gamma(1), \gamma'(1)) \leq (1 - t)L + tL = L.$$

Thus  $\text{diam } \Gamma_{AB}(t) \leq L$  as required.  $\square$

**REMARK 2.6.** Under the assumptions of Proposition 2.5 the space  $\mathcal{H}_L(X)$  does not have to be a CAT(0) space itself. Indeed, let  $X = \mathbb{R}$ ,  $A = [0, 1]$  and  $B = [1, 2]$ . The paths  $\Gamma(t) = [t, 1 + t]$  and

$$\Gamma'(t) = \begin{cases} [t, 1 + t] \setminus (1/2, 1/2 + 2t), & 0 \leq t \leq 1/2; \\ [t, 1 + t] \setminus (2t - 1/2, 3/2), & 1/2 \leq t \leq 1 \end{cases}$$

are both geodesics from  $A$  to  $B$  in  $\mathcal{H}_1(\mathbb{R})$ . Nonuniqueness of geodesics immediately implies that  $\mathcal{H}_1(\mathbb{R})$  is not a CAT(0) space [9, p. 160].

**THEOREM 2.7.** *Let  $X$  be a proper CAT(0) space. If  $X$  is  $\delta$ -hyperbolic, then  $\mathcal{H}_L(X)$  is  $\delta'$ -hyperbolic, where  $\delta'$  depends on  $\delta$  and  $L$ . Furthermore, the Gromov boundaries of  $X$  and  $\mathcal{H}_L(X)$  are quasimetrically equivalent.*

**PROOF.** Let  $s : \mathcal{H}_L(X) \rightarrow X$  be a mapping such that  $s(A) \in A$  for every  $A \in \mathcal{H}_L(X)$  (the existence of such  $s$  follows from the Axiom of Choice). Let  $A$  and  $B$  be two sets in  $\mathcal{H}_L(X)$ , and let  $D = D(A, B)$ ,  $d = d(s(A), s(B))$ . Since  $s(A) \in N_D(B) \subset N_{D+L}(\{s(B)\})$ , we have  $d \leq D + L$ . Conversely,  $A \subset N_L(\{s(A)\}) \subset N_{d+L}(B)$  and  $B \subset N_{d+L}(A)$ , which means that  $d \geq D - L$ . Thus the mapping  $s$  is a  $(1, L)$ -quasi-isometry. Since  $\mathcal{H}_L(X)$  admits a quasi-isometric embedding into a  $\delta$ -hyperbolic space, it is  $\delta'$ -hyperbolic with  $\delta' = \delta'(L, \delta)$  [9, p. 402]. By [9, p. 436] the mapping  $s$  induces a quasimetric mapping  $\partial s : \partial \mathcal{H}_L(X) \rightarrow \partial X$ . Since  $s$  is surjective, so is  $\partial s$ .  $\square$

We now turn to the hyperspace of convex sets  $\mathcal{H}^c(X)$  and its subspace  $\mathcal{H}_L^c(X) = \mathcal{H}^c(X) \cap \mathcal{H}_L(X)$ . Corollary 2.2 fails for  $\mathcal{H}^c(X)$ , which can be seen as follows. As before, let  $S(R) = \{x \in \mathbb{R}^n : |x| = R\}$  be a sphere equipped with the intrinsic metric. Choose  $h \in (0, R)$  and let  $A = \{x \in S(R) : x_n \geq h\}$ ,  $B = \{x \in S(R) : x_n = -R\}$ . Clearly  $A, B \in \mathcal{H}^c(S(R))$  and  $D(A, B) = \pi R$ . Suppose that  $\gamma : [0, 1] \rightarrow \mathcal{H}^c(S(R))$  is a geodesic segment joining  $A$  to  $B$ . Choose three points  $a, b, c \in A$  so that the geodesic triangle  $\Delta(a, b, c)$  in  $S(R)$  with vertices  $a, b$  and  $c$  contains the point  $p = (0, 0, \dots, R)$  in its interior. For all sufficiently small  $t > 0$  the set  $\gamma(t)$  must contain points  $a', b'$  and  $c'$  such that the interior of the geodesic triangle  $\Delta(a', b', c')$  also contains  $p$ . Since  $A_t$  is geodesically convex, we have  $P \in A_t$ , hence  $D(A_t, B) = \pi R = D(A, B)$ . This is a contradiction.

Once again, the condition CAT(0) saves the day. When  $X$  is a CAT(0) space, the geodesics in  $X$  are unique, which allows us to define the convexification map  $\text{conv} : \mathcal{H}(X) \rightarrow \mathcal{H}^c(X)$  as follows:  $\text{conv}(A)$  is the intersection of all geodesically convex closed sets containing  $A$ . To prove that the hyperspaces of convex subsets of CAT(0) spaces are geodesic, we require the following lemma.

**LEMMA 2.8.** *If  $X$  is a proper CAT(0) space, then the convexification map  $\text{conv} : \mathcal{H}(X) \rightarrow \mathcal{H}^c(X)$  is a contraction.*

**PROOF.** Given  $A \in \mathcal{H}(X)$ , one can write  $\text{conv}(A)$  as the closure of an increasing union of sets  $A_m$ , where  $A_0 = A$  and  $A_{m+1}$  is the union of all geodesics joining points of  $A_m$ . To prove that  $\text{conv}$  is a contraction, it suffices to show  $D(\overline{A_{m+1}}, \overline{B_{m+1}}) \leq D(\overline{A_m}, \overline{B_m})$  for all  $m$ , or equivalently, just for  $m = 0$ . For every  $a \in A_1$  there exist a geodesic segment  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0), \gamma(1) \in A$  and  $\gamma(t) = a$  for some  $t$ . Choose another geodesic  $\gamma' : [0, 1] \rightarrow X$  so that  $\gamma'(0), \gamma'(1) \in B$ ,  $d(\gamma(0), \gamma'(0)) \leq D(A, B)$ , and  $d(\gamma(1), \gamma'(1)) \leq D(A, B)$ . Using the convexity of the distance function in CAT(0) spaces [9, p. 176], we obtain

$$d(a, \gamma'(t)) \leq td(\gamma(0), \gamma'(0)) + (1-t)d(\gamma(1), \gamma'(1)) \leq D(A, B).$$

Since  $\gamma'(t) \in B_1$ , it follows that  $\overline{A_1} \subset N_{D(A, B)}(\overline{B_1})$ . Interchanging the roles of  $A$  and  $B$ , we obtain  $D(\overline{A_1}, \overline{B_1}) \leq D(A, B)$ , as required.  $\square$

**COROLLARY 2.9.** *If  $X$  is a proper CAT(0) space, then  $\mathcal{H}^c(X)$  and  $\mathcal{H}_L^c(X)$  are geodesic spaces for all  $L \geq 0$ .*

**PROOF.** By virtue of Corollary 2.2, Proposition 2.5 and Lemma 2.8 the proof reduces to the following simple observation. If  $\gamma$  is a geodesic in a metric space

$Y$ , and  $T : Y \rightarrow Y$  is a contraction that fixes the endpoints of  $\gamma$ , then  $T \circ \gamma$  is a geodesic in  $T(Y)$ .  $\square$

### 3. Mappings of hyperspaces

Every continuous mapping  $f : X \rightarrow Y$  induces a mapping  $G_f : \text{CL}(Y) \rightarrow \text{CL}(X)$  as follows:  $G_f(A) = f^{-1}(A)$ . In general  $G_f$  is not continuous, and can even map two sets at finite distance from each other into sets at infinite distance. Indeed, let  $X = \{(x, y) \in \mathbb{R}^2 : x > 0\} \cup \{(x, 0) \in \mathbb{R}^2 : x \leq 0\}$ ,  $Y = \mathbb{R}$ , and define  $f : X \rightarrow Y$  by  $f(x, y) = x$ . Obviously  $D(G_f(\{0\}), G_f(\{1\})) = \infty$ . Observe that  $f$  is an open mapping, i.e. for any  $x \in X$  and  $r > 0$  there is  $\rho > 0$  such that  $B(f(x), \rho) \subset f(B(x, r))$ . However,  $\rho$  cannot be chosen independently of  $r$ . This leads us to the following definition, which appeared, e.g. in [19].

**DEFINITION 3.1.** A mapping  $f : X \rightarrow Y$  between two metric spaces  $X, Y$  is called *co-uniformly continuous* if there exists an increasing function  $\tilde{\omega}_f : (0, \infty) \rightarrow (0, \infty)$  such that  $B(f(x), \tilde{\omega}_f(r)) \subset f(B(x, r))$  for all  $x \in X$  and all  $r > 0$ .

If  $f : X \rightarrow Y$  is co-uniformly continuous, then  $G_f$  is uniformly continuous. Indeed, given two sets  $A, B \in \text{CL}(Y)$  such that  $D(A, B) \leq \tilde{\omega}_f(r)$ , we easily obtain  $f^{-1}(A) \subset N_r(f^{-1}(B))$  and  $f^{-1}(B) \subset N_r(f^{-1}(A))$ . Similarly, if  $f$  is a Lipschitz quotient (defined in the introduction), then  $G_f$  is bi-Lipschitz (cf. Lemma 6.1 [24]).

Every uniformly continuous homeomorphism  $f : X \rightarrow Y$  between metric spaces lift to a continuous mapping  $F_f : \mathcal{H}(X) \rightarrow \mathcal{H}(Y)$  of the corresponding hyperspaces:

$$F_f(A) = \{f(a) : a \in A\}.$$

If  $f$  is bi-Lipschitz, then  $F_f$  is also bi-Lipschitz. In other words, the class of bi-Lipschitz mappings is invariant under the hyperspace functor. The same is true for isometries, and moreover, for many spaces  $X$  every isometry of  $\mathcal{H}(X)$  coincides with  $F_f$  where  $f$  is an isometry of  $X$  [3, 12]. The following result shows that quasisymmetric mappings are not preserved by the hyperspace functor. Before stating it, let us introduce the pointwise Lipschitz constant of  $f$  at  $x \in X$  by

$$L_f(x) := \limsup_{y \rightarrow x} \frac{d(f(x), f(y))}{d(x, y)}.$$

We say that a map  $f : X \rightarrow Y$  is *hyperquasisymmetric* if  $F_f$  is quasisymmetric. Considering the action of  $F_f$  on singletons, we see that such  $f$  must be quasisymmetric itself.

**PROPOSITION 3.2.** *Let  $f : X \rightarrow Y$  be a homeomorphism of connected metric spaces which is Lipschitz on a neighborhood of one point in  $X$ , and whose pointwise Lipschitz constant is infinite at another point in  $X$ . Then  $f$  is not hyperquasisymmetric.*

For example, the map  $f : [0, 1] \rightarrow [0, 1]$ ,  $f(t) = \sqrt{t}$ , is quasisymmetric, but not hyperquasisymmetric.

**PROOF.** We use primes to denote images under  $f$  and  $F_f$ , i.e.,  $u' = f(u)$ ,  $A' = F_f(A)$ , etc. Let  $U$  be a neighborhood of a point  $p \in X$  such that  $f|_U$  is  $L$ -Lipschitz, and let  $q \in X \setminus U$  satisfy  $L_f(q) = +\infty$ . We may assume without loss of generality that  $U = B(p, \epsilon)$  for some

$$0 < \epsilon < \min \left\{ \frac{1}{3}d(p, q), \frac{1}{2L+1}d(p', q') \right\}.$$

Choose points  $v_n \rightarrow q$  so that  $d(v'_n, q') > nd(v_n, q)$ . By restricting our attention to sufficiently large  $n$ , we may assume that

$$\max\{d(v_n, q), d(v'_n, q')\} \leq \epsilon.$$

Then

$$\min\{d(v_n, p), d(v'_n, p')\} \geq \epsilon$$

for all  $n$ .

Since  $X$  is connected, we may choose points  $u_n \in U$  so that  $\delta_n := d(p, u_n) = d(q, v_n)$  for all  $n$ . Let  $A_n = \{p, v_n\}$ ,  $B_n = \{p, u_n, v_n\}$ , and  $C_n = \{p, v_n, q\}$ . Then  $D(A_n, B_n) = D(A_n, C_n) = \delta_n$ . On the other hand,

$$D(A'_n, B'_n) = \min\{d(p', u'_n), d(u'_n, v'_n)\} \leq L\delta_n$$

while

$$D(A'_n, C'_n) = \min\{d(p', q'), d(v'_n, q')\} > n\delta_n.$$

Thus  $F_f$  is not quasimetric.  $\square$

The main theorem of this section provides two characterizations of the class of hyperquasimetric maps and incidentally shows that this class is invariant under the hyperspace functor. In other words, hyperquasimetric maps lift to quasimetric maps on iterated hyperspaces such as  $\mathcal{H}(\mathcal{H}(X))$ . See [2, 29, 32] for more on iterated hyperspaces. Some preliminaries are required to state the theorem. Given a metric space  $X$ , let  $\text{Dist}(X)$  be its distance set:

$$\text{Dist}(X) = \{d_X(x_1, x_2) : x_1, x_2 \in X\} \subset [0, \infty).$$

The modulus of continuity of a mapping  $f : X \rightarrow Y$  is defined as

$$\omega_f(\delta) = \sup\{d_Y(f(x_1), f(x_2)) : d_X(x_1, x_2) \leq \delta\}, \quad \delta > 0.$$

We say that  $\omega_f$  is controlled by a homeomorphism  $\eta : [0, \infty) \rightarrow [0, \infty)$  provided that  $\omega_f(\delta) < \infty$  for all  $\delta > 0$  and

$$\omega_f(t\delta) \leq \eta(t)\omega_f(\delta), \quad \delta \in \text{Dist}(X), \quad t > 0.$$

Whether or not  $\omega_f$  is controlled by  $\eta$  may depend on  $X$  as well as on  $\omega_f$  and  $\eta$ .

**LEMMA 3.3.** *Let  $C \in \text{CL}(X)$ . For any  $\delta \in \text{Dist}(\mathcal{H}(X; C))$  there exists a non-decreasing sequence  $\{\delta_n\} \subset \text{Dist}(X)$  such that  $\delta_n \rightarrow \delta$ .*

**PROOF.** Choose  $A, B \in \mathcal{H}(X; C)$  so that  $D(A, B) = \delta$ . Without loss of generality we may assume that for any  $\epsilon > 0$  there is  $a \in A$  such that  $\text{dist}(a, B) \geq \delta - \epsilon$ . Then there is  $b \in B$  such that  $\delta - \epsilon \leq d_X(a, b) \leq \delta$ . Since  $\epsilon > 0$  is arbitrary, the claim follows.  $\square$

**THEOREM 3.4.** *Let  $f$  be a homeomorphism of a metric space  $X$  into a metric space  $Y$ . The following are equivalent:*

- (i)  $f$  is hyperquasimetric;
- (ii) there exists a homeomorphism  $\eta : [0, \infty) \rightarrow [0, \infty)$  such that for any four distinct points  $x_1, \dots, x_4 \in X$

$$(3.1) \quad \frac{d_Y(f(x_1), f(x_2))}{d_Y(f(x_3), f(x_4))} \leq \eta \left( \frac{d_X(x_1, x_2)}{d_X(x_3, x_4)} \right);$$

(iii)  $\omega_f$  is controlled by some homeomorphism  $\eta : [0, \infty) \rightarrow [0, \infty)$  and there exists  $c > 0$  such that for all  $x_1, x_2 \in X$

$$(3.2) \quad d_Y(f(x_1), f(x_2)) \geq c\omega_f(d_X(x_1, x_2));$$

(iv)  $F_f$  is hyperquasisymmetric.

REMARK 3.5. The mappings satisfying (3.1) have previously appeared in the literature on several occasions. Ivascu [18] called them “freely quasisymmetric”; we do not use this term to avoid confusion with Väisälä’s well-known terminology [33]. Aseev and Shalaginov [1] studied such mappings in the context of self-similar spaces. They proved, among other things, that a mapping  $f : X \rightarrow Y$  satisfies (3.1) if and only if it lifts to a quasisymmetric mapping  $f \times f : X \times X \rightarrow Y \times Y$ . See also [28]. Ghamsari and Herron [13] and Herron and Mayer [16] studied this class of maps in connection with the question of characterizing bi-Lipschitz homogeneous Jordan curves. Combining Theorem 3.4 with Theorem E of [16] leads to the following result: a Jordan curve  $\Gamma$  in a doubling metric space is bi-Lipschitz homogeneous and bounded turning if and only if it admits a hyperquasisymmetric parametrization.

PROOF OF THEOREM 3.4. (i) $\Rightarrow$ (ii). Suppose that  $F_f$  is  $\eta$ -quasisymmetric. Choose distinct points  $x_1, \dots, x_4 \in X$  and set  $A = \{x_1, x_3\}$ ,  $B = \{x_2, x_3\}$  and  $C = \{x_1, x_4\}$ . Then

$$D(A, C) = \min\{d_X(x_3, x_4), \max\{d_X(x_1, x_3), d_X(x_1, x_4)\}\} \geq \frac{1}{2}d_X(x_3, x_4)$$

and

$$D(A, B) = \min\{d_X(x_1, x_2), \max\{d_X(x_1, x_3), d_X(x_2, x_3)\}\} \leq d_X(x_1, x_2).$$

Similar considerations show that

$$D(F_f(A), F_f(B)) \geq \frac{1}{2}d_Y(f(x_1), f(x_2))$$

and

$$D(F_f(A), F_f(C)) \leq d_Y(f(x_3), f(x_4)).$$

Using the quasisymmetry of  $F_f$ , we find

$$\frac{d_Y(f(x_1), f(x_2))}{d_Y(f(x_3), f(x_4))} \leq 2\eta \left( 2 \frac{d_X(x_1, x_2)}{d_X(x_3, x_4)} \right).$$

Thus  $f$  satisfies (ii) with the function  $\eta_1(t) = 2\eta(2t)$ .

(ii) $\Rightarrow$ (iii). The first part of (iii) follows immediately from (ii). To prove (3.2), let  $\delta = d_X(x_1, x_2)$ , pick  $\epsilon > 0$  and choose  $x_3, x_4 \in X$  so that  $d_X(x_3, x_4) \leq \delta$  and  $d_Y(f(x_3), f(x_4)) \geq \omega_f(\delta) - \epsilon$ . Using (3.1) with  $x_i$  appropriately rearranged, we obtain

$$d_Y(f(x_1), f(x_2)) \geq \eta(1)^{-1}(\omega_f(\delta) - \epsilon),$$

and (iii) follows with  $c = 1/\eta(1)$ .

(iii) $\Rightarrow$ (i). First we prove that  $F_f$  itself has the properties listed in (iii). It is easy to see that  $F_f$  has the same modulus of continuity as  $f$ . Given  $\delta \in \text{Dist}(\mathcal{H}(X))$ , let  $\{\delta_n\}$  be as in Lemma 3.3. For any  $t > 0$  we have

$$\omega_f(t\delta) \leq \eta(t\delta_n/\delta)\omega_f(\delta_n) \leq \eta(t\delta_n/\delta)\omega_f(\delta).$$

Letting  $n \rightarrow \infty$  yields  $\omega_f(t\delta) \leq \eta(t)\omega_f(\delta)$ . Thus the modulus of continuity of  $F_f$  is controlled by  $\eta$ .

To prove that  $F_f$  satisfies (3.2), pick  $A, B \in \mathcal{H}(X)$  and let  $s = D(A, B)$ . We may assume that there exists a sequence  $\{a_n\} \subset A$  such that  $s_n := \text{dist}_X(a_n, B) \rightarrow s$ ; otherwise interchange  $A$  and  $B$ . Passing to a subsequence, we can ensure that one of the following two cases occurs.

*Case 1.* For each  $n$  there is  $b_n \in B$  such that  $d_X(a_n, b_n) = s_n$ . Inequality (3.2) implies

$$(3.3) \quad \text{dist}_Y(f(a_n), f(B)) \geq c\omega_f(s_n).$$

Since  $s_n \in \text{Dist}(X)$ , it follows that

$$(3.4) \quad \lim_{n \rightarrow \infty} \omega_f(s_n) \geq \lim_{n \rightarrow \infty} \eta(s/s_n)^{-1} \omega_f(s) = \eta(1)^{-1} \omega_f(s).$$

Combining (3.3) and (3.4) we obtain  $D(f(A), f(B)) \geq c\eta(1)^{-1} \omega_f(s)$ . Thus  $F_f$  satisfies (3.2) with the constant  $c' = c/\eta(1)$ .

*Case 2.* the distance from  $a_n$  to  $B$  is not achieved for any  $n$ . Inequality (3.2) implies

$$(3.5) \quad \text{dist}_Y(f(a_n), f(B)) \geq c \lim_{\epsilon \downarrow 0} \omega_f(s_n + \epsilon).$$

Choose a sequence  $\{b_{mn}\} \subset B$  such that  $d_X(a_n, b_{mn}) \downarrow s_n$  as  $m \rightarrow \infty$ . Since  $d_X(a_n, b_{mn}) \in \text{Dist}(X)$ , we have

$$\omega_f(s) \leq \eta(s/s_n) \lim_{\epsilon \downarrow 0} \omega_f(s_n + \epsilon)$$

The latter inequality and (3.5) yield  $D(f(A), f(B)) \geq c\eta(1)^{-1} \omega_f(s)$ . This completes the proof of property (3.2) for  $F_f$ .

Finally, for any distinct  $A, B, C \in \mathcal{H}(X)$  we have

$$\frac{D(f(A), f(B))}{D(f(A), f(C))} \leq \frac{\omega_f(D(A, B))}{c' \omega_f(D(A, C))} \leq \frac{1}{c'} \eta \left( \frac{D(A, B)}{D(A, C)} \right),$$

which proves (i).

(i) $\Rightarrow$ (iv). If  $f$  is hyperquasisymmetric, then it verifies (iii), and so does  $F_f$  (by the previous step). Since (iii) implies (i),  $F_f$  is hyperquasisymmetric.  $\square$

**REMARK 3.6.** The proof of implication (iii) $\Rightarrow$ (i) works for unbounded closed sets as well. Therefore, a hyperquasisymmetric mapping  $f : X \rightarrow Y$  induces (hyper-)quasisymmetric mappings on all components of  $\text{CL}(X)$ .

**COROLLARY 3.7.** *Let  $X$  be a Gromov hyperbolic space, and let  $d'$  and  $d''$  be two visual metrics on its boundary  $\partial X$ . Then the identity map  $\text{id} : (\partial X, d') \rightarrow (\partial X, d'')$  is hyperquasisymmetric.*

**PROOF.** There exist  $\alpha > 0$  and  $C > 1$  such that

$$C^{-1}(d'(p, q))^\alpha \leq d''(p, q) \leq C(d'(p, q))^\alpha$$

for all  $p, q \in \partial X$  (see, e.g. Proposition III.H.3.21 in [9]). This and Theorem 3.4 imply that  $\text{id}$  is hyperquasisymmetric.  $\square$

Corollary 3.7 can be generalized as follows. Bonk and Schramm [7] call  $f : X \rightarrow Y$  a *snowflake map* if there exist  $\alpha > 0$  and  $C \geq 1$  such that

$$C^{-1}d_X(x_1, x_2)^\alpha \leq d_Y(f(x_1), f(x_2)) \leq Cd_X(x_1, x_2)^\alpha$$

for all  $x_1, x_2 \in X$ . By Theorem 3.4 snowflake maps are hyperquasisymmetric. Examples of snowflake maps are provided by Theorem 6.5 [7], which asserts that

rough similarities between Gromov hyperbolic spaces induce snowflake maps on their boundaries.

#### 4. Hyperconformal dimension

The following definition is motivated by Corollary 3.7.

DEFINITION 4.1. Let  $X$  be a metric space. The *hyperconformal* dimension of  $X$  is defined by

$$\text{HCdim } X = \inf\{\text{Hdim } f(X) : f \text{ is hyperquasisymmetric}\},$$

where  $\text{Hdim}$  stands for the Hausdorff dimension [23].

Definition 4.1 is modeled after the more commonly used definition of *conformal dimension* [6, 15, 25]:

$$\text{Cdim } X = \inf\{\text{Hdim } f(X) : f \text{ is quasisymmetric}\}.$$

Evidently  $\text{Cdim } X \leq \text{HCdim } X \leq \text{Hdim } X$ . By virtue of Corollary 3.7 both conformal and hyperconformal dimensions of the boundary of a hyperbolic space are well-defined. Moreover, conformal dimension of  $\partial X$  is invariant under quasi-isometries of  $X$ , while hyperconformal dimension is invariant under rough similarities of  $X$  [7].

Since hyperquasisymmetric maps of metric spaces preserve more structure than general quasisymmetric maps, one can expect  $\text{HCdim } X$  to reveal some features of  $X$  that are not recorded by  $\text{Cdim } X$ . For instance, conformal dimension does not distinguish between spaces  $X$  with  $\text{Cdim } X < 1$  [21, 31]. This is in contrast with the following result.

THEOREM 4.2. *For every  $s \geq 0$  there exists a compact metric space  $X$  with  $\text{HCdim } X = s$ .*

PROOF. For each  $s \in \{0\} \cup [1, \infty)$  there exists a compact metric space  $X$  such that  $\text{Cdim } X = \text{Hdim } X = s$  and therefore  $\text{HCdim } X = s$  [5, 8, 25, 30]. It remains to consider the case  $0 < s < 1$ . For any such  $s$  we shall construct  $X \subset \mathbb{R}$  as a union of two generalized Cantor sets [23, 4.11]. Given an infinite set of positive integers  $E \subset \{1, 2, \dots\}$ , define a sequence  $\lambda^E = \{\lambda_n^E : n \geq 1\}$  so that  $\lambda_n^E = 1/2$  for  $n \notin E$  and  $\prod_{k=1}^n \lambda_k^E = 2^{-n/s}$  for  $n \in E$ . Let

$$C(\lambda^E) = \bigcap_{n=0}^{\infty} \bigcup_{j=1}^{2^n} I_j^n,$$

where  $I_1^0 = [0, 1]$  and the other closed intervals  $I_j^n$  are defined as follows. The intervals  $\{I_j^n\}_{j=1}^{2^n}$  are disjoint, each of length equal to  $\mu_n^E := \prod_{k=1}^n \lambda_k^E$ ; furthermore, each  $I_j^n$  is contained in an interval  $I_i^{n-1}$  and shares an endpoint with it.

By virtue of [23, 4.11] we have  $\text{Hdim } C(\lambda^E) = s$  for any  $E$  as above. Let  $n_i$  be a strictly increasing sequence of integers such that

$$(4.1) \quad n_{i+2} - n_i > \frac{n_{i+1} - n_i}{s} + 1, \quad i \geq 1,$$

and

$$(4.2) \quad \lim_{i \rightarrow \infty} \frac{n_i}{i} = \infty.$$

Define  $E = \{n_{2i-1} : i \geq 1\}$ ,  $F = \{n_{2i} : i \geq 1\}$ , and  $X = C(\lambda^E) \cup (1 + C(\lambda^F)) \subset \mathbb{R}$ . Clearly  $\text{Hdim } X = s$ . We claim that  $\text{HCdim } X = s$ , or, equivalently,  $\text{Hdim } f(X) \geq s$  for any hyperquasisymmetric mapping  $f$ .

A generalized Cantor set such as  $C(\lambda^E)$  contains the endpoints of all intervals  $I_j^n$  used in its construction. Therefore, for any  $i \geq 1$  the set  $X$  contains a pair of points  $a_i, b_i$  such that  $b_i - a_i = 2^{-n_i/s}$ . Furthermore,  $X$  contains the points that split  $[a_i, b_i]$  into  $2^{n_{i+2} - n_i - 1}$  equal subintervals. Suppose that  $f : X \rightarrow Y$  is hyperquasisymmetric, and  $c > 0$  is as in (3.2). For every integer  $j$  between 1 and  $n_{i+2} - n_i - 1$  we have

$$(4.3) \quad \omega_f(2^{-j-n_i/s}) \geq 2^{-j} d_Y(f(a_i), f(b_i)) \geq 2^{-j} c \omega_f(2^{-n_i/s}).$$

Setting  $j$  equal to the smallest integer greater than  $(n_{i+1} - n_i)/s$  (cf. (4.1)), we obtain

$$\omega_f(2^{-n_{i+1}/s}) \geq 2^{-(n_{i+1} - n_i)/s - 1} c \omega_f(2^{-n_i/s}).$$

Therefore, for all  $i > 1$

$$(4.4) \quad \omega_f(2^{-n_i/s}) \geq \left(\frac{c}{2}\right)^{i-1} 2^{-(n_i - n_1)/s} \omega_f(2^{-n_1/s}) =: C_1 \left(\frac{c}{2}\right)^{i-1} 2^{-n_i/s}.$$

For any  $\epsilon > 0$  inequality (4.4) implies

$$\lim_{i \rightarrow \infty} 2^{(1+\epsilon)n_i/s} \omega_f(2^{-n_i/s}) \geq C_1 \lim_{i \rightarrow \infty} 2^{\epsilon n_i/s} \left(\frac{c}{2}\right)^{i-1} = \infty,$$

where in the last step we used (4.2). This together with (4.3) yield

$$\lim_{\delta \downarrow 0} \omega_f(\delta) \delta^{-1-\epsilon} = \infty.$$

Using (3.2) again, we conclude that

$$\text{Hdim } f(X) \geq (1 + \epsilon)^{-1} \text{Hdim } X.$$

Since  $\epsilon$  was arbitrary,  $\text{Hdim } f(X) \geq \text{Hdim } X$  as desired.  $\square$

A metric space  $X$  is called Ahlfors regular if there exist  $s > 0$  and  $C \geq 1$  such that the  $s$ -dimensional Hausdorff measure  $H^s$  satisfies

$$C^{-1} r^s \leq H^s(B(x, r)) \leq C r^s$$

for all  $x \in X$  and  $0 < r \leq 2 \text{diam } X$ . Since the boundary of a Gromov hyperbolic group is Ahlfors regular, it is natural to ask the following

**QUESTION 4.3.** Are there any Ahlfors regular metric spaces with hyperconformal dimension between 0 and 1?

The following proposition shows that  $\text{HCdim}$  is much more rigid than  $\text{Cdim}$ .

**PROPOSITION 4.4.** *If  $X$  contains a nontrivial rectifiable curve, then  $\text{HCdim } X = \text{Hdim } X$ .*

**PROOF.** Let  $\gamma : [0, L] \rightarrow X$  be a curve parametrized by arclength and such that  $\gamma(0) \neq \gamma(1)$ . Suppose that  $f : X \rightarrow Y$  is surjective and hyperquasisymmetric. Our goal is to prove that  $\text{Hdim } Y \geq \text{Hdim } X$ . Given  $n \geq 1$ , let  $x_i = \gamma(Li/n)$ ,  $i = 0, \dots, n$ . Since

$$\sum_{i=1}^n d_Y(f(x_i), f(x_{i-1})) \geq d_Y(f(\gamma(0)), f(\gamma(1))) =: s$$

and  $d_X(x_i, x_{i-1}) \leq L/n$  for all  $i$ , it follows that  $\omega_f(L/n) \geq s/n$ . Using the fact that  $\omega_f$  is nondecreasing, we obtain  $\liminf_{\delta \rightarrow 0} \delta^{-1} \omega_f(\delta) > 0$ . By property (iii) in Theorem 3.4,  $f^{-1}$  is Lipschitz on small scales. Thus  $\text{Hdim } Y \geq \text{Hdim } X$ .  $\square$

Proposition 4.4 implies that the hyperconformal dimension of the Sierpiński carpet  $S$  is equal to  $\text{Hdim } S = \log 8 / \log 3$ . Although the precise value of  $\text{Cdim } S$  remains unknown, it is known to be strictly less than  $\text{Hdim } S$  [6, 20].

Bonk and Kleiner [6] introduced another version of  $\text{Cdim}$ , called *Ahlfors regular conformal dimension*. It is defined only for Ahlfors regular metric spaces  $X$ :

$$\text{ARCdim } X = \inf\{\text{Hdim } f(X) : f \text{ is quasymmetric and } f(X) \text{ is Ahlfors regular}\}.$$

Clearly  $\text{Cdim } X \leq \text{ARCdim } X \leq \text{Hdim } X$ . It is therefore of interest to compare  $\text{ARCdim}$  with  $\text{HCdim}$ . The Sierpiński carpet  $S$  has  $\text{ARCdim } S < \text{HCdim } S$  [20]. It remains unclear if there is an Ahlfors regular space  $X$  with  $\text{ARCdim } X > \text{HCdim } X$ . No such spaces would exist if hyperquasymmetric mappings preserved Ahlfors regularity; however, this is not the case.

PROPOSITION 4.5. *There exists a hyperquasymmetric mapping of the real line  $\mathbb{R}$  onto a metric space  $Y$  that is not Ahlfors regular.*

PROOF. Let  $Y$  be the real line  $\mathbb{R}$  equipped with metric  $d_Y(a, b) = \rho(|a - b|)$ , where  $\rho : [0, \infty) \rightarrow [0, \infty)$  is an increasing concave function to be determined. The identity map  $\text{id} : \mathbb{R} \rightarrow Y$  is  $\eta$ -quasymmetric provided that

$$(4.5) \quad \frac{\rho(tu)}{\rho(u)} \leq \eta(t), \quad t, u > 0.$$

We shall choose  $\rho$  so that it is  $C^1$  smooth on  $(0, \infty)$  and

$$(4.6) \quad C^{-1} \leq \frac{u\rho'(u)}{\rho(u)} \leq C, \quad u > 0$$

for some  $C \geq 1$ . It is easy to see that (4.6) yields (4.5) with  $\eta(t) = \max\{t^C, t^{1/C}\}$ . Theorem 3.4 implies that  $\text{id} : \mathbb{R} \rightarrow Y$  is hyperquasymmetric as long as it is quasymmetric. It remains to construct  $\rho$  so that (4.6) holds but  $Y$  is not Ahlfors regular. One possible choice is

$$\rho(u) = \begin{cases} u \log(3/u), & 0 < u \leq 1; \\ (\log 3 - 1)u + 1, & u > 1. \end{cases}$$

Indeed,

$$\rho'(u) = \begin{cases} \log(3/u) - 1, & 0 < u \leq 1; \\ \log 3 - 1, & u > 1, \end{cases}$$

hence  $\rho$  is concave and

$$1 - \frac{1}{\log 3} \leq \frac{u\rho'(u)}{\rho(u)} \leq 1, \quad u > 0.$$

It remains to observe that  $\text{Hdim } Y = 1$  but the 1-dimensional Hausdorff measure on  $Y$  is not locally finite.  $\square$

## 5. Embeddings into hyperspaces

In this section we consider isometric and bi-Lipschitz embeddings of metric spaces into hyperspaces. Of course, every space  $X$  embeds isometrically into  $\mathcal{H}(X)$  via the map  $x \mapsto \{x\}$ ; our goal is to embed  $X$  into the hyperspace of a simpler metric space, such as  $\mathbb{R}$ . Before stating our first result, we introduce the notation  $l_+^\infty = \{x \in l^\infty : x_k \geq 0 \ \forall k\}$ .

**THEOREM 5.1.** *Let  $X$  be a connected metric space. If  $C \in \text{CL}(X)$  is unbounded, then  $\mathcal{H}(X; C)$  contains an isometric copy of  $l_+^\infty$ . Furthermore, every separable metric space admits a  $\sqrt{2}$ -bi-Lipschitz embedding into  $\mathcal{H}(X; C)$ , and every bounded separable metric space admits an isometric embedding. The constant  $\sqrt{2}$  is sharp.*

**PROOF.** Given  $x = (x_1, x_2, \dots) \in l_+^\infty$ , let  $x_{ij} = \min\{x_i, j\}$  for  $i, j = 1, 2, \dots$ . The sequence  $\tilde{x} := (x_{11}, x_{21}, x_{12}, x_{31}, x_{22}, x_{13}, \dots)$  verifies  $0 \leq \tilde{x}_k \leq k$  for all  $k \geq 1$ . Also, the map  $x \mapsto \tilde{x}$  is an isometry of  $l_+^\infty$  into itself.

Since  $C$  is unbounded, it contains a sequence  $\{c_n\}$  such that

$$d(c_n, c_0) \geq d(c_{n-1}, c_0) + 2n, \quad n \geq 1.$$

For  $x \in l_+^\infty$  let

$$A(x) = \left( C \setminus \bigcup_{n=1}^{\infty} B(c_n, \tilde{x}_n) \right) \cup \bigcup_{n=1}^{\infty} \partial B(c_n, \tilde{x}_n).$$

Since  $\tilde{x} \in l_+^\infty$ , it follows that  $A(x) \in \mathcal{H}(X; C)$ . It is straightforward to check that  $D(A(x), A(y)) = \|\tilde{x} - \tilde{y}\| = \|x - y\|$  for all  $x, y \in l_+^\infty$ .

Every separable metric space embeds isometrically into  $l^\infty$  by Frechét's theorem. The map  $(x_1, x_2, \dots) \mapsto \sqrt{2}(x_1^+, x_1^-, x_2^+, x_2^-, \dots)$  is easily seen to be a  $\sqrt{2}$ -bi-Lipschitz embedding of  $l^\infty$  into  $l_+^\infty$ . Also, a bounded subset of  $l^\infty$  can be isometrically mapped into  $l_+^\infty$  by translation  $x \mapsto x + (M, M, \dots)$ , where  $M$  is sufficiently large.

It remains to show the sharpness of the constant  $\sqrt{2}$ . Suppose that  $F : \mathbb{Z} \rightarrow \mathcal{H}(\mathbb{R}; \mathbb{R})$  is an  $L$ -bi-Lipschitz embedding. Observe that for any sets  $A, B \in \mathcal{H}(\mathbb{R}; \mathbb{R})$  we have  $D(A, B) \leq \max\{D(A, \mathbb{R}), D(B, \mathbb{R})\}$ . Therefore, for every  $n \in \mathbb{Z}$

$$\begin{aligned} 2n/L &\leq D(F(n), F(-n)) \\ &\leq \max\{D(F(n), \mathbb{R}), D(F(-n), \mathbb{R})\} \\ &\leq \max\{D(F(n), F(0)), D(F(-n), F(0))\} + D(F(0), \mathbb{R}) \\ &\leq Ln + D(F(0), \mathbb{R}). \end{aligned}$$

Letting  $n \rightarrow \infty$ , we obtain  $L \geq \sqrt{2}$ .  $\square$

Despite the last statement of Theorem 5.1, many spaces of the form  $\mathcal{H}(X; C)$  do contain an isometric copy of  $l^\infty$ . The following is a partial result in this direction.

**PROPOSITION 5.2.** *Let  $X$  be a geodesic metric space. Suppose that  $C \in \text{CL}(X)$  contains a sequence  $\{c_n : n \geq 1\}$  such that  $\text{dist}(c_n, C \setminus \{c_n\}) \rightarrow \infty$  as  $n \rightarrow \infty$ . Then  $l^\infty$  isometrically embeds into  $\mathcal{H}(X; C)$ .*

**PROOF.** For each  $n \geq 1$  let  $d_n = \text{dist}(c_n, C \setminus \{c_n\})$  and choose a geodesic arc  $\gamma_n : [0, 1] \rightarrow X$  of length  $d_n/2$  so that  $\gamma(0) = c_n$ . For  $x = (x_1, x_2, \dots) \in l^\infty$  set  $x_{ij} = \min\{d_j/4, \max\{-d_j/4, x_i\}\}$ ,  $i, j \geq 1$ . The sequence

$$\tilde{x} := (x_{11}, x_{21}, x_{12}, x_{31}, x_{22}, x_{13}, \dots)$$

verifies  $|\tilde{x}_n| \leq d_n/4$ , and the map  $x \mapsto \tilde{x}$  is an isometry of  $l^\infty$  into itself.

Let  $A(x) = (C \setminus \{c_n : n \geq 1\}) \cup \{\gamma_n(2\tilde{x}_n/d_n + 1/2) : n \geq 1\}$ . It is clear that  $A(x) \in \mathcal{H}(\mathbb{R}; C)$  and  $D(A(x), A(y)) \leq \|\tilde{x} - \tilde{y}\|_{l^\infty}$  for all  $x, y \in l^\infty$ . To prove the reverse inequality, choose  $k$  so that  $|\tilde{x}_k - \tilde{y}_k| = \|\tilde{x} - \tilde{y}\|_{l^\infty}$  and observe that

$$\text{dist}(\gamma_n(2\tilde{x}_n/d_n + 1/2), A(y)) = |\tilde{x}_n - \tilde{y}_n|.$$

This completes the proof.  $\square$

For example, Proposition 5.2 implies that every separable metric space admits an isometric embedding into  $\mathcal{H}(\mathbb{R}, \{n^2 : n \in \mathbb{Z}\})$ .

### References

- [1] V. V. Aseev and A. A. Shalaginov, *Mappings that boundedly distort distance ratios*, Dokl. Akad. Nauk **335** (1994), no. 2, 133–134; English translation in Russian Acad. Sci. Dokl. Math. **49** (1994), no. 2, 248–250.
- [2] T. O. Banach and M. M. Zarichnyĭ, *Compactifications of the iterated hyperspace functor*, Izv. Vyssh. Uchebn. Zaved. Mat. (1987), no. 10, 3–6; English translation in Soviet Math. (Iz. VUZ) **31** (1987), no. 10, 1–4.
- [3] C. Bandt, *On the metric structure of hyperspaces with Hausdorff metric*, Math. Nachr. **129** (1986), 175–183.
- [4] S. Bates, W. B. Johnson, J. Lindenstrauss, D. Preiss, and G. Schechtman, *Affine approximation of Lipschitz functions and nonlinear quotients*, Geom. Funct. Anal. **9** (1999), 1092–1127.
- [5] C. J. Bishop and J. T. Tyson, *Locally minimal sets for conformal dimension*, Ann. Acad. Sci. Fenn. Math. **26** (2001), no. 2, 361–373.
- [6] M. Bonk and B. Kleiner, *Conformal dimension and Gromov hyperbolic groups with 2-sphere boundary*, Geom. Topol. **9** (2005), 219–246.
- [7] M. Bonk and O. Schramm, *Embeddings of Gromov hyperbolic spaces*, Geom. Funct. Anal. **10** (2000), no. 2, 266–306.
- [8] M. Bourdon, *Au bord de certains polyèdres hyperboliques*, Ann. Inst. Fourier (Grenoble), **45** (1995), 119–141.
- [9] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.
- [10] V. W. Bryant, *The convexity of the subset space of a metric space*, Compositio Math. **22** (1970), 383–385.
- [11] M. Csörnyei, *Can one squash the space into the plane without squashing?*, Geom. Funct. Anal. **11** (2001), 933–952.
- [12] T. Foertsch, *Isometries of spaces of convex compact subsets of globally non-positively Busemann curved spaces*, Colloq. Math. **103** (2005), no. 1, 71–84.
- [13] M. Ghamsari and D. A. Herron, *Higher dimensional Ahlfors regular sets and chordarc curves in  $\mathbb{R}^n$* , Rocky Mountain J. Math. **28** (1998), no. 1, 191–222.
- [14] M. Gromov, *Metric structures for Riemannian and non-Riemannian spaces*, with appendices by M. Katz, P. Pansu and S. Semmes. Progress in Mathematics, vol. 152. Birkhäuser Boston, Inc., Boston, MA, 1999.
- [15] J. Heinonen, *Lectures on analysis on metric spaces*, Springer-Verlag, New York-Heidelberg, 2001.
- [16] D. A. Herron and V. Mayer, *Bi-Lipschitz group actions and homogeneous Jordan curves*, Illinois J. Math. **43** (1999), no. 4, 770–792.
- [17] A. Hohti, *On Lipschitz homogeneity of the Hilbert cube*, Trans. Amer. Math. Soc., **291** (1985), no. 1, 75–86.
- [18] D. Ivaşcu, *Remarks on a class of quasimetric mappings*, in “Romanian-Finnish Seminar on Complex Analysis” 125–131, Lecture Notes in Math., vol. 743, Springer, Berlin, 1979.
- [19] W. B. Johnson, J. Lindenstrauss, D. Preiss, and G. Schechtman, *Uniform quotient mappings of the plane*, Michigan Math. J. **47** (2000), 15–31.
- [20] S. Keith and T. Laakso, *Conformal Assouad dimension and modulus*, Geom. Funct. Anal. **14** (2004), no. 6, 1278–1321.

- [21] L. V. Kovalev, *Conformal dimension does not assume values between 0 and 1*, to appear in Duke Math. J.
- [22] O. Maleva, *Lipschitz quotient mappings with good ratio of constants*, *Mathematika* **49** (2002), parts 1/2, No. 97/98, 159–165.
- [23] P. Mattila, *Geometry of sets and measures in Euclidean spaces*, Cambridge Univ. Press, Cambridge, 1995.
- [24] M. Mendel and A. Naor, *Euclidean quotients of finite metric spaces*, *Adv. in Math.* **189** (2004), no. 2, 451–494.
- [25] P. Pansu, *Dimension conforme et sphère à l'infini des variétés à courbure négative*, *Ann. Acad. Sci. Fenn. Ser. A I Math.* **14** (1989), no. 2, 177–212.
- [26] B. Randrianantoanina, *On the structure of level sets of uniform and Lipschitz quotient mappings from  $\mathbb{R}^n$  to  $\mathbb{R}$* , *Geom. Funct. Anal.* **13** (2003), 1329–1358.
- [27] J. Roe, *Lectures on coarse geometry*, University Lecture Series, vol. 31. Amer. Math. Soc., Providence, RI, 2003.
- [28] A. A. Shalaginov, *Mappings of self-similar curves*, *Sibirsk. Mat. Zh.* **34** (1993), no. 6, 210–215; English translation in *Siberian Math. J.* **34** (1993), no. 6, 1190–1195.
- [29] H. Toruńczyk and J. West, *A Hilbert space limit for the iterated hyperspace functor*, *Proc. Amer. Math. Soc.* **89** (1983), no. 2, 329–335.
- [30] J. T. Tyson, *Sets of minimal Hausdorff dimension for quasiconformal maps*, *Proc. Amer. Math. Soc.* **128** (2000), no. 11, 3361–3367.
- [31] J. T. Tyson, *Lowering the Assouad dimension by quasisymmetric mappings*, *Illinois J. Math.* **45** (2001), no. 2, 641–656.
- [32] J. T. Tyson, *Bi-Lipschitz embeddings of hyperspaces of compact sets*, *Fund. Math.*, to appear.
- [33] J. Väisälä, *Free quasiconformality in Banach spaces. I*, *Ann. Acad. Sci. Fenn. Ser. A I Math.* **15** (1990), no. 2, 355–379.

DEPARTMENT OF MATHEMATICS, MAILSTOP 3368, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368

*E-mail address:* lkovalev@math.tamu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 WEST GREEN STREET, URBANA, IL 61822

*E-mail address:* tyson@math.uiuc.edu