

# On the Number of Vertices with Specified Eccentricity

Dhruv Mubayi and Douglas B. West  
Department of Mathematics  
University Of Illinois  
Urbana, IL 61801

## Abstract

The *eccentricity* of a vertex  $v$  in a graph is the maximum of the distances from  $v$  to all other vertices. The *diameter* of a graph is the maximum of the eccentricities of its vertices. Fix the parameters  $n, d, c$ . Over all graphs with order  $n$  and diameter  $d$ , we determine the maximum (within 1) and the minimum of the number of vertices with eccentricity  $c$ .

We solve an extremal problem involving distances in connected graphs. Let  $G$  be a connected graph. The *distance*  $d(u, v)$  between two vertices  $u$  and  $v$  in  $G$  is the length of a shortest  $u, v$ -path. The *eccentricity* of a vertex  $v$  in  $G$  is  $\epsilon(v) = \max_{u \in V(G)} d(u, v)$ . The *radius* and *diameter* of  $G$  are the minimum and maximum eccentricities, respectively. A *diametric path* is a shortest  $u, v$ -path when  $d(u, v)$  is the diameter of  $G$ .

Let  $\mathcal{G}_{n,d}$  be the family of connected graphs with order  $n$  and diameter  $d$ . Over  $\mathcal{G}_{n,d}$ , let  $h_{n,d}(c)$  and  $f_{n,d}(c)$ , respectively, be the minimum and maximum number of vertices with eccentricity  $c$ . In this paper, we determine these functions almost completely. Since diameter is maximum eccentricity, these values are 0 if  $c > d$ . Also, no vertex  $w$  has eccentricity less than half  $\text{diam}G$ , since  $d(u, v) \leq d(u, w) + d(w, v)$  for all  $u, v, w$ . Furthermore, the existence of a diametric path requires at least  $d + 1$  vertices. We therefore may assume that  $d/2 \leq c \leq d < n$ . We also assume that  $d > 1$ , since otherwise  $G$  is a clique and all vertices have eccentricity  $d$ .

We consider only simple graphs. A *nontrivial* graph is a graph with at least one edge. Let  $N_G(v)$  or simply  $N(v)$  denote the set of neighbors of vertex  $v$  in graph  $G$ . A *vertex expansion* in a graph  $G$  is a replacement of a vertex  $v \in V(G)$  by a clique  $Q$  of new vertices, such that the neighborhood outside  $Q$  of each vertex of  $Q$  is  $N_G(v)$ . Let  $C_n$  and  $P_n$  denote the cycle and the path of order  $n$ .

## 1 Preliminary Lemmas

We begin with elementary remarks useful for both the minimization and the maximization problems. A simple class of graphs provides extremal examples in many cases.

**Lemma 1** *For  $k > 0$ , let  $G(k, 0) = C_{2k}$ . For  $k \geq l > 0$ , let  $G(k, l)$  be the graph obtained by extending a path of length  $l - 1$  from one vertex of  $C_{2k}$  and adding one pendant edge at the diametrically opposite vertex of  $C_{2k}$ . The graph  $G(k, l)$  has order  $2k + l$ , diameter  $k + l$ , and no vertices of eccentricity less than  $k$ . The count of vertices in  $G(k, l)$  with various eccentricities is*

case	eccentricity	multiplicity
$l > 1$	$k$	$2k - 2l + 2$
	$k < \epsilon(v) < k + l - 1$	3
	$k + l - 1 \leq \epsilon(v) \leq k + l$	2
$l = 1$	$k$	$2k - 1$
	$k + 1$	2
$l = 0$	$k$	$2k$

**Proof:** Vertices on the cycle have eccentricity at least  $k$ ; vertices on the pendant paths have eccentricity greater than  $k$ . From the peripheral vertices, eccentricity decreases by one with each step toward the center until eccentricity  $k$  is reached (see Fig. 1).  $\square$




Fig. 1.  $G(6, 4)$ , labeled by eccentricity

**Lemma 2** *A vertex expansion in a nontrivial graph does not change the eccentricity of any vertex, and the eccentricity of the new vertices equals the eccentricity of the vertex they replaced. In particular,  $h_{n+1,d}(c) \leq h_{n,d}(c)$  and  $f_{n+1,d}(c) \geq f_{n,d}(c) + 1$ .*

**Proof:** Suppose that  $G'$  is obtained from  $G$  by a vertex expansion at  $v$ . No chordless path of length more than one contains two vertices of the new clique  $Q$ , since those vertices have the same neighborhoods outside  $Q$ . Thus every shortest  $x, y$ -path in  $G'$  is a copy of an  $x, y$ -path in  $G$  (or of a path ending at  $v$  if  $y \in Q$ ). Thus distances and eccentricities do not change. The remark about  $h$  follows by expanding a vertex of eccentricity other than  $c$  in an extremal graph for the parameters  $n, d, c$ . For the remark about  $f$ , expand a vertex of eccentricity  $c$ .  $\square$

**Lemma 3** *Suppose that  $G \in \mathcal{G}_{n,d}$ . If  $n - d \leq c \leq d$  and  $c \geq d/2$ , then the vertices  $v_c$  and  $v_{d-c}$  on a diametric path  $P$  with vertices  $v_0, \dots, v_d$  have eccentricity  $c$ . If  $n < c + d$ , then  $\epsilon(v_i) < c$  for  $d - c < i < c$ .*

**Proof:** By symmetry, we need only consider  $v_i$ , where  $d/2 \leq i \leq c$ . Define  $c'$  by

$$c' = \begin{cases} c & \text{if } i = c \\ c - 1 & \text{if } i < c. \end{cases}$$

Every vertex of  $P$  is within distance  $c'$  of  $v_i$ , and when  $i = c$  we have  $d(v_i, v_0) = c'$ . If  $v_i$  does not have eccentricity at most  $c'$ , then there is a vertex  $w \notin V(P)$  such that  $d(v_i, w) = c' + 1$ . Let  $Q$  be a shortest  $w, v_i$ -path, and let  $v_q$  be the vertex at which  $Q$  first enters  $P$ . For  $q > c'$ , let  $R$  be a shortest  $w, v_0$ -path; when  $q \leq c'$ , let  $R$  be a shortest  $w, v_d$ -path. Let  $v_r$  be the vertex where  $R$  first enters  $P$ . If  $R$  contains  $v_{c'}$ , then the portion of  $R$  before  $v_{c'}$  is a  $w, v_{c'}$ -path, and hence the length of  $R$  is at least  $c' + 1 + c'$  or at least  $(c' + 1) + (d - c')$ . Since each of these exceeds  $d$ , we conclude that  $v_r$  and  $v_q$  are on opposite sides of  $v_{c'}$  along  $P$  (although  $q$  may equal  $c'$ ). Fig. 2 shows one such arrangement.




Fig. 2. Forcing eccentricity  $c$

Let  $z$  be the last vertex of  $Q \cap R$ . Because each is a shortest path from  $w$ , we may assume that  $Q$  and  $R$  coincide before  $z$ . We consider four distances along  $P, Q, R$ . Let  $\alpha = d(v_i, v_r), \beta = d(v_i, v_q), \gamma = d(v_r, z), \delta = d(v_q, z)$ . Because  $P$  is a shortest  $v_0, v_d$ -path, we have  $\gamma + \delta \geq \alpha + \beta$ . Because  $Q$  is a shortest  $v_i, w$ -path, we have  $\alpha + \gamma \geq \beta + \delta$ . Summing the two inequalities yields  $\gamma \geq \beta$ . Thus

$$n \geq |V(P \cup Q \cup R)| = (d + 1) + (d(v_i, w) - \beta) + (\gamma - 1) \geq d + 1 + c'.$$

When  $i = c = c'$ , this contradicts the hypothesis  $n \leq c + d$ . When  $i < c = c' + 1$ , this contradicts the hypothesis  $n < c + d$ .  $\square$

These tools, which will be useful when considering  $f_{n,d}$ , enable us to compute  $h_{n,d}$ .

**Theorem 4** For  $n > d$  and  $d/2 \leq c \leq d$ ,

$$h_{n,d}(c) = \begin{cases} 2 & \text{if } c = d \\ 0 & \text{if } n > c + d \text{ and } d/2 \leq c < d \\ 2 & \text{if } n \leq c + d \text{ and } d/2 < c < d \\ 1 & \text{if } n \leq c + d \text{ and } c = d/2. \end{cases}$$

**Proof:** The endpoints of a diametric path have eccentricity  $d$ , and  $P_{d+1}$  with Lemma 2 shows that this is the minimum when  $c = d$ . For  $d/2 \leq c < d$ , the graph  $G(c + 1, d - c - 1)$  of Lemma 1 has order  $c + d + 1$ , diameter  $d$ , and no vertices of eccentricity  $c$ . With Lemma 2, this completes the case  $n > c + d$ . The lower bound for the case  $n \leq c + d$  is given by Lemma 3. For the upper bound when  $n \leq c + d$  we use  $P_{d+1}$  and Lemma 2.  $\square$

## 2 The Maximization Problem

We now study the maximum number of vertices with eccentricity  $c$ . We begin with constructions for the lower bound.

**Lemma 5** For  $n > d$  and  $d/2 \leq c \leq d$ ,

$$f_{n,d}(c) \geq \begin{cases} n - 2(d - c) & \text{if } n \geq c + d \text{ and } c \geq d - 1 \\ n - 3(d - c) + 2 & \text{if } n \geq c + d \text{ and } \frac{2d-1}{3} \leq c < d - 1 \\ n - d + 1 & \text{if } n < c + d \text{ or } \frac{d}{2} < c \leq \frac{2d-1}{3} \\ n - d & \text{if } c = \frac{d}{2}. \end{cases}$$

**Proof:** Since these formulas increase by one with each augmentation of  $n$ , by Lemma 2 it suffices to consider the smallest value of  $n$  in each case. For Case 1 and Case 2, let  $n = c + d$  and let  $G = G(c, d - c)$ . By Lemma 1, the number of vertices in  $G$  with eccentricity  $c$  is  $2c - (d - c) = (c + d) - 2(d - c)$  when  $d - c \leq 1$  and is  $2c - 2(d - c) + 2 = (c + d) - 3(d - c) + 2$  when  $d - c > 1$ . For Case 3 and Case 4, let  $n = d + 1$  and let  $G = P_{d+1}$ . Here the number of vertices with each eccentricity is 2, except that there is only one central vertex when  $d$  is even.  $\square$

When  $n \geq c + d$  and  $c < (2d - 1)/3$ , the constructions of Cases 2 and 3 in Lemma 5 both apply, but the construction in Case 3 provides more vertices of eccentricity  $c$ .

To obtain upper bounds on  $f_{n,d}(c)$ , the formulas in Lemma 5 suggest that we obtain lower bounds on the number of vertices that must have eccentricity other than  $c$ , since the desired bound is independent of  $n$  in each case. Let  $g_{n,d}(c) = n - f_{n,d}(c)$ . Our approach is to start with a diametric path  $P = v_0, v_1, \dots, v_d$  in  $G \in \mathcal{G}_{n,d}$  and to deduce the existence of many vertices with eccentricity not  $c$ . The vertices of  $P$  before  $v_{d-c}$  and after  $v_c$  have eccentricity greater than  $c$ ; thus  $g_{n,d}(c) \geq 2(d - c)$ . This proves optimality of the constructions in Lemma 5 when  $c \geq d - 1$  and when  $c = d/2$ . For  $d/2 < c < d - 2$ , we must improve the lower bound on  $g_{n,d}(c)$ .

Although we will try to show that the bounds of Lemma 5 are optimal, in fact we can only prove that  $g_{n,d}(c) \geq 3(d - c) - 3$  when  $\frac{2d-2}{3} \leq c < d - 1$ . The cases in which we cannot show that  $g_{n,d}(c) \geq 3(d - c) - 2$  lead in fact to surprising examples where  $g_{n,d}(c) = 3(d - c) - 3$ .

**Theorem 6** For  $n > d$  and  $d/2 < c < d - 1$ ,

$$g_{n,d}(c) \geq \begin{cases} 3(d - c) - 2 & \text{if } n \geq c + d \text{ and } \frac{2d-2}{3} \leq c < d - 1 \text{ and } d - c \text{ is even} \\ 3(d - c) - 3 & \text{if } n \geq c + d \text{ and } \frac{2d-2}{3} \leq c < d - 1 \text{ and } d - c \text{ is odd} \\ d - 1 & \text{if } n < c + d \text{ or } \frac{d}{2} < c \leq \frac{2d-1}{3}. \end{cases}$$

Furthermore, if  $g_{n,d}(c) = 3(d - c) - 3$ , then a graph achieving this has a diametric pair at the end of paths of length  $\frac{d-c+1}{2}$  that emerge from antipodal vertices on a cycle of length  $2c - 2$ .

**Proof:** Suppose that  $n > d$  and  $d/2 < c < d - 1$ , and consider  $G \in \mathcal{G}_{n,d}$ . Let  $P = v_0, \dots, v_d$  be a diametric path in  $G$ . For  $n < c + d$ , Lemma 3 implies that  $\epsilon(v_i) \neq c$  when  $i \notin \{d - c, c\}$ . Thus we may assume that  $n \geq c + d$ .

In this range, the desired lower bound is  $\min\{3(d - c) - 2, d - 1\}$ , so it suffices to obtain either  $3(d - c) - 2$  or  $d - 1$  as a lower bound. If no vertex of  $P$  between  $v_{d-c}$  and  $v_c$  has eccentricity  $c$ , then at least  $d - 1$  vertices of  $G$  have eccentricity other than  $c$ . Thus we may assume that  $P$  has a vertex of eccentricity  $c$  between  $v_{d-c}$  and  $v_c$ .

Let  $v_i$  be the last vertex on  $P$  before  $v_c$  that has eccentricity  $c$ , and let  $z$  be a vertex at distance  $c$  from  $v_i$ . Since  $d(v_i, u) < c$  for  $u \in V(P)$ , we have  $z \notin V(P)$ . Let  $Q$  be a shortest  $z, v_d$ -path, and let  $v_t$  be the first vertex of  $Q$  that is also on  $P$ . Since  $d(z, v_d) \leq d$  and  $d(z, v_i) = c$ , we have  $t > i$ .

We define several sets of vertices that will have eccentricity other than  $c$ .

$$\begin{aligned} A &= \{v_j: i < j < c\} \\ B &= \{u \in V(Q - P): d(u, v_d) < d - c\} \\ C &= \{u \in V(Q - P): d(u, v_d) > c\} \end{aligned}$$

For  $u \in A$ , the choice of  $i$  yields  $\epsilon(v_j) \neq c$ . By the triangle inequality, every vertex of  $G$  having distance less than  $d - c$  from one end of  $P$  has distance more than  $c$  from the other end; thus  $\epsilon(u) > c$  for  $u \in B$ . For  $u \in C$ , the definition of  $C$  yields  $\epsilon(u) \geq d(u, v_d) > c$ . Thus  $\epsilon(u) \neq c$  for  $u \in A \cup B \cup C$ . Note also, since  $c > d - c$ , that  $A, B, C$  are pairwise disjoint.

Define  $v_{d-i'}, z', Q', v_{d-t'}, A', B', C'$  symmetrically from the other end of  $P$ . As above,  $\epsilon(u) \neq c$  for  $u \in A' \cup B' \cup C'$ , and  $A', B', C'$  are pairwise disjoint. Fig. 3 shows a case with  $t, t' > c$  and  $(A' \cup B' \cup C') \cap (A \cup B \cup C) = \emptyset$ . By construction,  $i \geq d - i'$ . Since  $d(u, v_d) < d - i$  for  $u \in A \cup B$  and  $d(u, v_0) < d - i'$  for  $u \in A' \cup B'$  a vertex common to these two sets would yield a  $v_0, v_d$ -path shorter than  $P$ . Thus we may assume that  $(A \cup B) \cap (A' \cup B') = \emptyset$ .




Fig. 3. Sets of vertices with eccentricity not  $c$

To obtain the desired lower bound on  $g_{n,d}(c)$ , we will show that  $\gamma = |A \cup B \cup C \cup A' \cup B' \cup C'|$  is large. In addition to the  $2(d - c)$  outermost vertices of  $P$ , it suffices to show that  $\gamma \geq d - c - 2$  or that  $\gamma \geq 2c - d - 1$ .

Let  $k = d(z, v_d) - c$ . Since  $Q$  is a shortest  $z, v_d$ -path, we have  $d(v_t, z) = d(z, v_d) - (d - t)$ . By the triangle inequality,  $d(v_t, z) \geq d(v_i, z) - (t - i)$ . Thus  $(c + k) - (d - t) \geq c - (t - i)$ , which yields  $k \geq d - 2t + i$ . If  $t \leq c$ , then  $k \geq (d - c) + (i - c)$ , and we obtain  $|A \cup C| \geq c - i - 1 + k \geq d - c - 1$ .

Thus we may assume that  $t > c$ , which yields  $|B| = \min\{t - c - 1, c + k - (d - t)\}$ . Similarly, we define  $k' = d(z', v_0) - c$  and may assume that  $t' > c$ . Fig. 3 illustrates the case where  $k, k' > 0$  and the six relevant sets are pairwise disjoint. For all cases, we have computed the inequalities  $2t \geq d - k + i$  and  $2t' \geq d - k' + i'$ . Invoking these and then  $i, i' \leq c - 1$  yields the computations below and the analogous lower bounds on  $|A' \cup B' \cup C'|$  in terms of  $k'$ .

range	$ A $	$ B $	$ C $	$ A \cup B \cup C  \geq$
$k \geq 0$	$c - i - 1$	$t - c - 1$	$k$	$\frac{d - c - 3 + k}{2}$
$0 > k \geq d - 2c - 1$	$c - i - 1$	$t - c - 1$	0	$\frac{d - c - 3 - k}{2}$
$d - 2c - 1 > k \geq 1 - c$	$c - i - 1$	$c + k - d + t$	0	$\frac{3c - d + k - 1}{2}$

With  $k \geq 1 - c$ , the last bound becomes  $(2c - d)/2$ . Each of these bounds is at least half the needed amount, except when  $k = 0$ . We may sum the bounds for  $|A \cup B \cup C|$  and  $|A' \cup B' \cup C'|$  unless these two sets intersect. When  $C \cap B' = C' \cap B = \emptyset$ , summing the two lower bounds and

subtracting  $|C \cap C'|$  yields at least the desired amount unless  $C = C'$ , in which case it yields  $\gamma \geq d - c - 3$ . We postpone the analysis of that case.

**Case 1:**  $C \cap B'$  or  $C' \cap B$  is nonempty. By symmetry, we may assume  $C \cap B' \neq \emptyset$ . In following  $Q'$  from  $v_0$ , let  $y$  be the first vertex encountered in  $C$ . Let  $H$  be the cycle formed by the  $v_t, v_{d-t'}$ -subpath of  $P$ , the  $v_{d-t'}, y$ -subpath of  $Q'$ , and  $y, v_t$ -subpath of  $Q$ . We define several vertex sets with respect to  $H$  (see Fig. 4). Let  $D = \{u \in V(H) - V(P) : d(u, v_0) < d - c\}$ . Since  $d(u, v_d) \leq c$  for  $u \in V(Q) - V(C)$ , we have  $D \subseteq B' \cup C$ . Also  $|D| = t' - c - 1$ . Let  $C_0 = C - V(H)$ , and let  $C_1 = C - D - C_0$ .

The cycle  $H$  consists of  $t - (d - t')$  edges on  $P$ ,  $t' - c - 1$  edges from  $v_{d-t'}$  through  $D$ ,  $|C_1|$  edges to absorb the vertices of  $C_1$ ,  $c + 1 - (d - t)$  edges on  $Q$  from  $C$  to  $v_t$ . Thus  $H$  has length  $2(t + t' - d) + |C_1|$ . The distance from  $z$  to  $v_i$  is at most  $|C_0|$  plus half the length of  $H$ ; thus  $c \leq |C_0| + |C_1|/2 + t + t' - d$ . Now we compute

$$|B \cup B' \cup C| \geq |B \cup D \cup C_0| + |C_1|/2 \geq (t - c - 1) + (t' - c - 1) + (c - t - t' + d) = d - c - 2.$$




Fig. 4. Case 1:  $C \cap B' \neq \emptyset$

**Case 2:**  $B \cap C' = B' \cap C = \emptyset$  and  $C = C'$ . If  $\gamma = d - c - 3$ , then the inequalities used in obtaining  $\gamma \geq d - c - 3$  hold with equality, and we need only one more vertex with eccentricity other than  $c$ . The equalities yield  $i = i' = c - 1$ ,  $k = k' \geq 0$ , and  $t = t' = (d - k + c - 1)/2$ . Thus  $t - i = d - t - k$ , which implies that the  $v_i, z$ -path along  $P$  and  $Q$  is a shortest  $v_i, z$ -path (similarly for  $v_{d-i'}$  and  $z'$ ). Between  $B$  and  $C$  there are  $1 + 2c - d$  vertices on  $Q$ .

Let  $x$  be the first vertex of  $Q$  after  $B$  (see Fig. 4); we have  $d(x, v_d) = d - c$ . By the triangle inequality,  $d(x, v_0) \geq c$ . We are finished if  $\epsilon(x) > c$ , so we may assume that  $d(x, v_0) = c$ . A shortest  $x, v_0$ -path and the  $x, v_d$ -portion of  $Q$  together form a  $v_0, v_d$ -path  $P^*$  of length  $d$ .

We show next that  $d(x, u) < c$  for  $u \in B \cup C \cup B'$  ( $A, A'$  are empty and  $C' = C$ ). Using  $Q$  and  $P$ , we have  $d(x, u) < c$  for  $u \in B \cup C \cup \{v_c, \dots, v_{d-1}\}$ . Because  $\epsilon(u) > c$  when  $d(u, v_0) < d - c$ , we may assume that  $P^*$  enters the set of vertices within distance  $d - c - 1$  of  $v_0$  at  $v_{d-c-1}$  or at the most distant vertex of  $B'$ . Since  $d - t' - k' = (d - i') - (d - t')$ , we can modify the last part of the  $x, v_0$ -portion of  $P^*$  to obtain  $d(x, u) < c$  for  $u \in B' \cup \{v_1, \dots, v_{d-c}\}$ .

We may apply the same analysis to  $P^*$  that we applied to  $P$ ; we are finished unless this analysis also falls into Case 2. Let  $k^*$  and  $z^*$  be the value and vertex in the analysis of  $P^*$  that play the role of  $k$  and  $z$  in the analysis of  $P$ . If  $k^* > 0$ , then  $\epsilon(z^*) > c$ . Since  $d(x, z^*) = c$ , we have shown that  $z^*$  is not in  $B \cup C \cup B'$  and not on  $P$  within distance  $d - c - 1$  of either end. Hence  $z^*$  is a new vertex of large eccentricity and we are finished. Thus we may assume that  $k^* = 0$ .

Since we could have started with  $P^*$  as  $P$ , we may also assume that  $k = 0$ . Then when we generate the new  $P^*$ , again we are finished unless  $k^* = 0$ . The unresolved case is the case in which the theorem statement claims only  $g_{n,d}(c) \geq 3(d-c) - 3$ .  $\square$

The remaining unresolved case is when  $n \geq c + d$ ,  $\frac{2d-2}{3} \leq c < d - 1$ , and  $d - c$  is odd. In this situation we have proved that  $3(d-c) - 3 \leq g_{n,d}(c) \leq 3(d-c) - 2$ . Surprisingly, more subtle constructions achieve the lower bound for some such values of  $(d, c)$  when  $n$  is sufficiently large.

**Theorem 7** *If  $n \geq 3d + c - 3$ ,  $\frac{3d}{4} \leq c < d - 1$ , and  $d - c$  is odd, then  $g_{n,d}(c) = 3(d-c) - 3$ .*

**Proof:** For such values we construct a graph  $G$  illustrated in Fig. 5. Each segment in the illustration represents a path of indicated length. Let  $s = (d - c - 1)/2$ ; this is an integer since  $d - c$  is odd. To form  $G$ , begin with a cycle  $C$  of length  $2c - 2$  and two paths of length  $s + 1$  extending from antipodal vertices on  $C$  to the unique vertices of eccentricity  $d$ . In the language we have been using, we may label the vertices of the diametric path around the top of  $C$  as the path  $P$  with vertices  $v_0, \dots, v_d$ . Here  $v_{d-t}$  is the entrance of  $P$  to  $C$  and  $v_t$  is its departure. We have seen that  $g_{n,d}(c) < 3(d-c) - 2$  requires  $t = (d + c - 1)/2$ . Let  $P'$  be the other  $v_0, v_d$ -path of length  $d$ , with vertices  $v'_0, \dots, v'_d$ , of which  $2s + 4$  are shared with  $P$ .

We next add two paths  $W, W'$  of length  $2\lfloor c/2 \rfloor + 3$ . Listed by vertices, these are  $W = v_{d-c}, w_1, \dots, w_{2\lfloor c/2 \rfloor + 2}, v'_c$  and  $W' = v'_{d-c}, w'_1, \dots, w'_{2\lfloor c/2 \rfloor + 2}, v_c$ . Note that since  $2s + 1 = d - c$ , the endpoints of  $W$  and  $W'$  have distance  $s$  from  $v_{d-t}$  and  $v_t$  along  $P$  and  $P'$ . Note also that  $W$  and  $W'$  have no common vertex. We next add a path  $X$  from  $w_{\lfloor c/2 \rfloor - 2s + 1}$  to  $w'_{\lfloor c/2 \rfloor + 2s + 2}$  and a path  $X'$  from  $w'_{\lfloor c/2 \rfloor - 2s + 1}$  to  $w_{\lfloor c/2 \rfloor + 2s + 2}$ . The length of  $X$  and  $X'$  is  $2s - 1$  if  $c$  is even;  $2s$  if  $c$  is odd. Finally, we add the four edges  $\{v_{d-c+1}w_1, v'_{d-c+1}w'_1, v_{c-1}w'_{2\lfloor c/2 \rfloor + 2}, v'_{c-1}w_{2\lfloor c/2 \rfloor + 2}\}$ .




Fig. 5. A general construction for  $g_{n,d}(c) = 3(d-c) - 3$

Altogether, we have created  $3d + c - 3$  vertices. By Lemma 2, the construction is valid for  $n \geq 3d + c - 3$ . Vertices  $v_0$  and  $v_d$  have eccentricity  $d$ . All of  $v_j$  and  $v'_j$  with  $j < d - c$  or  $j > c$  have eccentricity greater than  $c$ ; there are  $2(s + 1) + 2(2s - 1) = 6s = 3(d - c) - 3$  such vertices. It remains to be shown that all other vertices have eccentricity exactly  $c$ . We use the four-fold symmetry of the construction to shorten the analysis.

We first show that all distances involving the remaining vertices are at most  $c$ . A convenient computation to avoid most discussion by parity is that  $2\lfloor c/2 \rfloor$  plus the length of  $X$  or  $X'$  equals  $c + 2s - 1$ . Also,  $c \leq 3d/4$  yields  $3s + 2 \leq \lceil c/2 \rceil$ , using  $s = (d - c - 1)/2$  and the opposite parity of  $d$  and  $c$ .

First consider  $G' = G - \{v_0, \dots, v_{d-t-1}\} - \{v_{t+1}, \dots, v_d\}$ . Within  $G'$ , we define a collection of cycles such that every two vertices appear together in at least one of the cycles and each cycle has length at most  $2c + 1$ . The cycle  $C$  has length  $2c - 2$ . The path from  $v_{d-c}$  to  $v_c$  via  $W, X, W'$  has length  $c - 2s + 1$ , which exceeds the distance along  $C$  by 2. Modifying  $C$  by replacing the shorter with the longer path yields a cycle  $C'$  of length  $2c$ . Combining the two paths yields a cycle  $C''$  of length  $2c - 4s$ . The cycle  $A$  consisting of  $X \cup X'$  plus the central portions of  $W$  and  $W'$  has length at most  $12s + 2$ , which is less than  $2c$ . Let  $B$  be the cycle consisting of  $X$ , the central portion of  $W'$ , the initial portions of  $W$  and  $W'$ , and the portion of  $C$  between  $v_{d-c}$  and  $v'_{d-c}$  via  $v_{d-t}$ . The length of  $B$  is  $c + 4s + 2$ , which is less than  $2c$ . Finally, let  $D$  be the cycle formed by  $W - v'_c$ , the edge from  $W - v'_c$  to  $v'_{c-1}$ , and the portions of  $P'$  and  $P$  meeting at  $v_{d-t}$ . The length of  $D$  is  $2c$  when  $c$  is odd and  $2c + 1$  when  $c$  is even. Together with all the cycles isomorphic to these by symmetry, we have the desired list of cycles in  $G'$ .

Now consider vertices in  $P \cap P'$ . It suffices by symmetry to specify paths of length at most  $c$  from  $v_d$  to  $W \cup X \cup (P - \{v_0, \dots, v_{d-c-1}\})$ . We reach those on  $P$  since the length of  $P$  is  $d$ . We reach  $w_1$  within distance  $c$  by traveling along  $P$  until the last step. From  $v_d$ , the path along  $P$  to  $W'$ , along  $W'$  to  $X$ , along  $X$  to  $W$ , and along  $W$  to  $w_2$  has length exactly  $c$ . If instead we travel in the other direction on  $W$  for the last segment, we can reach  $s$  of those vertices within distance  $c$  because  $3s + 2 \leq \lceil c/2 \rceil$  yields  $s \leq \lfloor c/2 \rfloor - 2s - 1$ . We reach the remainder of  $W$  from  $v_d$  along  $P'$  and  $W$ ; the length is at most  $\lfloor c/2 \rfloor + 3s + 2$ , which is at most  $c$ .

It now remains only to exhibit, for each  $u \in V(G')$ , a vertex  $f(u) \in V(G)$  such that  $d_G(u, f(u)) = c$ . By symmetry, we need only consider (roughly) half the vertices in each of  $X, W, P'$ . Let  $f(v_{d-c}) = f(w_1) = v_d$ ; these distances we have computed. For  $2 \leq j \leq 1 + \lfloor c/2 \rfloor$ , let  $f(w_j) = v'_{c-j+1}$ , except that when  $c$  is even and  $j = 1 + c/2$ , let  $f(w_j) = v'_{c-j+2}$ .

Observe that  $w_j$  and  $f(w_j)$  lie on the cycle  $D$ . Except for the exception, the distance between them on  $D$  via  $v'_{d-c}$  is  $j + 2s + (c - j + 1) - (d - c) = c$ . When  $j = 1 + \lfloor c/2 \rfloor$ , the path from  $w_j$  to  $v'_{c-j+1}$  through the junctions  $w_{\lfloor c/2 \rfloor + 2s + 2s}$ ,  $w'_{\lfloor c/2 \rfloor - 2s + 1}$ ,  $w'_1$ , and  $v'_{d-c+1}$  also has length  $c$  if  $c$  is odd, but it is a shortcut of length  $c - 1$  if  $c$  is even. In this case, setting  $f(w_j)$  to  $v'_{c-j+2}$  brings the length of the "shortcut" back to  $c$ . This distance computed along the *other half* of  $D$  is now also  $c$ . By checking the possible alternative routes, it can be verified that no other shortcuts arise. For example, when  $j \leq \lfloor c/2 \rfloor$ , one may try the route along  $X, W', v_c, v'_c$ ; the length is again  $c$ . One may try the route along  $w_1, v_{d-c+1}, v_c, v'_c$ ; the length is  $c - 3 + 2j$ , which exceeds  $c$  for  $j \geq 2$  and explains our choice of  $f(w_1)$ .

Since distance is symmetric, we may assign  $f(f(w_j)) = w_j$ . Since  $c - \lfloor c/2 \rfloor + 1 \leq d/2$ , we have thus also established eccentricity  $c$  for a vertex of each isomorphism class on  $P'$ . It remains only to consider  $X$ . Let  $x_j$  be the vertex of  $X$  whose distance from the junction  $w_{\lfloor c/2 \rfloor - 2s + 1}$  is the

same as that of  $w_j$ , where  $j > \lfloor c/2 \rfloor - 2s$ . It suffices to consider  $\lfloor c/2 \rfloor - 2s < j \leq \lfloor c/2 \rfloor - s$ , by symmetry. Let  $f(x_j) = f(w_j) = v'_{c-j+1}$ . Observe that  $v'_{c-j+1}$  is the vertex diametrically opposite  $x_j$  on the cycle  $C'$  of length  $2c$ . There are no shortcuts.  $\square$

We make no attempt to show that  $n = 3d + c - 3$  is the minimum number of vertices that permits such a construction. When  $d - c = 3$ , the constraint  $c \geq 3d/4$  requires  $d \geq 12$ . Since all cases with  $c \leq 2d/3$  are settled, this leaves  $(d, c) = (10, 7)$  and  $(d, c) = (11, 8)$ . In Figure 6 we illustrate special constructions for these cases. These constructions are not unique. Another small family of constructions with  $f_{n,d}(c) = n - 3(d - c) + 3$  resolves the open cases for  $d - c = 5$  and  $16 \leq d \leq 19$  (also for  $d - c = 3$  using a few more vertices than Figure 6). The family also has constructions for  $(d, c) = (22, 15)$  and  $(d, c) = (23, 16)$ ; in Figure 7 we show its largest member. Except for  $(22, 15)$  and  $(23, 16)$ , the gap of 1 remains between the upper bound and construction when  $2d/3 < c < 3d/4$  and  $d - c$  is odd and  $d - c \geq 7$ .




Fig. 6.  $g_{n,10}(7) = 6$  for  $n \geq 30$ , and  $g_{n,11}(8) = 6$  for  $n \geq 35$ .




Fig. 7.  $g_{n,23}(16) = 18$  for  $n \geq 96$ .